



# Mining the Flavoproteome of *Brucella ovis*, the Brucellosis Causing Agent in *Ovis aries*

 Martha Minjárez-Sáenz,<sup>a,b</sup>  Marta Martínez-Júlvez,<sup>a,b,d</sup>  Inmaculada Yruea,<sup>c,d</sup>  Milagros Medina<sup>a,b,d</sup>

<sup>a</sup>Departamento de Bioquímica y Biología Molecular y Celular, Facultad de Ciencias, Universidad de Zaragoza, Zaragoza, Spain

<sup>b</sup>Instituto de Biocomputación y Física de Sistemas Complejos (BIFI), Universidad de Zaragoza, Zaragoza, Spain

<sup>c</sup>Estación Experimental de Aula Dei, CSIC, Zaragoza, Spain

<sup>d</sup>Group of Biochemistry, Biophysics and Computational Biology "GBsC" (BIFI, Unizar) Joint Unit to CSIC, Zaragoza, Spain

**ABSTRACT** Flavoproteins are a diverse class of proteins that are mostly enzymes and contain as cofactors flavin mononucleotide (FMN) and/or flavin adenine dinucleotide (FAD), which enable them to participate in a wide range of physiological reactions. We have compiled 78 potential proteins building the flavoproteome of *Brucella ovis* (*B. ovis*), the causative agent of ovine brucellosis. The curated list of flavoproteins here reported is based on (i) the analysis of sequence, structure and function of homologous proteins, and their classification according to their structural domains, clans, and expected enzymatic functions; (ii) the constructed phylogenetic trees of enzyme functional classes using 19 *Brucella* strains and 26 pathogenic and/or biotechnological relevant alphaproteobacteria together with *B. ovis*; and (iii) the evaluation of the genetic context for each entry. Candidates account for ~2.7% of the *B. ovis* proteome, and 75% of them use FAD as cofactor. Only 55% of these flavoproteins belong to the core proteome of *Brucella* and contribute to *B. ovis* processes involved in maintenance activities, survival and response to stress, virulence, and/or infectivity. Several of the predicted flavoproteins are highly divergent in *Brucella* genus from revised proteins and for them it is difficult to envisage a clear function. This might indicate modified catalytic activities or even divergent processes and mechanisms still not identified. We have also detected the lack of some functional flavoenzymes in *B. ovis*, which might contribute to it being nonzoonotic. Finally, potentiality of *B. ovis* flavoproteome as the source of antimicrobial targets or biocatalyst is discussed.

**IMPORTANCE** Some microorganisms depend heavily on flavin-dependent activities, but others maintain them at a minimum. Knowledge about flavoprotein content and functions in different microorganisms will help to identify their metabolic requirements, as well as to benefit either industry or health. Currently, most flavoproteins from the sheep pathogen *Brucella ovis* are only automatically annotated in databases, and only two have been experimentally studied. Indeed, certain homologues with unknown function are not characterized, and they might relate to still not identified mechanisms or processes. Our research has identified 78 members that comprise its flavoproteome, 76 of them flavoenzymes, which mainly relate to bacteria survival, virulence, and/or infectivity. The list of flavoproteins here presented allows us to better understand the peculiarities of *Brucella ovis* and can be applied as a tool to search for candidates as new biocatalyst or antimicrobial targets.

**KEYWORDS** *Brucella ovis*, flavoproteome, alpha-proteobacteria, flavoenzymes, metabolic function

Flavoproteins contain as cofactors the riboflavin (RF) derivatives flavin mononucleotide (FMN) and/or flavin adenine dinucleotide (FAD), and participate in a wide range of physiological reactions and metabolic pathways (1–4). Most of them are essential

**Editor** Francisco Uzal, University of California, Davis

**Copyright** © 2022 Minjárez-Sáenz et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Milagros Medina, [mmedina@unizar.es](mailto:mmedina@unizar.es), or Marta Martínez-Júlvez, [mmartine@unizar.es](mailto:mmartine@unizar.es).

The authors declare no conflict of interest.

**Received** 17 November 2021

**Accepted** 19 February 2022

**Published** 22 March 2022

mediators in oxido-reduction processes, where they can either exchange one or two electrons, whereas other cofactors or coenzymes exclusively catalyze one- (iron-sulfur clusters, haem groups, etc.) or two- (nicotinamide adenine nucleotides) electron transfer processes. This makes flavoenzymes exhibit their redox versatility in a large number of metabolic redox processes. Moreover, among the around 500 different flavin-dependent proteins so far identified ~10% catalyze nonredox reactions or act as signaling and sensing molecules (5–7). Examples would include signal transduction in programmed cell death, embryonic development, chromatin remodeling, nucleotide synthesis, tRNA methylation, protein folding, and defense against oxidative stress (8–15), among others. Some flavoproteins are also involved in the xenobiotic metabolism of aromatic compounds, in pathogens virulence, and in light-dependent processes in which flavin receives photons: as luciferase light-emission, DNA reparation, plant phototropism, and cellular clocks (16–22). Many flavoenzymes are also suitable biocatalysts due to their selectivity, control, and efficiency of the reactions they catalyze (3, 23), as well as therapeutic targets both in the treatment of infectious diseases and in mammalian pathological situations (24–32). Despite their potential, only a few flavoproteins are widely exploited (33). To expand their applicability, it is essential to investigate the flavoprotein content and diversity in different organisms. So far, detailed information of flavoproteomes has only been reported in *Homo sapiens*, *Saccharomyces cerevisiae*, and *Arabidopsis thaliana* (5, 6, 34, 35), to which has to be added a comparative analysis in a bunch of archaeal, eubacterial, protozoan, and eukaryotic genomes (7). These studies indicate that whereas some organisms depend heavily on flavin-dependent activities, others maintain a minimum of flavoproteins. A better knowledge about content of flavoproteomes would surely help in the understanding of metabolic requirements in different organisms, and benefit either industry or health.

In this context, we aimed to trace flavin-binding proteins in *Brucella ovis* (*B. ovis*), a Gram-negative bacteria that provokes placentitis in sheep and genital lesions in rams that affect the quality of the semen and the fertility, and causes major economic impacts in countries and regions with sheep (*Ovis aries*) breeding activity (36). The virulence of each particular *Brucella* species depends on enzymes and cell envelope proteins that act as virulence factors, and on the ability to fight against the host response (37). Nonetheless, which genes and proteins are essential in these processes, as well as how they interact during intracellular virulence, still remain unclear. Here, we have identified 78 candidates to constitute the *B. ovis* flavoproteome. We envisage a metabolic function for many of them upon evaluation of their presence in *Brucella* and pathogenic and biotechnological relevant alphaproteobacteria (Table SP1 in the supplemental material), as well as of their evolutionary fingerprint and genetic context. Our studies point to a list of flavoproteins with high probability to contribute to the *B. ovis* survival, virulence, and/or infectivity, some of which have not yet been characterized in any homologue. This list is also discussed as a tool in the search for candidates as new biocatalyst or antimicrobial targets.

## RESULTS

**Overall features of the *Brucella ovis* flavoproteome.** The *Brucella ovis* ATCC 25840 genome contains 2,890 genes organized in two chromosomes, CI (2.10 Mb, 1,928 genes) and CII (1.15 Mb, 962 genes) (18). Among them, we identified 78 flavoproteins encoded by 49 genes of CI (2.4%) and 37 genes of CII (3.5%). They constitute the curated flavoproteome of *B. ovis* (Tables 1 and 2) and represent ~2.7% of the *B. ovis* proteome. This percentage agrees with average estimations from the study of other flavoproteomes (5–7, 34).

Fifty (64%) and 22 (28%) of these flavoproteins are predicted to bind, respectively, FAD and FMN as cofactor (Tables 1 and 2). Four (5%) would bind both: the NADH dependent dihydropyrimidine dehydrogenase, binding FMN and FAD respectively at its PreA and PreT subunits; the sarcosine oxidase beta subunit; the glutamate synthase, binding FMN and FAD respectively at the large and short subunits; and the assimilatory

**TABLE 1** Predicted flavoproteins encoded in the chromosome I of *Brucella ovis* ATCC 25840

Entry	EC	Protein	Pfam domain <sup>a</sup>	Locus tag	Protein code	Flavin	Coenzyme or ligand	More similar PDB (% identity)
1	1.1.5.3	Glycerol-3-phosphate dehydrogenase	DAO (PF01266) 19-347 DAO_C (PF16901) 399-508	BOV_RS00950 (glpD)	ABQ60174.1	FAD	Quinone	2QCU (50)
2	1.1.99.1	Choline dehydrogenase (Glucose-methanol-choline GMC family)	GMC_oxred_N (PF00732) 4-292 GMC_oxred_C (PF05199) 384-521	BOV_RS02765 (betA)	ABQ61350.1	FAD		2JBV (35)
3	1.1.99.1	Choline dehydrogenase (GMC family)	GMC_oxred_N (PF00732) 4-295 GMC_oxred_C (PF05199) 388-524	BOV_RS07775	ABQ60630.1	FAD		4HA6 (31)
4	1.1.-.- <sup>b</sup>	<b>Potential FAD-binding oxygen oxidoreductase (glcE)?<sup>c</sup></b>	FAD_binding_4 (PF01565) 54-187 FAD-oxidase_C (PF02913) 223-463	BOV_RS06750	ABQ60928.1	FAD		3PM9 (29)
5	1.1.-.- <sup>b</sup>	<b>Potential FAD-binding oxygen oxidoreductase (glcE)?</b>	FAD_binding_4 (PF01565) 43-180 FAD-oxidase_C (PF02913) 219-469	BOV_RS02095	ABQ61939.1	FAD		3PM9 (60)
6	1.3.1.1	NADH dependent Dihydropyrimidine dehydrogenase subunit PreA	DHO_dh (PF01180) 4-307 Fer4_21 (PF14697) 340-400	BOV_RS01510 (preA)	ABQ60560.1	FMN	2x(4Fe-4S) 4Fe-4S	2B4G (24) 1GTE (37)
7	1.3.1.88	NADH dependent Dihydropyrimidine dehydrogenase subunit PreT	Fer4_20 (PF14691) 31-139 Pyr_redox_2 (PF07992) 153-441	BOV_RS01515 (preT)	ABQ61103.1	FAD	NAD(P)H	5JCA_L (32) 5VJ7_A (35)
8	1.3.1.91	tRNA dihydrouridine synthase B	Dus (PF01207) 22-316	BOV_RS05355 (DusB)	ABQ61416.1	FMN	NAD(P)H	6EI9 (41)
9	1.3.1.98	tRNA dihydrouridine 20/20a synthase UDP-N-acetylmuramate dehydrogenase	Dus (PF01207) 21-320 FAD_binding_4 (PF01565) 42-172 murB_C (PF02873) 206-304	BOV_RS04255 (dusA) BOV_RS06850 (MurB)	ABQ61966.1 ABQ61769.1	FMN FAD	NAD(P)H NADPH	3B0P (46) 3TX1 (35)
10	1.3.5.1	Succinate dehydrogenase flavoprotein subunit	FAD_binding_2 (PF00890) 24-419 Succ_DH_flav_C (PF02910) 474-613	BOV_RS08985 (sdhA)	ABQ61077.1	FAD		2H88_A (63)
11	1.3.5.2	PyrD dihydroorotate dehydrogenase 2 (quinone)	DHO_dh (PF01180) 44-336	BOV_RS01655 (pyrD)	ABQ61413.1	FMN	Quinone	4ORI (48)
12	1.3.8.1	Short Chain Acyl-CoA dehydrogenase	Acyl-CoA_dh_N (PF02771) 37-155 Acyl-CoA_dh_M (PF02770) 160-268 Acyl-CoA_dh_1 (PF00441) 288-457 Acyl-CoA_dh_C (PF12806) 470-585	BOV_RS02120	ABQ60180.1	FAD		1BUC (29)
13	1.3.8.4	Isovaleryl-CoA dehydrogenase	Acyl-CoA_dh_N (PF02771) 7-118 Acyl-CoA_dh_M (PF02770) 122-217 Acyl-CoA_dh_1 (PF00441) 229-377	BOV_RS00090 (ivd)	ABQ60382.1	FAD		4KTO (83) 4OSM (100)
14	1.3.8.- <sup>b</sup>	Acyl-CoA dehydrogenase	Acyl-CoA_dh_N (PF02771) 12-123 Acyl-CoA_dh_M (PF02770) 127-221 Acyl-CoA_dh_1 (PF00441) 235-382	BOV_RS06310	ABQ61585.1	FAD		1RX0 (57)
15	1.4.3.5	Pyridoxamine 5'-phosphate oxidase	Putative_PNPOx (PF01243) 29-113 PNP_phzG_C (PF10590) 166-208 DAO (PF01266) 3-313 HI0933_like (PF03486) 5-391	BOV_RS02140 (pdxH)	ABQ60142.1	FMN		1NDL (45)
16	1.4.3.19	Glycine oxidase ThiO		BOV_RS01020 (thiO)	ABQ60316.1	FAD		4YSH (27)
17	1.4.3.- <sup>b</sup>	<b>Potential Aminoacetone oxidase family FAD-binding enzyme/ NAD (P)/FAD-dependent dehydrogenase</b>		BOV_RS06670	ABQ60616.1	FAD		3V76 (67)
18	1.4.3.- <sup>b</sup>	<b>Potential Aminoacetone oxidase family FAD-binding enzyme/ NAD (P)/FAD-dependent dehydrogenase</b>		BOV_RS04985	ABQ60524.1	FAD		2IOZ (24)
19	1.4.99.- <sup>b</sup>	Predicted D-amino acid dehydrogenase small subunit	DAO (PF01266) 10-400	BOV_RS08480	ABQ61937.1	FAD		6J38 (23)
20	1.4.-.- <sup>b</sup>	<b>Pyridoxamine 5'-phosphate oxidase family protein</b>	Pyridox_ox_2 (PF12900) 10-141	BOV_RS06575	ABQ61684.1	FMN		3U0I (99) 2HQ9 (30)

(Continued on next page)

TABLE 1 (Continued)

Entry	EC	Protein	Pfam domain <sup>a</sup>	Locus tag	Protein code	Flavin	Coenzyme or ligand	More similar PDB (% identity)
21	1.5.1.20	Methylenetetra-hydrofolate reductase	MTHFR (PF02219) 14-291	BOV_RS06945 (metF)	ABQ60279.1	FAD	NAD(P)H	3FST (48)
22	1.5.1.- <sup>b</sup>	<b>Flavin reductase domain containing protein</b>	Flavin_Reduct (PF01613) 22-172	BOV_RS05125	ABQ60228.1	FMN	NAD(P)H	1EJE (29)
23	1.5.3.1	Sarcosine oxidase beta subunit	DAO (PF01266) 36-161	BOV_RS01075 (soxB_1)	ABQ60177.1	FAD &		2GAG_B (64)
		Sarcosine oxidase alpha subunit	DAO (PF01266) 8-222	BOV_RS01090 (soxB_2)	ABQ61310.1	FMN		2GAG_B (58)
			Fer2_4 (PF13510) 16-102	BOV_RS01100	ABQ61036.1	FMN	NADH	2GAG_A (47)
			FAD_oxidore (PF12831) 171-218				4Fe-4S	
			GCV_T (PF01571) 526-790					
			GCV_T_C (PF08669) 815-901					
24	1.5.5.1	Electron transferring flavoprotein-ubiquinone oxidoreductase (ETF-QO)	Thi4 (PF01946) 12-52	BOV_RS03100	ABQ61337.1	FAD	4Fe-4S	2GMH (49)
			ETF_QO (PF05187) 451-560					
25	1.6.5.2	WrpA type FMN-dependent NADH:quinone oxidoreductase	FMN_red (PF03358) 13-145	BOV_RS05025	ABQ60884.1	FMN	quinone	5F4B (98)
26	1.7.- <sup>b</sup>	<b>Predicted NAD(P)H nitroreductase</b>	Nitroreductase (PF00881) 23-167	BOV_RS05130	ABQ60834.1	FMN	quinone	3K6H (43)
27	1.8.1.4	Dihydropyridyl dehydrogenase (lpdA-2)	Pyr_redox_2 (PF07992) 57-399	BOV_RS05390 (lpdA-2)	ABQ60398.1	FAD	NAD(P)H	2A8X (43)
			Pyr_redox_dim (PF02852) 420-528				NADH	
28	1.8.1.4	Dihydropyridyl dehydrogenase (lpdA-3)	Pyr_redox_2 (PF07992) 3-329	BOV_RS09065 (lpdA-3)	ABQ61458.1	FAD	NADH	3URH (78)
			Pyr_redox_dim (PF02852) 348-456					
29	1.8.1.7	Glutathione-disulphide reductase	Pyr_redox_2 (PF07992) 6-321	BOV_RS04850 (gor)	ABQ61016.1	FAD	NADPH	4DNA (69)
			Pyr_redox_dim (PF02852) 341-449					
30	1.8.1.9	Thioredoxin-disulphide reductase	Pyr_redox_2 (PF07992) 8-302	BOV_RS07155 (trx)	ABQ60123.1	FAD	NAD(P)H	4JNQ (100)
31	1.8.1.9	Predicted thioredoxin-disulphide reductase	Pyr_redox_2 (PF07992) 9-171	BOV_RS04925	ABQ61134.1	FAD	NAD(P)H	5YQG (49) <sup>d</sup> 1NHS (30)
								5VJ7 (34)
32	1.14.13.1	Predicted Salicylate hydroxylase	FAD_binding_3 (PF01494) 6-349	BOV_RS03690	ABQ60137.1	FAD	NAD(P)H	4BJZ (32)
33	1.14.13.1	Predicted Salicylate hydroxylase	FAD_binding_3 (PF01494) 2-328	BOV_RS04715	ABQ60978.1	FAD	NAD(P)H	3RP8 (27)
34	1.14.13.- <sup>b</sup>	Predicted UbiH/COQ6 monooxygenase family	FAD_binding_3 (PF01494) 14-322	BOV_RS08970	ABQ60166.1	FAD	NAD(P)H	4K22 (37)
35	1.14.14.3	Bacterial luciferase	Bac_luciferase (PF00296) 8-255	BOV_RS09695	ABQ60348.1	FMN		3FGC (20)
36	1.16.1.4	Cob(II)alamin reductase	Flavin_Reduct (PF01613) 60-208	BOV_RS06210	ABQ60249.1	FMN		3CB0 (99)
37	1.17.1.4	Xanthine dehydrogenase, small subunit	Fer2 (PF00111) 13-59	BOV_RS01845 (xdhA)	ABQ61298.1		NADH	2W3S_A (48)
			Fer2_2 (PF01799) 88-161				2(2Fe-2S)	
			FAD_binding_5 (PF00941) 204-367			FAD		
38	1.18.1.2	Ferredoxin-NADP <sup>+</sup> reductase	CO_deh_flav_C (PF03450) 376-476	BOV_RS01770 (fpr)	ABQ61707.1	FAD		6RRA (100)
			FAD_binding_6 (PF00970) 19-102				NADPH	
			NAD_binding_1 (PF00175) 113-230				NADH	3G5S (46)
39	2.1.1.74	Methylenetetrahydrofolate-tRNA(uracil54-C5)-methyltransferase	GIDA (PF01134) 10-378	BOV_RS04425 (trmFO)	ABQ61275.1	FAD		
40	2.1.1.229	tRNA (carboxymethyluridine34-5-O)-methyltransferase	GIDA (PF01134) 10-399	BOV_RS09735 (mmg)	ABQ60378.1	FAD	NADH	2ZXI (50)
			GIDA_assoc (PF13932) 402-612					
41	2.2.1.6	Acetolactate synthase 3 catalytic subunit	TPP_enzyme_N (PF02776) 1-165	BOV_RS06655 (ilvB)	ABQ60081.1	FAD	Thiamine dIPP	6DEN (46)
			TPP_enzyme_M (PF02005) 191-327					
			TPP_enzyme_C (PF02775) 393-540					

(Continued on next page)

TABLE 1 (Continued)

Entry	EC	Protein	Pfam domain <sup>a</sup>	Locus tag	Protein code	Flavin	Coenzyme or ligand	More similar PDB (% identity)
42	2.5.1.9	Riboflavin synthase alpha subunit	Lum_binding (PF00677) 20-258	BOV_RS03790 (riBE)	ABQ60518.1	RF		4E0F (98)
43	4.2.3.5	Chorismate synthase	Chorismate_synt (PF01264) 10-355	BOV_RS02190 (aroC)	ABQ60200.1	FMN		1UM0 (42)
44	7.1.1.2	NADH-quinone oxidoreductase subunit F (H <sup>+</sup> translocating)	Complex1_51K (PF01512) 47-216 SLB8 (PF10531) 242-292	BOV_RS04000 (nuoF)	ABQ60521.1	FMN	NADH	6Q9C_D (46)
45		Electron transferring flavoprotein alpha subunit (ETFa)	NADH_4Fe-4S (PF10589) 332-414 ETF (PF01012) 44-210	BOV_RS09295 (etfA)	ABQ61011.1		4Fe-4S	1EFP_A (69)
		Electron transferring flavoprotein beta subunit (ETFb)	ETF_alpha (PF00766) 229-312 ETF (PF01012) 29-206	BOV_RS09300 (etfB)	ABQ60428.1	FAD	AMP	1EFP_B (72)

<sup>a</sup>Pfam domains include name, code, and residues in the *B. ovis* flavoprotein making the domain.

<sup>b</sup>Identified as flavoenzyme, but available information does not allow to fully predict its activity.

<sup>c</sup>Shown in bold and italics are those candidates for which a clear function cannot be depicted.

<sup>d</sup>More than one structure to represent the different regions of the protein.

**TABLE 2** Predicted flavoproteins encoded in the chromosome II of *Brucella ovis* ATCC 25840

Entry	EC	Protein	Pfam domains <sup>a</sup>	Locus tag	Protein code	Flavin	Coenzyme or Ligand	More similar PDB (% identity)
46	1.1.2.3	L-lactate dehydrogenase (cytochrome c o b2)	FMN_dh (PF01070) 14-377	BOV_RS14715 (lldD)	ABQ62635.1	FMN		5ZBM (40)
47	1.1.-.- <sup>b</sup>	<b>Potential L-gulonolactone oxidase FAD-binding oxygen oxidoreductase<sup>c</sup></b>	FAD_binding_4 (PF01565) 22-149 nothing up to 444	BOV_RS14405	ABQ62001.1	FAD		4AUT (34)
48	1.1.1.402	D-erythritol 1-phosphate dehydrogenase	DAO (PF01266) 8-333 DAO_C (PF16901) 386-482	BOV_RS14450 (eryB)	ABQ62056.1	FAD	Quinone	2OCU (51)
49	1.1.99.1	Choline dehydrogenase (GMC family, membrane bound)	GMC_oxred_N (PF00732) 5-295 GMC_oxred_C (PF05199) 387-523	BOV_RS14905	ABQ62100.1	FAD		4HA6 (34)
50	1.1.99.2	Predicted L-2-hydroxyglutarate dehydrogenase	DAO (PF01266) 5-392	BOV_RS15155 (lhgO)	ABQ62911.1	FAD	NADH	3DME (37)
51	1.1.99.14	Glycolate dehydrogenase GlcD subunit	FAD_binding_4 (PF01565) 55-193 FAD-oxidase_C (PF02913) 229-470	BOV_RS11160 (glcD)	ABQ62237.1	FAD		3PM9 (29)
52	1.3.1.- <sup>b</sup> / 1.7.1.81	Predicted alkene reductase: N-ethylmaleimide reductase, glycerol trinitrate reductase or xenobiotic reductase B	Oxidored_FMN (PF00724) 3-349	BOV_RS14625	ABQ62490.1	FMN	NAD(P)H	5N6G (69)
53	1.3.8.- <sup>b</sup>	Acyl-CoA dehydrogenase	AidB_N (PF18158) 17-173 Acyl-CoA_dh_M (PF02770) 188-282 Acyl-CoA_dh_1 (PF00441) 292-447	BOV_RS13205	ABQ62576.1	FAD		5EZ3 (100)
54	1.3.8.- <sup>b</sup>	Acyl-CoA dehydrogenase	Acyl-CoA_dh_N (PF02771) 4-116 Acyl-CoA_dh_M (PF02770) 120-214	BOV_RS12330	ABQ62784.1	FAD		4N5F (63)
55	1.3.8.- <sup>b</sup>	Acyl-CoA dehydrogenase	Acyl-CoA_dh_1 (PF00441) 227-375 Acyl-CoA_dh_N (PF02771) 9-115	BOV_RS14135	ABQ62889.1	FAD		5LNX (36)
56	1.3.8.- <sup>b</sup>	Acyl-CoA dehydrogenase	Acyl-CoA_dh_M (PF02770) 120-218 Acyl-CoA_dh_1 (PF00441) 231-378 Acyl-CoA_dh_N (PF02771) 3-157	BOV_RS14115	ABQ62082.1	FAD		6JUC (61)
57	1.3.99.- <sup>b</sup>	Predicted Ksd-like steroid dehydrogenase	Acyl-CoA_dh_1 (PF00441) 282-451 Acyl-CoA_dh_C (PF12806) 467-593	BOV_RS13530	ABQ62061.1	FAD		1D4D (25)
58	1.4.1.13	Glutamate synthase large subunit (alpha subunit)	FAD_binding_2 (PF00890) 5-533 GATase_2 (PF00310) 56-480 Glu_syn_central (PF04898) 508-794 Glu_synthase (PF01645) 856-1230 GXGXG (PF01493) 1309-1498	BOV_RS10585 (glbB)	ABQ61996.1	FMN	3Fe-4S	1EA0 (45)
59	1.4.99.- <sup>b</sup>	Glutamate synthase small subunit (beta subunit)	Fer4_20 (PF14691) 24-131 Pyr_redox_2 (PF07992) 148-465	BOV_RS10590 (glbD)	ABQ62546.1	FAD	NADPH	6S6U_G (37)
60	1.4.-.- <sup>b</sup>	D-amino acid dehydrogenase Predicted D-amino acid dehydrogenase	DAO (PF01266) 3-397 DAO (PF01266) 7-397	BOV_RS14735 (dadA) BOV_RS13345	ABQ62278.1 ABQ62519.1	FAD FAD	4Fe-4S	4YSH (26) 4YSH (26)
61	1.4.-.- <sup>b</sup>	Predicted D-amino acid dehydrogenase	DAO (PF01266) 34-384	BOV_RS13970	ABQ62405.1	FAD		4YSH (24)
62	1.5.3.1	Predicted monomeric Sarcosine oxidase	DAO (PF01266) 31-383	BOV_RS13970	ABQ62932.1	FAD		1ZOV (22)
63	1.6.99.1	<b>NADPH dehydrogenase (Old yellow enzyme)</b>	Oxidored_FMN (PF00724) 2-340	BOV_RS11390	ABQ62422.1	FMN	NADPH	3GR7 (34)
64	1.6.-.- <sup>b</sup>	<b>NADH dehydrogenase</b>	Pyr_redox_2 (PF07992) 8-325	BOV_RS12460	ABQ62704.1	FAD	NADH	4NWZ (31)
65	1.8.1.2	Assimilatory sulphite reductase (NADPH) alpha component cluster	PepSY_TM (PF03929) (3-149) Hypothetical protein (81aa) PepSY_TM (PF03929) (2-128) Flavodoxin_1 (PF00258) (74-147) FAD_binding_6 (PF00970) (80-171)	BOV_RS11420 BOV_RS11425 BOV_RS11430	WP_006015252.1 WP_006015255.1 WP_006015257.1	FAD FAD FAD		TM helices  6EFV (~30 B. melitensis)
66	1.8.1.4	Dihydropolypyl dehydrogenase (lpdA-1)	NAD_binding_1 (PF00175) (189-291) Pyr_redox_2 (PF07992) 7-326 Pyr_redox_dim (PF02852) 345-453	BOV_RS12670 (lpdA-1)	ABQ62466.1	FAD	NADPH NAD(P)H	6CMZ (60)

(Continued on next page)

TABLE 2 (Continued)

Entry	EC	Protein	Pfam domains <sup>a</sup>	Locus tag	Protein code	Flavin	Coenzyme or Ligand	More similar PDB (% identity)
67	1.8.5.81	Peptide-methionine (S)-S-oxide reductase (quinone) (Msr), MsrP catalytic subunit. MsrQ heme-binding subunit	Ferric_reduct (PF01794) 55-166	BOV_RS15075 (msrQ)	ABQ62365.1	FMN	heme b	6HCY_A (17)
68	1.13.11.32	Nitronate monooxygenase (formerly 2-nitropropane dioxygenase NPD)	Oxidored_molyb (PF00174) 98-252 NMO (PF03060) 124-342 (formerly NPD)	BOV_RS15070 (msrP) BOV_RS14290	ABQ62343.1 ABQ62537.1	Molybdopterin FMN	Quinone	1XDY (54) 3BW2 (26)
69	1.13.11.79	Predicted aerobic 5,6-dimethylbenzimidazole synthase (BluB)	Nitroreductase (PF00881) 74-240	BOV_RS15390 (bluB)	ABQ62404.1	FMN	NADH	2ISK (36)
70	1.14.13. <sup>b</sup>	UbiH/UbiF family hydroxylase	FAD_binding_3 (PF01494) 49-377	BOV_RS13080	ABQ62553.1	FAD	NAD(P)	5KOX_A (20)
71	1.14.13.2	4-hydroxybenzoate 3-monooxygenase	FAD_binding_3 (PF01494) 2-342	BOV_RS13400 (pobA)	ABQ62030.1	FAD	NAD(P)H	1PBE (63)
72	1.18.1.3/5	Predicted Ferredoxin/rubredoxin/putidaredoxin NAD <sup>+</sup> Reductase	1: Pyr_redox_2 (PF07992) 4-301 2: Reductase_C (PF14759) 320-404	BOV_RS13795	ABQ62051.1	FAD	NADH	3FG2 (45)
73	1. <sup>c-c</sup> <sup>b</sup>	<b>Predicted nitroreductase family protein</b>	Nitroreductase (PF00881) 48-216	BOV_RS12545	ABQ62091.1	FMN		2IFA (55)
74	2.7.7.2	Bifunctional riboflavin kinase/FAD synthase	FAD_syn (PF06574) 18-172	BOV_RS11255 (ribF)	ABQ62831.1	FMN & FAD RF & FMN		2X0K (36)
75	2.7.1.26	FAD:protein FMN transferase	Flavokinase (PF01687) 190-313	BOV_RS11440	ABQ62066.1	FAD		5MGY (38) 6NXI (36)
76	2.7.13.3	Blue-light-activated histidine kinase	ApbE (PF02424) 18-296					6PPS (100)
			PAS_9 (PF13426) 34-136 PAS_3 (PF08447) 184-259	BOV_RS13160	ABQ62113.1	FMN		
77	4.1.1.36 6.3.2.5	Coenzyme A biosynthesis bifunctional protein: Phosphopantothenoyl-cysteine decarboxylase/Phosphopantothenate-cysteine ligase (CTP)	HWE_HK (PF07536) 285-367 Flavoprotein (PF02441) 7-180 DFP (PF04127) 189-375	BOV_RS15430 (coaBC)	ABQ62036.1	FMN	CTP	1E20 (34) 4QJI (42)
78		Protein NrdI	Flavodoxin_NrdI (PF07972) 5-122	BOV_RS11810 (nrdI)	ABQ62891.1	FMN		2XOD (36)

<sup>a</sup>Pfam domains include name, code, and residues in the *B. ovis* flavoprotein making the domain.<sup>b</sup>Identified as flavoenzyme, but available information does not allow to fully predict its activity.<sup>c</sup>Shown in bold and italics are those candidates for which a clear function cannot be depicted.




sulfite reductase (NADPH) alpha component cluster where FMN and FAD bind respectively at two of its subunits. The bifunctional riboflavin kinase/FAD synthase would bind RF, FMN, and FAD, and the riboflavin synthase alpha subunit would bind RF as product. Despite the lower content of proteins binding FMN, the *B. ovis* flavoproteome is slightly biased toward FMN when considering overall kingdoms, where the majority of flavoenzymes bind FAD (75%) (34), or some eukaryotic flavoproteomes, as the *Homo sapiens* one (with 84% FAD-dependent proteins) (6, 38). Thirty-three of the identified flavoproteins in *B. ovis* are expected to use NAD(P)<sup>+</sup>/H as coenzyme, whereas a few would bind haem, iron-sulfur clusters, quinones, CTP, or thiamine phosphate. It is worth mentioning that currently many of these proteins are only computationally annotated and, in many cases, with vague confidence regarding ligands (searching motifs for FAD also find NAD(P)H binding proteins), nature of the protein, and metabolic functions.

**The structure conformational space in the *B. ovis* flavoproteome.** 3D structures of proteins from *B. ovis* are scarce: 10 different proteins in 14 Protein Data Bank (PDB) entries (Table SP2). Only one corresponds to a flavoprotein that is also NADP<sup>+</sup>/H dependent, Ferredoxin-NADP<sup>+</sup> reductase (FPR) (39), and three more correspond to structures, or subunits, of NAD(P)<sup>+</sup>/H dependent proteins (Table SP2). The PDB contains also 11 structures of potential flavoproteins from other *Brucella* (Table SP3 and Fig. SP2). Ten correspond to nine *B. ovis* homologues sharing more than 98.5% identity, and two are also NAD(P)<sup>+</sup>/H dependent. For the remaining structure, the best match in *B. ovis* ATCC 25840 only shares 33% identity, but an identical sequence is found in *B. ovis* IntaBari-2002-82-58. Five of these flavoproteins are FAD-dependent (only two show FAD in the structure), three are FMN-dependent, and one might bind either RF, FMN, or FAD (Table SP3). Regarding function, seven are oxidoreductases with NAD(P)<sup>+</sup>/H (quinone) dehydrogenase, acyl-CoA dehydrogenase, thioredoxin reductase, or monooxygenase activities. Three are transferases; one is riboflavin synthase and two relate to the LOV domain of sensory histidine kinase. Notably, the PDB entry 3U0I from *Brucella melitensis*, identified as a pyridoxamine 5'-phosphate oxidase family protein of unknown function, shares nearly 100% identity with *B. ovis* ABQ61684.1. Most of these structures come from structural genomic projects on *B. melitensis*, *Brucella abortus*, and *Brucella suis*, and their functions are not experimentally curated. Tables 1 and 2 show that in 66 (85%) and 39 (50%) of the candidates there are structures of homologues with more than 30% and 45% sequence identity, respectively, providing good structural models for at least ~50% of the flavoproteins.

*B. ovis* flavoproteins use up to 26 Pfam clans and 73 domain families (Tables 1 and 2, Fig. 1, and Fig. SP1), and in 80% of the cases they fold using more than one domain. Nonetheless, only 13 clans, plus six domains not assigned to any clan, are implicated in flavin binding (Fig. SP1), in agreement with the diversity in structural topologies for the interaction of these cofactors. The NADP\_Rossmann and TIM-barrel clans are the most widely represented and preferred respectively for FAD and FMN binding in *B. ovis* flavoproteins, being highly spread in proteins and particularly within flavoproteins (5, 7). The NADP\_Rossmann fold appears in 38 flavoproteins in *B. ovis*, but presumably it is not involved in flavin binding in domains FAD\_oxidored of sarcosine oxidase alpha subunit and DFP of coenzyme A biosynthesis bifunctional protein, where it respectively binds NADH and CTP. From the 36 remaining flavoproteins, 35 bind FAD and riboflavin synthase binds RF (Lum\_binding domain). NADP\_Rossmann flavoproteins use mostly domains DAO (some completed by DAO\_C domain caps) and Pyr\_redox\_2 (might contain also Pyr\_redox\_dim or Fer4\_20 domains). Domain families FAD\_binding\_3, GMC\_oxired\_N (also having GMC\_oxired\_C), FAD\_binding\_2 (some completed with Succ\_DH\_flav\_C), GIDA (holding also GIDA\_assoc), HI0933\_like, and Thi4 are also represented. In this clan we want to notice a particular case: two proteins with DAO domains shorter than regular ones, ABQ60177.1 (BOV\_RS01075, *soxB\_1*) and ABQ61310.1 (BOV\_RS01090, *soxB\_2*), cover the primary sequence of the sarcosine oxidase beta subunit binding FAD in other *Brucella* orthologues (Table 1, Fig. 2A). On its side, the TIM\_barrel





**FIG 1** Distribution of *B. ovis* ATCC 25840 flavoproteins in structural Pfam clans and domain families. (A) Pie chart of the clans and domains found within the identified flavoproteins. Clan names are highlighted in bold. Names of domain families directly involved in flavin binding are colored in orange, whereas domains not involved in flavin binding but present in the flavoproteins are in black. Overall percentages are based on the number of domain families in each clan (including (Continued on next page)

clan is represented by nine flavoproteins that use up to seven different domains: Oxidored\_FMN, Dus, DHO\_dh, FMN\_dh, Glu\_syn\_central, Glu-synthase, and NMO.

Clans AcylCoA\_ox\_dh\_N and AcylCoA\_dh appear in seven FAD dependent acyl-CoA dehydrogenases. Normally, they fold in Acyl-CoA\_dh\_N (in one case replaced by Aidb\_N) and Acyl-CoA\_dh\_1 domains and use connecting Acyl-CoA\_dh\_M and/or ending Acyl-CoA\_dh\_C domains. The six flavoproteins of the FAD\_PCMH clan bind FAD; five and one use, respectively, FAD\_binding\_4 and FAD\_binding\_5 domains (completed with FAD-oxidase\_C, CO\_deh\_flav\_C, or murB\_C domains).

The FMN-binding clan is predicted in four flavoproteins that bind FMN with Putative\_PNPOx (with PNP\_phzG\_C domain), Flavin\_Reduct, and Pyridox\_ox\_2 domains. Three also FMN binding flavoproteins fall in each of the FMN-dep-Nrtase and Flavoprotein clans. The first have nitroreductase domains, presumably involved in reduction of nitrogen-containing compounds, whereas those in the second clan belong each to a different family: FMN\_red (NAD(P)H-quinone dehydrogenase), Flavodoxin\_NrdI (electron transport NrdI protein), and Flavodoxin\_1 (assimilatory sulfite reductase alpha component). This subunit of assimilatory sulfite reductase contains also a FAD\_binding\_6 domain, of the FAD\_Lum\_binding clan, that binds FAD. A FAD\_binding\_6 domain is also present in ferredoxin-NADP<sup>+</sup> reductase. The assimilatory sulfite reductase alpha component in *B. ovis* shows noticeable features (Fig. 2B). In the current genome assembly, it is annotated as codified by four sequential genes (BOV\_RS11420, BOV\_RS11425, BOV\_RS11430, and BOV\_RS11435), which would make up four protein subunits (WP\_006015252.1, WP\_006015255.1, WP\_006015257.1, and WP\_006015259.1), whereas in other *Brucella* all these components are encoded by either one or two genes (Fig. 2B). Its central Flavodoxin\_1 and FAD\_binding\_6 domains allocate FMN and FAD, respectively, whereas the additional domains will attach the protein to the membrane and bind the NADPH coenzyme. The functional protein will be complemented with a sulfite reductase (NADPH) haemoprotein beta-component (ABQ61351) codified in Cl.


Only two *B. ovis* flavoproteins are members of the FADS\_DHS clan. They adopt a Rossmann fold similar to clan NADP\_Rossmann, but are distinguished since the FAD cofactor binds in the opposite direction. It is represented by the TPP\_enzyme\_M domain in the catalytic subunit of a synthase and ETF\_alpha domain in the electron transferring flavoprotein alpha subunit (ETFa). Four clans are only found once in the *B. ovis* flavoproteome: the FAD\_oxidored in the FAD dependent MTHFR domain of methylenetetrahydrofolate reductase; the PAS\_Fold with the PAS\_9 domain that binds FMN in the blue-light-activated histidine kinase; the HUP that binds FMN/FAD in the FAD\_syn domain of bifunctional FAD synthase/flavokinase (FADS); and the 2heme\_cytochrom binding FMN at the Ferric\_reduct domain of MsrQ subunit of peptide-methionine (S)-S-oxide reductase (Msr).

Six domains not assigned to any clan also bind flavins in *B. ovis* flavoproteins: the Flavokinase domain binds RF/FMN in bifunctional FADS; the Bac\_luciferase domain binds FMN in bacterial luciferase; the Chorismate\_synt domain binds FMN in chorismate synthase; the NADH\_4Fe\_4S domain binds FMN at the NADH-quinone oxidoreductase subunit F; the ApbE domain binds FAD in the FAD:protein FMN transferase; and the Flavoprotein domain binds FMN in Coenzyme A biosynthesis bifunctional protein.


In agreement with the use of FAD-dependent enzymes for novel or unusual functions requiring the adaptation of already existing topologies or new structural designs (7), the less populated clans and domains in the *B. ovis* flavoproteome mainly contribute to FMN binding.

### FIG 1 Legend (Continued)

domains involved and not involved in flavin binding). Most populated clans concerning structural folding, NADP\_Rossmann and TIM\_barrel, are shown individually, while the rest are grouped according to the number of families found in each: three (gray), two (orange), or one (green). The blue portion includes domain families with no clan assigned. The number for a particular domain presented in each chromosome is denoted in brackets, as *N* in Cl/*N* and ClI/*N*. Details for only flavin binding domains are shown in Figure SP1. (B) Clans involved in flavin cofactor binding according to the flavin type. (C) Distribution of flavin binding clans by chromosomal location.



**FIG 2** Genomic context for *SoxB* and sulfite reductase component genes in selected *Brucella* species. (A) Organization of genes encoding different subunits of the membrane bound sarcosine oxidase subunit B (*SoxB*). In *B. ovis*, two genes, *soxB\_1* and *soxB\_2*, separated by a IS5 transposase gene, encode together the full length of the *SoxB* protein. On the top, the homology structural models of *SoxB\_1* and *SoxB\_2* proteins and their superposition on the *SoxB* from *Stenotrophomonas maltophilia* (2GAG\_B) are shown. (B) Organization of genes encoding for the assimilatory sulfite reductase alpha component. In both panels, gene senses are denoted by arrows and *B. ovis* gene codes are written next to the corresponding arrow. Structural Pfam domain families for subunits contributing to flavin binding are shown on the top of genes.



**FIG 3** Structural models for *B. ovis* enzymes predicted to covalently bind the flavin cofactor. (A) succinate dehydrogenase flavoprotein subunit and (B) sarcosine oxidase beta subunit models (orange) for the covalent linking to C8M of flavins in *B. ovis*. Homology models were built using as templates the structures of *Ascaris suum* (PDB 4YSX\_A, light blue in A) and *S. maltophilia* (2GAG, green in B), respectively, and the RaptorX server. Parameters predicting models' quality are summarized below the figure. (C) Structure of the blue-light-activated histidine kinase from *Brucella abortus* 2308 (6PPS), with 100% identity to the *B. ovis* enzyme. Relevant residues as well as cofactors belonging to the templates are shown in sticks. Images generated with PyMol (134).

From the 78 identified candidates, only two flavoproteins are envisaged to bind flavins covalently. One is the succinate dehydrogenase flavoprotein subunit (SdhA), where covalent binding through H60 (in the conserved FPTRSHTV motif) to the FAD isoalloxazine (C8M) is predicted, as observed for H45, H56, or H79 in *E. coli* (PDB 2WDQ), *Gallus* *Sudha* (PDB 2H88), or *Ascaris suum* (PDB 4YSX) proteins (Fig. 3A). Covalent attachment of FAD to SdhA is essential for Sdh function in other bacteria (19). Covalent linking to FMN (C8M) is also predicted in the sarcosine oxidase beta subunit through H198 (H173 in the *Stenotrophomonas maltophilia* protein, PDB 2GAG), placing FMN at the alpha and beta subunits interface (Fig. 3B). Sequence and structural modeling also point to blue-light-activated histidine kinase as undergoing photochemistry with cysteinyl-flavin adduct formation between C69 and the isoalloxazine C4a of FMN (Fig. 3C), altogether these observations point out that the cofactor is noncovalently bound in most *B. ovis* flavoproteins.

#### Enzymatic classification and metabolic functions of *Brucella ovis* flavoproteins.

Most proteins of the *B. ovis* flavoproteome (76 out of 78) are flavoenzymes. We assigned enzymatic classes to all of them and full Enzyme Commission (EC) codes to nearly 70% (Table 3, Fig. 4, 5 and 6). The rest (30%) might have either divergent mechanisms from known flavoproteins or still not reported full functions. Flavoenzymes fall into the oxidoreductases (EC 1.) (86%, 66 out of 76), transferases (EC 2.) (9.2%, 7 out of 76, one of them with two transferase activities), lyases (EC 4.) (2.6%, 2 out of 76), and translocases (EC 7.) (1.3%, 1 out of 76) classes (Tables 1 and 2, and Table SP6; Fig. 7A). Therefore, most of *B. ovis* flavoenzymes participate in redox processes, in agreement with previous reported classifications (5, 7).

Two flavoproteins are predicted not holding catalytic activity by themselves: ETFa and NrdI. The ETFa together with the beta subunit (ETFb) specifically transfer electrons from different dehydrogenases to the respiratory electron transfer chain (ETC) via the electron transferring flavoprotein-ubiquinone oxidoreductase (ETF-QO). Noticeably, *etfA* and *etfB* encoding genes in *B. ovis* sit next to a gene encoding for an NADP<sup>+</sup> dependent butyryl-CoA dehydrogenase (Table SP4). This suggests that these three proteins might form a BCD bifurcating complex responsible for crotonyl-CoA reduction during butanol production. Homologues of *Clostridium acetobutylicum* have the same organization (40),



**TABLE 3** Metabolic functions and virulence potential envisaged for flavoproteins and flavoenzymes from *B. ovis* ATCC 25840. Search for metabolic functions and virulence prediction carried out as indicated in Materials and Methods

EC	Protein	Protein code	Metabolic pathway	Pathway category	VirulentPred <sup>b</sup>	VICMp <sup>pred</sup> function <sup>c</sup>
1.1.5.3	Glycerol-3-phosphate dehydrogenase	ABQ60174.1	Degradation of sugar alcohols	Carbohydrate metabolism	Non-virulent	Metabolism
1.1.1.402	D-erythritol 1-phosphate dehydrogenase	ABQ62056.1	Degradation of sugar alcohols	Carbohydrate metabolism	Non-virulent	Metabolism
1.1.2.3	L-lactate dehydrogenase (cytochrome c or b2)	ABQ62635.1	Lactate fermentation	<b>Virulence factor.</b> Fermentation and other catabolism	Non-virulent	Metabolism
1.1.3.8	<b>Potential L-gulonolactone oxidase FAD-binding oxygen oxidoreductase<sup>e</sup></b>	ABQ62001.1	Unclear function. Potentially involved in ascorbate metabolism	Potential role in nucleotide and cofactor metabolism	Non-virulent	Cellular Process
1.1.99.1	Choline dehydrogenase (Glucose-methanol-choline GMC family)	ABQ61350.1	Glycine betaine biosynthesis.	Amino acid Metabolism.	Non-virulent	<b>Virulence Factor</b>
1.1.99.1	Choline dehydrogenase (GMC family)	ABQ60630.1	Metabolism of disaccharides	Carbohydrate metabolism	<b>Virulent</b> Non-virulent	Cellular Process Metabolism / <b>Potential Virulence Factor</b>
1.1.99.1	Choline dehydrogenase (GMC family, membrane bound)	ABQ62100.1			<b>Virulent</b>	Metabolism
1.1.99.2	Predicted L-2-hydroxyglutarate dehydrogenase	ABQ62911.1	Glutamate and glutamine metabolism	Amino acid metabolism	Non-virulent	Metabolism
1.1.99.14	Glycolate dehydrogenase GlcD subunit	ABQ62237.1	Glycolate and Glyoxylate degradation	Central and energy metabolism	Non-virulent	Metabolism
1.1.-.-	<b>Potential FAD-binding oxygen oxidoreductase (glcE?)</b>	ABQ60928.1	It might work with GlcD	Probably central and energy metabolism	Non-virulent	Metabolism
1.1.-.-	<b>Potential FAD-binding oxygen oxidoreductase (glcE?)</b>	ABQ61939.1	It might work with GlcD	Probably central and energy metabolism	Non-virulent	Metabolism
1.3.1.1/2	NADH dependent Dihydropyrimidine dehydrogenase subunit PreA	ABQ60560.1	Pyrimidine and alanine metabolism	Nucleotide and cofactor metabolism. Amino acid metabolism	Non-virulent	Cellular Process / Metabolism
1.3.5.2	NADH dependent Dihydropyrimidine dehydrogenase subunit PreT	ABQ61103.1			Non-virulent	Cellular Process / Metabolism
1.3.5.2	PyrD dihydroorotate dehydrogenase 2 (quinone)	ABQ61413.1			Non-virulent	Metabolism
1.3.1.88	tRNA dihydrouridine synthase B	ABQ61416.1	Dihydrouridine modification of tRNA	Modification of cytoplasmic tRNAs	Non-virulent	Metabolism
1.3.1.91	tRNA dihydrouridine20/20a synthase	ABQ61966.1			Non-virulent	Metabolism
1.3.1.98	UDP-N-acetylmuramate dehydrogenase	ABQ61769.1	Peptidoglycan Biosynthesis. Cell wall biogenesis	Amino acid metabolism	<b>Virulent</b>	<b>Virulence Factor</b>
1.3.1.- /1.7.1.B1	Predicted alkene reductase: N-ethylmaleimide reductase, Glycerol trinitrate reductase or xenobiotic reductase B	ABQ62490.1	Degradation of toxic compounds	Xenobiotic metabolism and secondary metabolism	Non-virulent	Metabolism
1.3.5.1	Succinate dehydrogenase flavoprotein subunit	ABQ61077.1	Citric acid and methylaspartate cycles. Propionate fermentation	Fermentation and other catabolism. Central and energy metabolism.	Non-virulent	<b>Virulence Factor</b>
1.3.8.4	Isovaleryl-CoA dehydrogenase	ABQ60382.1	Leucine metabolism	<b>Virulence factor</b> Amino acid metabolism	<b>Virulent</b>	Cellular Process
1.3.8.-	Short Chain Acyl-CoA dehydrogenase	ABQ60180.1	Lipid metabolism. Butanone fermentation. Alanine, glutamate and glutamine metabolism. Ethylmalonyl-CoA pathway	Lipid metabolism. Fermentation and other catabolism. Amino acid metabolism. Central and energy metabolism	Non-virulent	Cellular Process

(Continued on next page)

TABLE 3 (Continued)

EC	Protein	Protein code	Metabolic pathway	Pathway category	VirulentPred <sup>b</sup>	VICMPred function <sup>c</sup>
1.3.8.1	Acyl-CoA dehydrogenase	ABQ62576.1	Bacterial resistance during alkylolation stress/Cell division/AidB domains	Lipid and steroid metabolism. Resistance during alkylolation stress. Defence from Host	<b>Virulent</b>	Metabolism
1.3.8.-	Acyl-CoA dehydrogenase	ABQ61585.1	Lipid metabolism. Butanone fermentation. Valine, alanine, tryptophan, glutamate and glutamine metabolisms.	Lipid metabolism,	Non-virulent	Metabolism
1.3.8.-	Acyl-CoA dehydrogenase	ABQ62784.1		Fermentation and other	Non-virulent	<b>Virulence Factor</b>
1.3.8.-	Acyl-CoA dehydrogenase	ABQ62889.1		catabolism, Amino acid metabolism, Central and energy metabolism.	Non-virulent	<b>Virulence Factor</b>
1.3.8.-	Acyl-CoA dehydrogenase	ABQ62082.1	Ethylmalonyl-CoA pathway. Phenyl acetate degradation (aerobic). Cyclohexanol degradation. Adipate degradation		Non-virulent	Metabolism
1.3.99.-	Predicted KsdD-like steroid dehydrogenase	ABQ62061.1	Androgen and steroid metabolism	Lipid metabolism	Non-virulent	Metabolism
1.4.1.13	Glutamate synthase large subunit (alpha subunit)	ABQ61996.1	Glutamate and glutamine metabolism	Amino acid metabolism	Non-virulent	Metabolism
1.4.3.5	Glutamate synthase small subunit (beta subunit)	ABQ62546.1				
1.4.3.5	Pyridoxamine 5'-phosphate oxidase	ABQ60142.1	Vitamin B6 metabolism	Nucleotide and cofactor metabolism	Non-virulent	Cellular Process / Metabolism
1.4.-.-	<b>Pyridoxamine 5'-phosphate oxidase family protein</b>	ABQ61684.1	Unclear function	Unclear function	<b>Virulent</b>	Cellular Process / Metabolism
1.4.3.19	Glycine oxidase ThiO	ABQ60316.1	Thiamine metabolism	Amino acid metabolism	Non-virulent	Metabolism
1.4.3.-	<b>Potential Aminoacetone oxidase family FAD-binding enzyme/ NAD(P)/FAD-dependent dehydrogenase</b>	ABQ60616.1	Potential role in amino acids and NAD metabolism	Amino acid metabolism. Nucleotide and cofactor metabolism	Non-virulent	<b>Virulence Factor</b>
1.4.3.-	<b>Potential Aminoacetone oxidase family FAD-binding enzyme/ NAD(P)/FAD-dependent dehydrogenase</b>	ABQ60524.1	Potential role in amino acids and NAD metabolism	Amino acid metabolism. Nucleotide and cofactor metabolism	Non-virulent	Metabolism
1.4.99.-	Predicted D-amino acid dehydrogenase small subunit	ABQ61937.1	Potential role in oxidative deamination of D-amino acids	Amino acid metabolism	Non-virulent	<b>Virulence Factor</b>
1.4.99.-	D-amino acid dehydrogenase	ABQ62278.1			<b>Virulent</b>	Metabolism / <b>Potential Virulence Factor</b>
1.4.-.-	Predicted D-amino acid dehydrogenase	ABQ62519.1			<b>Virulent</b>	Cellular Process
1.4.-.-	Predicted D-amino acid dehydrogenase	ABQ62405.1			Non-virulent	Metabolism
1.5.1.20	Methylenetetrahydrofolate reductase	ABQ60279.1	Tetrahydrofolate metabolism	Nucleotide and cofactor metabolism	Non-virulent	Cellular Process / Metabolism
1.5.1.-	<b>Flavin reductase domain containing protein</b>	ABQ60228.1	Flavin metabolism	Nucleotide and cofactor metabolism	<b>Virulent</b>	Cellular Process / Metabolism
1.5.3.1	Sarcosine oxidase beta subunit	ABQ60177.1	Creatine degradation	Fermentation and other catabolism	<b>Virulent</b>	Cellular Process / <b>Virulence Factor</b>
1.5.3.1	Sarcosine oxidase alpha subunit	ABQ61310.1			Non-virulent	
1.5.3.1	Predicted monomeric Sarcosine oxidase	ABQ61036.1			Non-virulent	
1.5.3.1	Predicted monomeric Sarcosine oxidase	ABQ62932.1				
1.5.5.1	Electron transferring flavoprotein-ubiquinone oxidoreductase (ETF-QO)	ABQ61337.1	Oxidative phosphorylation	Fermentation and other catabolism	Non-virulent	Cellular Process / Metabolism
1.5.5.1	Electron transferring flavoprotein-ubiquinone oxidoreductase (ETF-QO)	ABQ61337.1	Oxidative phosphorylation	Central and energy metabolism	Non-virulent	Metabolism

(Continued on next page)

TABLE 3 (Continued)

EC	Protein	Protein code	Metabolic pathway	Pathway category	VirulentPred <sup>b</sup>	VICMPred function <sup>c</sup>
1.6.5.2	WrpA type FMN-dependent NADH:quinone oxidoreductase	ABQ60884.1	Potential role in protection from stress damage	Defence from Host	Non-virulent	Metabolism
1.6.99.1	<b>NADPH dehydrogenase (Old yellow enzyme)</b>	ABQ62422.1	Unclear function	Unclear function	Non-virulent	Cellular Process
1.6.-.-	<b>NADH dehydrogenase</b>	ABQ62704.1	Unclear function	Unclear function	Non-virulent	Metabolism
1.7.-.-	<b>Predicted NAD(P)H nitroreductase</b>	ABQ60834.1	Oxidation-reduction of diverse nitrogen containing compounds	Nitrogen metabolism	Non-virulent	Metabolism
1.8.1.2	Assimilatory sulphite reductase (NADPH) alpha component cluster	WP_006015252.1 WP_006015255.1 WP_006015257.1 WP_006015259.1	Sulphate reduction	Amino acid metabolism	<b>Virulent</b>	Cellular Process / Metabolism
1.8.1.4	Dihydrolipoyl dehydrogenase (lpdA-1)	ABQ62466.1	Oxidative decarboxylation of pyruvate. Citric acid cycle.	Central and energy metabolism. Amino acid metabolism	Non-virulent	Cellular Process
1.8.1.4	Dihydrolipoyl dehydrogenase (lpdA-2)	ABQ60398.1	Glycine metabolism. Acetyl-CoA biosynthesis		Non-virulent	Metabolism
1.8.1.4	Dihydrolipoyl dehydrogenase (lpdA-3)	ABQ61458.1			Non-virulent	Metabolism
1.8.1.7	Glutathione-disulphide reductase	ABQ61016.1	Thiol redox pathway. Glutathione metabolism	Control the redox state of the cell	Non-virulent	Cellular Process
1.8.1.9	Thioredoxin-disulphide reductase	ABQ60123.1	Thiol thioredoxin related pathway. Reduction of cytoplasmic enzymes	Control the redox state of the cell	Non-virulent	Metabolism
1.8.1.9	Predicted thioredoxin-disulphide reductase	ABQ61134.1		Control the redox state of the cell	Non-virulent	Metabolism
1.8.5.B1	Peptide-methionine (S)-S-oxide reductase (quinone) (Msr). MsrP catalytic subunit. MsrQ heme-binding subunit	ABQ62365.1 ABQ62343.1	Methionine reparation of periplasmic proteins	Protection from stress damage. Defence from host	<b>Virulent</b>	Cellular Process / Metabolism
1.13.11.32	Nitronate monooxygenase (formerly 2-nitropropane dioxygenase NPD)	ABQ62537.1	Alkyl/nitronates degradation	Nitrogen metabolism	Non-virulent	Metabolism
1.13.11.79	Predicted aerobic 5,6-dimethylbenzimidazole synthase (BluB)	ABQ62404.1	Vitamin B12 metabolism	Nucleotide and cofactor metabolism	<b>Virulent</b>	Metabolism
1.14.13.1	Predicted Salicylate hydroxylase	ABQ60137.1	Phenol degradation	Xenobiotic metabolism and secondary metabolism	Non-virulent	Metabolism
1.14.13.1	Predicted Salicylate hydroxylase	ABQ60978.1	Phenol degradation	Xenobiotic metabolism and secondary metabolism	<b>Virulent</b>	<b>Virulence Factor</b>
1.14.13.-	Predicted UbiH/COQ6 monooxygenase family	ABQ60166.1	Ubiquinone biosynthesis	Central and energy metabolism	Non-virulent	<b>Virulence Factor</b>
1.14.13.-	UbiH/UbiF family hydroxylase	ABQ62553.1	Ubiquinone biosynthesis	Central and energy metabolism	Non-virulent	<b>Virulence Factor</b>
1.14.13.2	4-hydroxybenzoate 3-monooxygenase	ABQ62030.1	4-hydroxymandelate degradation	Fermentation and other catabolism	Non-virulent	Metabolism
1.14.14.3	Bacterial luciferase	ABQ60348.1	Bioluminescence	Bacterial luminescence	Non-virulent	Cellular Process
1.16.1.4	Cob(I)alamin reductase	ABQ60249.1	Vitamin B12 metabolism	Nucleotide and cofactor metabolism	<b>Virulent</b>	Metabolism
1.17.1.4	Xanthine dehydrogenase, small subunit	ABQ61298.1	Purine metabolism	Nucleotide and cofactor metabolism	Non-virulent	Cellular Process / <b>Potential Virulence factor</b>
1.18.1.3-5	Predicted Ferredoxin/rubredoxin/putidaredoxin NAD <sup>+</sup> Reductase	ABQ62051.1	Protection from ROS stress damage	Defence from host	Non-virulent	<b>Virulence Factor</b>
1.18.1.2	Ferredoxin-NADP <sup>+</sup> reductase	ABQ61707.1	Protection from ROS stress damage	Defence from host	Non-virulent	Metabolism


(Continued on next page)



TABLE 3 (Continued)

EC	Protein	Protein code	Metabolic pathway	Pathway category	VirulentPred <sup>b</sup>	VICMPred function <sup>c</sup>
1.-.-.-	<b>Predicted nitroreductase family protein</b>	<b>ABQ62091.1</b>	Reduction of nitrogen containing compounds.	Nitrogen metabolism	Non-virulent	Metabolism
2.1.1.74	Methylenetetrahydrofolate-tRNA-(uracil54-C5-)-methyltransferase	<b>ABQ61275.1</b>	Post-translationally modification of tRNA	Modification of cytoplasmic tRNAs	Non-virulent	<b>Virulence Factor</b>
2.1.1.229	tRNA (carboxymethyluridine34-5-O)-methyltransferase	<b>ABQ60378.1</b>	Post-translationally modification of tRNA	Modification of cytoplasmic tRNAs	Non-virulent	<b>Virulence Factor</b>
2.2.1.6	Acetolactate synthase 3 catalytic subunit	<b>ABQ60081.1</b>	Acetoin degradation, valine and isoleucine metabolism	Fermentation and other catabolism. Amino acid metabolism	Non-virulent	Metabolism
2.5.1.9	Riboflavin synthase alpha subunit	<b>ABQ60518.1</b>	Flavin biosynthesis	Nucleotide and cofactor metabolism	Non-virulent	Cellular Process / Metabolism
2.7.7.2	Bifunctional riboflavin kinase/FAD synthase	<b>ABQ62831.1</b>	Flavin biosynthesis	Nucleotide and cofactor metabolism	Non-virulent	Metabolism
2.7.1.26	FAD:protein FMN transferase	<b>ABQ62066.1</b>	Flavin transfer	Maturation of other enzymes	Non-virulent	Metabolism
2.7.1.180	Blue-light-activated histidine kinase	<b>ABQ62113.1</b>	Light activated phosphorylation	Signal transduction.	Non-virulent	Cellular Process / <b>Potential Virulence Factor</b>
4.1.1.36	Coenzyme A biosynthesis bifunctional protein	<b>ABQ62036.1</b>	Coenzyme A metabolism	Nucleotide and cofactor metabolism	Non-virulent	Metabolism
6.3.2.5	Chorismate synthase	<b>ABQ60200.1</b>	Chorismate metabolism	Amino acid metabolism	Non-virulent	Cellular Process
4.2.3.5	NADH-quinone oxidoreductase subunit F	<b>ABQ60521.1</b>	Oxidative phosphorylation	Central and energy metabolism	Non-virulent	Cellular Process
7.1.1.2	(H <sup>+</sup> translocating) Electron transferring flavoprotein alpha subunit (ETFa)	<b>ABQ61011.1</b>	Oxidative phosphorylation	Central and energy metabolism	Non-virulent	Cellular Process / Metabolism
	Electron transferring flavoprotein beta subunit (ETFb)	<b>ABQ60428.1</b>				
	Protein NrdI	<b>ABQ62891.1</b>	Electron exchange, nucleotide transport and metabolism	Nucleotide and cofactor metabolism	<b>Virulent</b>	Cellular Process

<sup>a</sup>Shown in bold and italics are those candidates for which a clear function cannot be depicted.  
<sup>b</sup>VirulentPred was used to predict potential virulent proteins based on amino-acid compositions, similarity, position specific scoring matrix, dipeptide composition, higher order dipeptide composition, and the bi-layer cascade Support Vector Machine module. When one of these approaches produced a match, the protein was labeled as virulent. Positive virulence is highlighted in bold.  
<sup>c</sup>The pattern based score of VICMPred was used to classify potential functions for proteins among cellular process, metabolism, signaling, or virulence factor. Positive virulence highlighted in bold.




and ETF proteins are also pointed to as potential targets for the treatment of some bacterial infections (41). NrdI is a flavodoxin-like electron-transport protein with potential analogous functions to ferredoxins. It belongs to the core proteome of *Brucella* but is hardly populated in the evaluated alphaproteobacteria (Table SP6).

**B. ovis flavo-oxidoreductases participate in a large variety of metabolic pathways.** Most *B. ovis* flavoenzymes are oxidoreductases and particularly belong to subclasses that use alcohols (EC 1.1.), CH-CH (EC 1.3.), CH-NH<sub>2</sub> (EC 1.4.), CH-NH (EC 1.5.), or sulfur (EC 1.8.) groups as hydrogen or electron donors, as well as paired donors with incorporation or reduction of molecular oxygen (EC 1.14.). Some use as donors NAD(P)H (EC 1.6.), metals (EC 1.16), or nitrogenous compounds (various EC 1. subclasses) (Tables 1–3, Fig. 7A).

The subclass alcohol flavo-oxidoreductase (Fig. 4A) includes three enzymes of the GMC family (EC 1.1.99.1). This large and diverse protein family shares structural fold and reaction mechanisms, with substrates ranging from sugars and alcohols to cholesterol and choline, which are usually related to the metabolism of methyl groups through betaine (Tables 1–3). They are present in yeasts, bacteria, insects and filamentous fungi, being the latest use for biomass utilization, biosensors, or food industry (42). This subclass has also two alcohol oxidases that share 55% identity and phylogenetically cluster together (bootstrap >98) (Fig. 4A). One is a canonical glycerol-3-phosphate dehydrogenase (GlpD) (EC 1.1.5.3, [ABQ60174.1](#)), and the other is envisaged erythritol oxidation activity (EC 1.1.1.402, [ABQ62056.1](#)) from gene context (BOV\_RS14450, *eryB*) and phylogenetic analysis (Tables 1–3, Fig. 4A). They likely supplement electrons for aerobic oxidative phosphorylation (OXPHOS) at the central junction of glycolysis, respiration, and phospholipid biosynthesis, being essential for aerobic growth on glycerol-like molecules (43). Noticeably, erythritol has a growth-promoting effect on intracellular *Brucella* pathogens (44, 45). GlpD is in the *Brucella* core proteome, but EryB lacks in some *Brucella* (Table SP6, Fig. 4A). These observations highlight the importance of the shuttle of electrons in the *Brucella* metabolism. Other enzymes in this subclass are oxygen oxidoreductases. One is the GlcD subunit of glycolate dehydrogenase that catalyzes oxidation of glycolate and D-lactate, respectively, to glyoxylate and pyruvate (EC 1.1.99.2), a key function in microbial redox oxidative metabolisms. Glycolate dehydrogenase is usually built by several subunits, including a GlcE one. In *B. ovis* the gene upstream of *glcD* is a *glcE* pseudogene (Table SP5). Nonetheless, the oxygen oxidoreductases [ABQ60928.1](#) and [ABQ61939.1](#) have similar features to GlcE and not precise function, so they might replace the non-functional GlcE protein (Table 2 and Table SP6). The protein [ABQ62001.1](#) (EC 1.1.3.8) is also proposed as oxygen oxidoreductase. It has a nonclassified long C-terminal (Table 2) and its closer available structural homologue is decaprenyl-phosphoryl-β-D-ribofuranose-2-oxidoreductase from *Mycobacterium smegmatis*, an essential enzyme in cell wall biosynthesis (46). This subclass also includes the hydroxyglutarate dehydrogenase (EC 1.1.99.2) and the FMN-dependent L-lactate dehydrogenase (EC 1.1.2.3), both in the core proteome of *Brucella* (Table SP6). Altogether, alcohol flavo-oxidoreductases contribute to the energetic intake metabolism for *B. ovis* survival.

#### FIG 4 Legend (Continued)

>90) three proteins of the GMC family, [ABQ61350.1](#), [ABQ60630.1](#), and [ABQ62100.1](#) (EC. 1.1.99.1). Clade 2 groups (bootstrap > 95) three proteins folding in DAO domains, [ABQ60174.1](#), [ABQ62056.1](#), and [ABQ62911.1](#). Clade 3 clusters five proteins distributed in monophyletic branches [ABQ61939.1](#), [ABQ62001.1](#), [ABQ62237.1](#), [ABQ60928.1](#), and [ABQ62635.1](#) with diverse functions (bootstrap >85). (B) Flavoenzymes of subclass EC 1.4. distribute in three clades (bootstrap >90). Clade 1 includes [ABQ61684.1](#) and [ABQ62546.1](#), present in a reduced number of alphaproteobacteria. Clade 2 comprises four proteins distributed in two subclades (bootstrap >88). [ABQ60316.1](#) clusters separately (bootstrap >92), while [ABQ62278.1](#), [ABQ62519.1](#), and [ABQ61937.1](#) are in the same subclade (bootstrap > 77). Clade 3 clusters [ABQ62405.1](#) in a divergent subclade (bootstrap > 80), and [ABQ60142.1](#), [ABQ60524.1](#), [ABQ60616.1](#), and [ABQ61996.1](#), separately in four branches of other subclades (bootstrap > 80). Phylogenetic cladograms include 222 sequences of subclass EC 1.1. and 198 sequences of subclass EC 1.4. from *B. ovis* ATCC 25840, *B. abortus* 2308, and 26 alphaproteobacteria related species. *E. coli* ApbE and *E. coli* GlpD were selected as outgroup in A and B, respectively, to highlight the clear evolutionary separation between clusters. The likelihood aLRT (approximate likelihood-ratio test) statistical test and a bootstrap value of 100 were used.






The subclass EC 1.3 includes seven acyl-CoA dehydrogenases (ACAD) (EC 1.3.8.-) widely represented in *Brucella* but not within alphaproteobacteria (Fig. 5A, Table SP6). The most divergent ACAD ([ABQ62576.1](#)) has an AidB domain instead of an Acyl-CoA\_dh\_N one, and is 100% identical to a protein from *B. melitensis* whose 3D structure is available (PDB [5E23](#), Table SP3, Fig. SP2). It has all features of *B. abortus* and *E. coli* homologues involved in the destruction of alkylating agents, suggesting it will provide resistance during alkylation stress as well as in cell division (47). The ACAD annotated as isovaleryl-CoA dehydrogenase ([ABQ60382.1](#)) in UniProtKB might participate in the catabolism of branched chain amino acids. The other five ACADs could participate in fatty acid  $\beta$ -oxidation (Tables 1–3), suggesting the use for *B. ovis* of lipids, probably recruited from host cells, as carbon sources (7). This subclass also includes the SdhA subunit of succinate dehydrogenase (Sdh) (EC 1.3.5.1) and the UDP-N-acetylmuramate dehydrogenase (MurB) (EC 1.3.1.98), two proteins that are in the core proteome of *Brucella* and conserved in all evaluated alphaproteobacteria (Fig. 5A, Table SP6). The Sdh complex is built by different subunits (Table SP4), localizes in the membrane of many bacteria, and catalyzes the oxidation of succinate to fumarate. It uses membrane quinone as electron acceptor and is the only enzyme linking the tricarboxylic acid cycle and the ETC (48). MurB catalyzes the NADPH dependent reduction of UDP-N-acetylglucosamine-enolpyruvate to UDP-N-acetylmuramic acid (EC 1.3.1.98) and participates in the biosynthesis of peptidoglycan building blocks (49). Subclass EC 1.3 also includes two subunits of the dihydropyrimidine dehydrogenase (EC 1.3.1.1) involved in the  $\beta$ -alanine metabolism, the biosynthesis of pantothenate and CoA, and the pyrimidine nucleotide metabolism; two enzymes involved in the modification of cytoplasmic tRNAs (EC 1.3.1.88, 1.3.1.91); one PyrD dihydroorotate dehydrogenase (EC 1.3.5.2) involved in nucleotide metabolism; and finally one alkene reductase (EC 1.3.1.-/1.7.1.B1) and one KsdD-like steroid dehydrogenase (EC 1.3.99.-) that might, respectively, contribute to toxic compounds degradation and oxidation/dehydrogenation of ketosteroids.

Flavoenzymes of subclass EC 1.4 are highly conserved within *Brucella*, but some are hardly present in alphaproteobacteria (Fig. 4B, Table SP6). They particularly include a pyridoxamine 5'-phosphate oxidase family protein of unclear function present in the *Brucella* core proteome, but only in five of the symbiotic alphaproteobacteria evaluated (*Agrobacterium tumefaciens*, *Mesorhizobium loti*, *Rhizobium etli*, *Rhizobium leguminosarum*, and *Sinorhizobium meliloti*). This suggests that it might be involved in nitrogen metabolism (Tables SP3 and SP6). Its crystal structure, solved for *B. melitensis*, predicts a dimer that might bind either FMN, FAD, or F420 (8-hydroxi-5-deazaflavina) (Fig. SP3). Its genomic context, next to a PhzF family phenazine biosynthesis protein gene (downstream), is pretty similar in homologues. Moreover, in three *Brucella* (*B. melitensis*, *B. canis*, and *B. microti*) a gene for a Nudix hydrolase follows the *phzF* gene, and in the alphaproteobacteria *Ochrobactrum anthropi* these genes are grouped together. Therefore, this protein might be somehow related to the hydrolysis of nucleoside diphosphates linked to other biomolecules. This subclass also includes glycine oxidase (ThiO, EC 1.4.3.19) involved in glycine oxidation to glyoxylate and in the thiamine metabolism; a potential D-amino acid oxidase (EC 1.4.99.-) with low similarity to other characterized enzymes of this type; pyridoxamine 5'-phosphate oxidase (EC 1.4.3.5)

#### FIG 5 Legend (Continued)

[ABQ61769.1](#) together and [ABQ61077.1](#) separately (bootstrap >98). Clade 2 (bootstrap >73) and clade 3 (bootstrap >70) contain, respectively, [ABQ61103.1](#) and [ABQ61413.1](#), given that these proteins are less conserved within studied species. Clade 4 distributes in four subgroups (bootstrap >95) including [ABQ62490.1](#), [ABQ60560.1](#), [ABQ61416.1](#), and [ABQ61966.1](#). Clade 5 includes seven ACAD homologues distributed in four subclades (bootstrap >90). The most divergent contains the entry [ABQ62576.1](#) (bootstrap >98). (B) Flavoreductases of the subclass EC 1.5 distribute in three clades (bootstrap >90). Clade 1 is the most divergent (bootstrap >98) and contains [ABQ60228.1](#). Clade 2 separates in two branches (bootstrap >90) [ABQ61036.1](#) and [ABQ61337.1](#) homologues. Clade 3 distributes in two subclades (bootstrap >85). One subclade includes [ABQ61310.1](#) and [ABQ60177.1](#) grouped in the same branch and [ABQ62932.1](#) in a separate branch, whereas the protein [ABQ60279.1](#) clusters separately. Phylogenetic cladograms include 319 sequences of subclass EC 1.3. and 108 sequences of subclass EC 1.5. from *B. ovis* ATCC 25480, *B. abortus* 2308, and 26 alphaproteobacteria related species. *E. coli* ApbE and *E. coli* GlpD were selected as outgroup in A and B, respectively, to highlight the clear evolutionary separation between clusters. The likelihood aLRT (approximate likelihood-ratio test) statistical test and a bootstrap value of 100 were used.



**FIG 6** Phylogeny of *B. ovis* flavoproteins of subclasses (A) EC 1.8, (B) EC 1.14, and (C) EC 1.16 related to nitrogen metabolism within alphaproteobacteria. (A) Flavoenzymes of subclass EC 1.8. cluster in three clades (bootstrap >90). Clade 1 contains [ABQ61016.1](#). Clade 2 distributes in two subclades organized in

(Continued on next page)

involved in the biosynthesis of pyridoxal 5'-phosphate (50); the two flavoprotein subunits of a glutamate synthase that through three distinct active centers (EC 1.4.1.13) uses of L-glutamine as carbon and nitrogen source during cell growth, particularly within the host as shown by *B. abortus* (51); and two proteins folding in HI0933-like domains that are potential NAD(P)/FAD-dependent dehydrogenases of the aminoacetone oxidase family (EC 1.4.3.-). We cannot unambiguously provide a clear function for these two last proteins, but they might have a particular role in *Brucella* because they are in its core proteome but poorly distributed in alphaproteobacteria (Table SP6).

The subclass EC 1.5 (Fig. 6B) includes the methylenetetrahydrofolate reductase (EC 1.5.1.20), a central enzyme in the carbon fixation and tetrahydrofolate metabolisms; the iron-sulfur flavoprotein ETF-QO (EC 1.5.5.1) that accepts electrons from ETF proteins and contributes to the respiratory chain and OXPHOS pathways (52); the flavin reductase domain containing protein (EC 1.5.1.-) that is the most divergent member and might have flavin reductase activity; and two sarcosine oxidase proteins (EC 1.5.3.1). One sarcosine oxidase (ABQ62932.1) relates to the monomeric soluble protein, whereas the other is composed by two subunits that bind flavins, alpha (ABQ61036.1) and beta (ABQ60177.1+ABQ61310.1), and is the membrane bound form. Sarcosine oxidases would catalyze demethylation of sarcosine as a way for *B. ovis* to grow with sarcosine, ensuring creatinine degradation and glycine, serine, and threonine metabolism. All these flavoproteins of subclass EC 1.5 are in the core proteome of *Brucella*, but are not conserved in alphaproteobacteria (Table SP6).


The subclass EC 1.6 is represented by three flavoenzymes. The WrpA-type FMN-dependent NADH:quinone oxidoreductase (ABQ60884.1, EC 1.6.5.2) is present in all *Brucella* (Table 3 and Table SP6). The *B. abortus* homologue structure (98.5% identity) relates it to NADH:quinone oxidoreductases, but *in vitro* its ability to bind redox cofactors and its oxidoreductase activity have not been proven. However, it modulates *B. abortus* interaction with the mammalian host and is suggested as a new functional class of WrpA/flavodoxin family proteins likely involved in cell survival under acute oxidative stress (53). This subclass also includes the FMN-dependent NADPH dehydrogenase of the old yellow enzyme family (EC 1.6.99.1). It is not present in all *Brucella* and alphaproteobacteria, and despite being highly studied in different species, its acceptor and physiological function remain unclear. Finally, this subclass includes a FAD-dependent NADH dehydrogenase with unknown function (EC 1.6.-.-).

Up to eight flavoenzymes fall in the subclass EC 1.8. The sequence identity of the *B. ovis* FAD dependent glutathione-disulphide reductase (EC 1.8.1.7) with the *S. meliloti* homologue indicates that it must contribute to maintain high levels of reduced glutathione to control redox homeostasis. This agrees with a recent report where disruptions in the gene encoding for it in *B. ovis* produce a significant disadvantage in bacterial growth (54, 55). The three predicted dihydrolipoyl dehydrogenases (ldpA-1, ldpA-2, ldpA-3, EC 1.8.1.4) are highly conserved in *Brucella* and alphaproteobacteria, with the exception of ldpA-1 poorly represented in alphaproteobacteria. Homologues in *B. suis* and *B. abortus* form part of complexes such as the alpha-ketoacid dehydrogenase, pyruvate dehydrogenase, and glycine cleavage multienzyme, implicated in the biosynthesis of Acetyl-CoA and secondary metabolites, oxidative decarboxylation of pyruvate, and glycine metabolism,

## FIG 6 Legend (Continued)

different subgroups that contain ABQ62466.1, ABQ60398.1, and ABQ61458.1. Clade 3 clusters in two subclades (bootstrap >91). The first subclade separates in two branches, WP\_006015257.1 and WP\_006015259.1 together and ABQ62365.1 separately. The second subclade clusters ABQ60123.1 and ABQ61134.1. (B) Flavoenzymes of subclass EC 1.14. and 1.16 distribute in three clades (bootstrap >80). Clade 1 includes the most divergent member, ABQ60249.1, which is the single member of the subclass 1.16. Clade 2 clusters ABQ60137.1, ABQ60978.1, and ABQ60348.1 (bootstrap >90). Clade 3 has three subclades and is more diverse. The most divergent subgroup includes ABQ62030.1 (bootstrap > 80), while ABQ60166.1 and ABQ62553.1 cluster in two branches of the other subclade (bootstrap >96). (C) FMN dependent flavoreductases predicted to act on nitrogenous compounds separately in three clades. Clade 1 has ABQ62091.1, which is barely conserved in alphaproteobacteria. Clade 2 contains ABQ62537.1, and clade 3 separates in two branches, ABQ62404.1 and ABQ60834.1 (bootstrap >75). Phylogenetic cladograms include 177 sequences of class EC 1.8., 160 sequences of class EC 1.14./1.16. and 71 sequences of nitrogen metabolism from *B. ovis* ATCC 25840, *B. abortus* 2308, and 26 alphaproteobacteria related species. *E. coli* GlpD and *E. coli* ApbE were selected as outgroup in A, B, and C, respectively, to highlight the clear evolutionary separation between clusters. The likelihood aLRT (approximate likelihood-ratio test) statistical test and a bootstrap value of 100 were used.





**FIG 7** Metabolic functions for the *B. ovis* flavoproteins. (A) Pie chart distribution of ECs. (B) Number of flavoproteins involved in different metabolic pathways. Blue portions of bars relate to entries predicted to act in a single pathway, whereas orange ones represent entries acting in several pathways.

which contribute to the bacteria pathogenicity (Table SP7) (56–58). The MsrQ subunit of peptide-methionine (S)-S-oxide reductase (quinone) (EC 1.8.5.B1) uses FMN and haem, and is complemented with molybdopterin and quinone at the MsrP subunit. Msr complex is essential for the maintenance of envelope integrity under bleach stress and protects proteins from oxidative-stress damage during host defense mechanisms (59). MsrQ is common to all *Brucella*, but not in alphaproteobacteria. This subclass also has the two flavo-protein subunits of the assimilatory sulfite reductase (NADPH): the alpha component cluster that, together with a beta subunit, catalyzes the six-electron reduction of sulfite to sulfide (EC 1.8.1.2). This protein is usually involved in sulfate and sulfur assimilation, and in microbial metabolism in diverse environments. As mentioned above, it is singular in *B. ovis*, since in other *Brucella* a single protein contains the four components (Fig. 2B). In addition, this subclass has two FAD-dependent thioredoxin-disulphide reductase like proteins (EC 1.8.1.9) with pyridine nucleotide-disulphide oxidoreductase activity potentially involved in the oxidation-reduction cycle of thioredoxin.

Most flavoenzymes of subclass EC 1.14 use NAD(P)H as donor and incorporate oxygen into the second substrate (Fig. 7B). Two of them are related to salicylate hydroxylase activity (EC 1.14.13.1) contributing to the degradation of aromatic compounds, and two are members of the UbiH/COQ6 monooxygenase and UbiH/UbiF hydroxylase families (EC 1.14.13.-) (Table SP6). These latter two enzymes are involved in the ubiquinone biosynthesis pathway (ARBA annotation: [ARBA00004749](#), “ubiquinone biosynthesis”) and share moderate sequence similarity with well-characterized flavoprotein monooxygenases, but close homologues have not been characterized yet (60–62). This subclass also has the 4-hydroxybenzoate 3-monooxygenase (EC 1.14.13.2) that participates in the benzoate degradation and favors microbial metabolism in diverse environments (63, 64). Another member is the bacterial luciferase (EC 1.14.14.3) that incorporates oxygen into reduced FMN to form a peroxyflavin-adduct that upon interaction with aliphatic long-chain aldehydes produces highly fluorescent species. This luciferase is particularly common in symbiotic nitrogen-fixing alphaproteobacteria (*A. tumefaciens*, *R. leguminosarum*, *S. meliloti*, except *R. pomeroyi*).

We have identified a single member of the subclass EC 1.16. This protein is assigned in UniProtKB as FMN-dependent 4-hydroxyphenylacetate 3-monooxygenase (EC 1.14.14.9, [ABQ60249.1](#)). However, it clearly diverges from subclass EC 1.14 (Fig. 6B) and is identical to the Cob(II)alamin reductase (EC 1.16.1.4) of *B. melitensis* with structure and activity experimentally proven (65). Therefore, [ABQ60249.1](#) must participate in the cobalamin (vitamin B<sub>12</sub>) biosynthetic pathway. The single member of subclass EC 1.17 is the small subunit of xanthine dehydrogenase (*xdhA*) (EC 1.17.1.4) that is present in most *Brucella* evaluated. This enzyme participates in the metabolism of purines and is made by several subunits.

The subclass EC 1.18. has two flavoenzymes that exchange electrons between pyridine nucleotides and iron-sulfur proteins. One is the bacterial type ferredoxin-NADP<sup>+</sup> reductase (EC 1.18.1.2), widely distributed in *Brucella* but not in alphaproteobacteria, and for which 3D structure and mechanism of action as NADPH oxidoreductase are reported (39). It probably delivers electrons from NADPH to the redox-based metabolism. But considering that a superoxide dismutase is expressed downstream, it might also oxidize NADPH to activate regulons that protect against oxidative damage (66) (Table SP4). The other member is the ferredoxin/rubredoxin/putidaredoxin NAD<sup>+</sup> reductase (EC 1.18.1.3-5), with also a potential role in cellular oxidative stress response or lipid metabolism.

Finally, four FMN dependent flavoreductases are predicted to act on nitrogenous compounds. The nitroreductase family protein (EC 1.-.-.-) has unknown precise function. The nitronate monooxygenase (EC 1.13.11.32) is predicted to use molecular oxygen to oxidize alkyl nitronates, and to produce enzyme-bound nitronate radicals and peroxyxynitroethane species. The aerobic dimethylbenzimidazole synthase (BluB, EC 1.13.11.79) putatively catalyzes the oxygen-dependent oxidative fragmentation of the reduced isoalloxazine of FMN to yield 5,6-dimethylbenzimidazole in the biosynthesis of cobalamin (67, 68). Finally, it is the NAD(P)H nitroreductase like protein (EC 1.7.-.-) that might oxidize diverse nitrogen-containing compounds (69).

#### Flavoenzymes of the transferase class show varied activities in *B. ovis*.

Flavotransferases (EC 2.) in *B. ovis* use different structural scaffolds and catalyze quite different reactions. Their distribution varies among *Brucella* and alphaproteobacteria species (Table SP6). Two of them fold in GIDA domains and act in the posttranslational modification of tRNAs (2.1) (Table 3). Three others participate in consecutive steps of the biosynthesis of flavin cofactors: the riboflavin synthase (EC 2.5.1.9); the bifunctional FADS with two independent transferase sites (EC 2.7.7.2, EC 2.7.1.26) that has recently been characterized showing species-specific traits in both of its activities (70); and a FAD:protein FMN transferase (EC 2.7.1.180) potentially involved in the transfer of the FMN moiety from FAD to a target flavoprotein. This class includes also the acetolactate synthase 3 (EC 2.2.) that transfers acetaldehyde from one pyruvate to either another pyruvate or 2-oxobutanoate (EC 2.2.1.6) in the respective valine and isoleucine biosynthetic pathways (71); and the blue-light-activated histidine kinase (EC 2.7.13.3). This latter enzyme undergoes photochemistry through

its FMN chromophore by formation of a cysteinyl-flavin adduct that allosterically controls the enzymatic activity at its kinase protein domain (Fig. 3C) (72).

**Lyases and translocases have a minor representation in the flavoproteome of *B. ovis*.** Flavoproteins acting as lyases use the less common folding. One is chorismate synthase (EC. 4.2.3.5) that catalyzes the formation of chorismate, a starting substrate in the biosynthesis of aromatic amino acids (73). The other is a bifunctional enzyme that catalyzes two sequential steps in coenzyme A biosynthesis: the CTP dependent conjugation of cysteine and 4'-phosphopantothenate to form 4-phosphopantothenoylcysteine (EC 6.3.2.5), followed by the FMN-dependent decarboxylation of this product to 4'-phosphopantotheine (EC 4.1.1.36) (74).

Only one flavoprotein of the core proteome of *Brucella* (Table SP6) is predicted as a translocase: the nuoF subunit of the complex I-like NADH quinone oxidoreductase that catalyzes the translocation of protons across membrane linked to a FMN dependent NADH dehydrogenase activity (EC 7.1.1.2) and that contributes therefore to aerobic respiration and OXPHOS (75).

**The *B. ovis* flavoproteome in virulence and infectivity.** In general, *Brucella* spp. do not show aggressive virulence mechanisms such as exotoxins, anti-phagocytic capsules, plasmids, fimbria, flagella, or antigenic variation. Nonetheless, they are highly pathogenic for their preferred or accidental hosts and their silent capacity to adapt to the intracellular environment. They are considered an evolutive virulence factor by themselves (76). In particular, *B. ovis* shows some peculiarities: (i) it does not produce H<sub>2</sub>S, does not hydrolyze urea, and does not reduce nitrate, contrary to most *Brucella* spp., (ii) its lipopolysaccharide protective envelope is naturally rough, and (iii) it is the unique *Brucella* able to oxidize ribitol with the exception of *B. neotomae* (18, 77). The flavoproteome partially contributes to these evolutionary abilities. An example is EryB, particularly present in species that cause abortions, as *B. ovis*, *B. melitensis* and *B. abortus* (78–81).

Thus, the *B. ovis* flavoproteome can be a source of virulence, infectivity, and survival factors, whose distribution varies among the close analyzed species (Tables 3 and Table SP7). Two potential virulent candidates belong to the core proteome of alphaproteobacteria: MurB and SdhA. Nonetheless, despite predictors indicating MurB as a potential virulent/infectivity factor, there is no experimental evidence beyond its essential housekeeping role to maintain the peptidoglycan cell wall (82). On the contrary, SdhA is a requirement for pathogenicity in *E. coli* (48), to stabilize the vacuole integrity during replication in the intracellular pathogen (like *B. ovis*) *Legionella pneumophila* (83), and its reduction is detected early during infection in *B. abortus* (84). Moreover, the *SdhB* gene is virulent in *L. pneumophila* (85), and the SdhB subunit plays a role in filamentation and virulence in *Candida albicans* (86). The integrity of Sdh subunits is also related to antibiotic resistance in *Salmonella enterica* and *Xanthomonas oryzae* pv. *Oryzae* (87, 88). Up to 17 of the flavoproteins predicted as potentially virulent factors in *B. ovis* belong to the core proteome of *Brucella* (or lack in a single species), and are already noticed as involved in the infection process of different pathogens (Table 3; Tables SP6 and SP7). Among them we can highlight the following: (i) Msr that maintains bacterial membrane integrity and contributes to adhesion with eukaryotic cells (59, 89); (ii) two D-amino acid dehydrogenases that could play a pleiotropic role in the production of important virulence factors and support pathological exclusive functions for different isoforms within an organism (90, 91); (iii) one isovaleryl-CoA dehydrogenase involved in vegetative growth, conidiation, and virulence of plant fungal pathogens (92); (iv) one glutamate synthase involved in chronic persistence of *B. abortus* infection in mice (51); (v) one cob(II)alamin reductase conserved in most alphaproteobacteria and whose deletion in *B. abortus* affects the pathogen survival (93); (vi) two tRNA methyltransferases with role in virulence, stress response, growth, and antibiotic susceptibility pathways (94); and (vii) NrdI, essential for the assembly of several subunits of class Ib ribonucleotide reductases expressed under oxidative stress and iron-limited growth conditions (95).

Other predicted virulent factors common in nearly all *Brucella* are the membrane bound sarcosine oxidase, the blue-light-activated histidine kinase, and the pyridoxamine 5'-phosphate oxidase family protein. The blue-light-activated histidine kinase increases its

**TABLE 4** *Brucella ovis* ATCC 25840 flavoproteins lacking homologues in *O. aries* and Mammalia<sup>a</sup>

Flavoprotein	Protein code	Flavoprotein	Protein code
Potential L-gulonolactone oxidase FAD-binding oxygen oxidoreductase	ABQ62001.1	Peptide-methionine (S)-S-oxide reductase (quinone). MsrP catalytic subunit.	ABQ62365.1
UDP-N-acetylmuramate dehydrogenase	ABQ61769.1	Nitronate monooxygenase (formerly 2-nitropropane dioxygenase NPD)	ABQ62537.1
Predicted alkene reductase: N-ethylmaleimide reductase, glycerol trinitrate reductase or xenobiotic reductase B	ABQ62490.1	Predicted aerobic 5,6-dimethylbenzimidazole synthase (BluB)	ABQ62404.1
Predicted KsdD-like steroid dehydrogenase	ABQ62061.1	Predicted Salicylate hydroxylase	ABQ60137.1
Pyridoxamine 5'-phosphate oxidase family protein <sup>b,c</sup>	ABQ61684.1	Predicted Salicylate hydroxylase <sup>b</sup>	ABQ60978.1
Glycine oxidase ThiO	ABQ60316.1	4-hydroxybenzoate 3-monooxygenase	ABQ62030.1
Potential Aminoacetone oxidase family FAD-binding enzyme/ NAD(P)/FAD-dependent dehydrogenase	ABQ60616.1	Bacterial luciferase	ABQ60348.1
Potential Aminoacetone oxidase family FAD-binding enzyme/ NAD(P)/FAD-dependent dehydrogenase	ABQ60524.1	Cob(II)alamin reductase	ABQ60249.1
Predicted D-amino acid dehydrogenase small subunit	ABQ61937.1	Ferredoxin-NADP <sup>+</sup> reductase	ABQ61707.1
Predicted D-amino acid dehydrogenase	ABQ62519.1	Predicted nitroreductase family protein <sup>b</sup>	ABQ62091.1
Flavin reductase domain containing protein	ABQ60228.1	Methylenetetrahydrofolate-tRNA-(uracil54-C5)-methyltransferase NAD(P)H oxidase	ABQ61275.1
Predicted monomeric Sarcosine oxidase	ABQ62932.1	Acetolactate synthase 3 catalytic subunit	ABQ60081.1
WrpA-type FMN-dependent NADH:quinone oxidoreductase	ABQ60884.1	Riboflavin synthase alpha subunit	ABQ605180.1
NADPH dehydrogenase (Old yellow enzyme) <sup>b</sup>	ABQ62422.1	Bifunctional riboflavin kinase/FAD synthase	ABQ62831.1
Predicted NAD(P)H nitroreductase <sup>c</sup>	ABQ60834.1	FAD:protein FMN transferase <sup>b</sup>	ABQ62066.1
Assimilatory sulphite reductase (NADPH) alpha component cluster <sup>c</sup>	WP_006015252.1 WP_006015255.1 WP_006015257.1	Blue-light-activated histidine kinase <sup>b</sup>	ABQ62113.1
		Chorismate synthase	ABQ60200.1
		Protein NrdI <sup>c</sup>	ABQ62891.1
Predicted thioredoxin-disulphide reductase	ABQ61134.1		

<sup>a</sup>Threshold set in at least 30% sequence identity over 50% of the query cover.<sup>b</sup>Proteins underrepresented in alpha-proteobacteria.<sup>c</sup>Proteins without homologues in any Eukarya.

own autophosphorylation to modulate the microorganism virulence in *B. abortus* (72, 96, 97). The pyridoxamine 5'-phosphate oxidase family protein is very rare in other bacteria, but its conservation within *Brucella* suggests a particular still unclear function. In addition, *B. ovis* contains an important number of flavoproteins (some highly conserved in *Brucella*, but not all) whose homologues are required for the survival of different pathogens upon infection by acting in key metabolic pathways and suppressing host defenses (Table SP7).

It is also interesting to note that the *B. ovis* genome contains a large amount of transposable recombinogenic elements and pseudogenes (up to 119 in CI and 125 in CII) that can contribute to its variability and adaptive and evolutionary capacities (98). Many of them sit next to or in flavoenzyme encoding genes (Tables SP4 and SP5). For example, one IS5 transposase is located between the gene encoding the pyridoxamine 5'-phosphate oxidase family protein and the gene encoding for the PhzF family phenazine biosynthesis protein. Other IS5 transposase interrupts the two *SoxB* genes (Fig. 2A, Table SP4), similarly to that reported in the *Pseudomonas aeruginosa* PAO1 and related to a reduction of pathogenicity (99). Regarding pseudogenes, the *B. ovis* xanthine dehydrogenase operon contains a pseudogene instead of a regular *xdhB* encoding region for the corresponding protein subunit (Table SP4). This *XdhB* subunit is not expected to bind flavin, but its lack will make xanthine dehydrogenase not functional (18). The *B. ovis* BOV\_RS11620 gene encoding for the NosR transcriptional regulator of the expression of the nitrous oxide reductase NosZ also has a deletion in its FMN\_bind domain (PF04205). The *nosX* gene, necessary for NosR covalent flavinylation, is in addition a pseudogene (BOV\_RS11650) (100). These features introduce defects in the nitrogen metabolism of *B. ovis*, and contrary to other *Brucella* make it particularly unable for full denitrification and nitrous oxide respiration (101). Thus, degradation of the *B. ovis* flavoproteome surely contributes to narrow its host range and to make it nonzoonotic (18).

#### The *B. ovis* flavoproteome as a source of antimicrobial targets and biocatalyst.

Up to 35 *B. ovis* flavoproteins, most of them belonging to the *Brucella* core proteome, lack homologues in *O. aries* and other mammals (Table 4). This list could be potential

targets for the search for antimicrobials. Some of them are already being explored as targets of inhibitors in other bacteria, as for example, UDP-N-acetylmuramate dehydrogenase (102, 103), riboflavin synthase (104), bifunctional riboflavin kinase/FAD synthase (25, 26), or chorismate synthase (105). In agreement, a comparative metabolomics study in *B. melitensis* also pointed to synthase as an attractive target (106). Others with certain homology to the here-identified thioredoxin-disulphide reductase (107) or FAD:protein FMN transferase are also considered antimicrobial targets (108). Noticeably, Table 4 includes an important number of the *B. ovis* flavoproteins for which the exact physiological function is difficult to envisage. Among them are predicted alkene reductase and KsdD-like steroid dehydrogenase, the pyridoxamine 5'-phosphate oxidase family protein, two potential aminoacetone oxidase family FAD-binding dehydrogenases, two potential D-amino acid dehydrogenases, the predicted nitroreductase family protein, and the NADPH dehydrogenase from the old yellow enzyme family. Of interest, the latter enzyme has been reported to participate in the oxidative stress response and detoxification in *B. subtilis* (109), which points to it as an interesting target to control pathogen survival. Moreover, seven of these *B. ovis* flavoproteins (Table 4) are underrepresented in alphaproteobacteria, suggesting that they might be explored also as potential selective antimicrobial targets. Among them are the pyridoxamine 5'-phosphate oxidase family protein, the NADPH dehydrogenase old yellow enzyme, and the predicted nitroreductase family protein, all of them of still unclear function. This group is completed with the predicted salicylate hydroxylase, the FAD:protein FMN transferase, the blue-light-activated histidine kinase, and the protein NrdI. Considering their above-mentioned envisaged roles for virulence upon infection in different bacteria (Table SP7), these four proteins might be also of particular relevance as drug targets (95, 96).

In addition, Table 4 might also contain flavoenzymes with particular properties for their use in organic synthesis, biocatalysis, and/or bioremediation. Some of them might be predicted nitroreductase family protein or nitronate monooxygenase. If, as envisaged, they contribute to the catabolism of nitroalkanes, widely used in chemical industry and as fuels, their low homology to other family members might provide them with particular stability or mechanistic features that would enlarge their applicative perspectives (110). In any case, before used as either antimicrobial targets or biocatalyst, these flavoproteins should be exhaustively characterized at the structural and functional levels to confirm their relevance for bacteria survival, and investigated for their species-specific features and/or the applicability of the chemical process they catalyze.

## DISCUSSION

The predicted flavoproteins of *B. ovis* are envisaged to catalyze an important number of reactions in a large number of metabolic pathways, being particularly involved in the shuttle of electrons to the bacterial metabolism, the primary and energy metabolism, the metabolism of fats, carbohydrates, proteins, and nucleotides, the oxidative stress response, and the tRNAs methylation (Fig. 7B, Table 3), according to previous reports (9, 12, 15, 111–116). Moreover, the *B. ovis* flavoproteome also contains enzymes that are candidates to favor the microbial metabolism in diverse environments, the xenobiotic metabolism for detoxification of aromatic compounds, the bacterial virulence, or the activation of metabolites (pyruvate, folate, pyridoxal 5'-phosphate, vitamin B2, vitamin B12, etc.). Therefore, flavoproteins and flavoenzymes are implicated in the transformation of a vast variety of metabolic bioactive compounds or are directly involved in suppressing the stress induced by the host cells upon infection, which can make some of these proteins potential targets in the treatment of brucellosis. Noticeably, in *B. ovis*, 55% of 78 predicted flavoproteins belong to the core proteome of *Brucella*, whereas only 18% lack in 25% of the *Brucella* species here evaluated (Table SP6). This indicates a heavy dependence of the *Brucella* metabolism on flavoproteins. Moreover, many of these core flavoproteins are particular to *Brucella*, since very few are found in all alphaproteobacteria evaluated: namely, Sdh, dihydrolipoyl dehydrogenases 2 and 3, thioredoxin-disulphide reductase, and MurB. Nonetheless, the study of



the *B. ovis* flavoproteome also shows some of its members are degraded, and probably unable to be functional, introducing variability in the capacities of this bacteria regarding other members of the genus.

**Concluding remarks.** In the last decades many efforts have been done in sequencing different genomes. Many proteins with undetermined or putative functions have been identified, but so far little has been done to elucidate or corroborate their biological activity. In this context, it is of relevance to predict and evaluate the functionality of candidates for flavoproteins in particular organisms. Here we provide the list of proteins making the flavoproteome in *B. ovis*, as well as data of their potential activities and prevalence in different *Brucella* and alphaproteobacteria species. Several predicted flavoproteins are highly divergent in this genus from revised proteins, and for them is difficult to envisage a clear function. This will probably relate to modified activities or divergent processes and mechanisms still not identified. Based on the compiled information here, we also identify some flavoproteins that might become potential antimicrobial and envisage that others might become new biocatalysts.

## MATERIALS AND METHODS

**Flavoprotein sequence searching.** Sequences for potential flavin-dependent proteins in *Brucella ovis* ATCC 25840 were retrieved from the National Center for Biotechnology Information (NCBI) and UniProtKB databases, and the genome and proteome ensembles of the bacteria. We searched for proteins binding RF, FMN, and FAD as ligands (117, 118). Sequences annotated in other species as consensus motifs for FMN- and FAD-binding were also used as queries to retrieve putative *B. ovis* flavoproteins using BLASTp online tools (7, 119, 120). The identified putative *B. ovis* flavoproteins were analyzed in the context of homologous flavin-dependent proteins reported in bacteria, archaea, eukaryotes, fungi, plants, and mammals.

*B. ovis* flavoprotein orthologues were retrieved in 19 *Brucella* species with complete genome sequenced and in 26 alphaproteobacteria representatives of this bacteria family with complete genome available (Table SP1).

**Flavoprotein classification.** The Pfam database was used to classify the retrieved flavoproteins in families and clans (121). Enzyme Commission numbers (EC numbers), protein names, or metabolic functions were assigned after examining each protein for their information in homologues from different organisms using different databases. These included the BRENDA database (122), the Enzyme Structures database (<http://www.ebi.ac.uk/thornton-srv/databases/enzymes/>), the KEGG PATHWAY database (<https://www.genome.jp/kegg/pathway.html>), the MetaCyc database collection (<https://metacyc.org/>), and the Pathogen-Host Interaction Data Integration Analysis System (Phidias), particularly the *Brucella* Bioinformatics Portal (BBP) containing 17 *Brucella* genomes (123, 124).

Potential virulent protein sequences were predicted using the pipelines of the servers VirulentPred (<http://203.92.44.117/virulent/>), a bi-layer cascade Support Vector Machine (SVM) methods developed for bacterial pathogens, (125) and VICmpred (<https://bio.tools/vicmpred>), specifically designed for Gram-negative bacterial proteins and also predicting general functional class (126).

**Structural modeling.** The BLASTp server was used to obtain identities and to search for the sequences of the most similar proteins with structures in the Protein Data Bank (PDB) (<https://www.rcsb.org/>), and for available 3D structures of flavin-dependent proteins of the *Brucella* genus. 3D structural homology models were built based on templates having at least 35% sequence identity using the Swiss-Model (127) and/or RaptorX (128) servers. Confidence scores calculated to indicate the quality of predicted 3D models were *P* value for the relative global quality, global distance test (GDT), and un-normalized GDT (uGDT) for the absolute global quality.

**Sequence alignment and phylogeny tree.** Sequence alignment and phylogenetic analysis were performed as described (129, 130). For phylogeny analysis the sequence profiles were globally aligned with Clustal Omega (<https://www.ebi.ac.uk/Tools/msa/clustalo/>) (131) and trimmed following the protocol of the TRIMAL software (132). A maximum likelihood phylogenetic tree using the Subtree Pruning and Regrafting (SPR) method was constructed with PHYML (<https://ngphylogeny.fr>) (133). The tree and cladogram were midpoint-rooted and plotted with FigTree (<http://tree.bio.ed.ac.uk/software/figtree/>). The approximate Likelihood-Ratio Test (aLRT) with a seed value of 123456 and bootstrap analyses with a value of 100 were performed. aLRT statistics: 0.022 proportion of invariant.

## SUPPLEMENTAL MATERIAL

Supplemental material is available online only.

**SUPPLEMENTAL FILE 1**, PDF file, 1.2 MB.

## ACKNOWLEDGMENTS

We thank the following grants for funding: PID2019-103901GB-I00 to M.M. by MCIN/AEI/10.13039/501100011033, and LMP13\_21 and E35\_20R to M.M. by the Government of Aragón-FEDER. M.M.-S. thanks CONACYT-México for a predoctoral fellowship.

## REFERENCES

- Massey V. 2000. The chemical and biological versatility of riboflavin. *Biochem Soc Trans* 28:283–296. <https://doi.org/10.1042/bst0280283>.
- Fraaije MW, Mattevi A. 2000. Flavoenzymes: diverse catalysts with recurrent features. *Trends Biochem Sci* 25:126–132. [https://doi.org/10.1016/S0968-0004\(99\)01533-9](https://doi.org/10.1016/S0968-0004(99)01533-9).
- Walsh CT, Wenczewicz TA. 2013. Flavoenzymes: versatile catalysts in biosynthetic pathways. *Nat Prod Rep* 30:175–200. <https://doi.org/10.1039/c2np20069d>.
- Leys D, Scrutton NS. 2016. Sweating the assets of flavin cofactors: new insight of chemical versatility from knowledge of structure and mechanism. *Curr Opin Struct Biol* 41:19–26. <https://doi.org/10.1016/j.sbi.2016.05.014>.
- Schall P, Marutschke L, Grimm B. 2020. The flavoproteome of the model plant *Arabidopsis thaliana*. *Int J Mol Sci* 21:5371. <https://doi.org/10.3390/ijms21155371>.
- Lienhart WD, Gudipati V, Macheroux P. 2013. The human flavoproteome. *Arch Biochem Biophys* 535:150–162. <https://doi.org/10.1016/j.abb.2013.02.015>.
- Macheroux P, Kappes B, Ealick SE. 2011. Flavogenomics—a genomic and structural view of flavin-dependent proteins. *FEBS J* 278:2625–2634. <https://doi.org/10.1111/j.1742-4658.2011.08202.x>.
- Gross E, Kastner DB, Kaiser CA, Fass D. 2004. Structure of Ero1p, source of disulfide bonds for oxidative protein folding in the cell. *Cell* 117:601–610. [https://doi.org/10.1016/S0092-8674\(04\)00418-0](https://doi.org/10.1016/S0092-8674(04)00418-0).
- Nishimasu H, Ishitani R, Yamashita K, Iwashita C, Hirata A, Hori H, Nureki O. 2009. Atomic structure of a folate/FAD-dependent tRNA T54 methyltransferase. *Proc Natl Acad Sci U S A* 106:8180–8185. <https://doi.org/10.1073/pnas.0901330106>.
- Murty CV, Adiga PR. 1982. Pregnancy suppression by active immunization against gestation-specific riboflavin carrier protein. *Science* 216:191–193. <https://doi.org/10.1126/science.7063879>.
- Susin SA, Lorenzo HK, Zamzami N, Marzo I, Snow BE, Brothers GM, Mangion J, Jacotot E, Costantini P, Loeffler M, Larochette N, Goodlett DR, Aebersold R, Siderovski DP, Penninger JM, Kroemer G. 1999. Molecular characterization of mitochondrial apoptosis-inducing factor. *Nature* 397:441–446. <https://doi.org/10.1038/17135>.
- Natarajan SK, Becker DF. 2012. Role of apoptosis-inducing factor, proline dehydrogenase, and NADPH oxidase in apoptosis and oxidative stress. *Cell Health Cytoskeleton* 12:11–27.
- Ferreira P, Villanueva R, Martínez-Júlviz M, Herguedas B, Marcuello C, Fernandez-Silva P, Cabon L, Hermoso JA, Lostao A, Susin SA, Medina M. 2014. Structural insights into the coenzyme mediated monomer-dimer transition of the pro-apoptotic apoptosis inducing factor. *Biochemistry* 53:4204–4215. <https://doi.org/10.1021/bi500343r>.
- Novo N, Ferreira P, Medina M. 2021. The apoptosis-inducing factor family: moonlighting proteins in the crosstalk between mitochondria and nuclei. *IUBMB Life* 73:568–581. <https://doi.org/10.1002/iub.2390>.
- Myllykallio H, Lipowski G, Leduc D, Filee J, Forterre P, Liebl U. 2002. An alternative flavin-dependent mechanism for thymidylate synthesis. *Science* 297:105–107. <https://doi.org/10.1126/science.1072113>.
- Zhang M, Wang L, Zhong D. 2017. Photolyase: dynamics and electron-transfer mechanisms of DNA repair. *Arch Biochem Biophys* 632:158–174. <https://doi.org/10.1016/j.abb.2017.08.007>.
- Spellerberg B, Cundell DR, Sandros J, Pearce BJ, Idanpaan-Heikkilä I, Rosenow C, Masure HR. 1996. Pyruvate oxidase, as a determinant of virulence in *Streptococcus pneumoniae*. *Mol Microbiol* 19:803–813. <https://doi.org/10.1046/j.1365-2958.1996.425954.x>.
- Tsolis RM, Seshadri R, Santos RL, Sangari FJ, Lobo JMG, de Jong MF, Ren Q, Myers G, Brinkac LM, Nelson WC, Deboy RT, Angiuoli S, Khouri H, Dimitrov G, Robinson JR, Mulligan S, Walker RL, Elzer PE, Hassan KA, Paulsen IT. 2009. Genome degradation in *Brucella ovis* corresponds with narrowing of its host range and tissue tropism. *PLoS One* 4:e5519. <https://doi.org/10.1371/journal.pone.0005519>.
- McNeil MB, Hampton HG, Hards KJ, Watson BNJ, Cook GM, Fineran PC. 2014. The succinate dehydrogenase assembly factor, SdhE, is required for the flavinylation and activation of fumarate reductase in bacteria. *FEBS Lett* 588:414–421. <https://doi.org/10.1016/j.febslet.2013.12.019>.
- Dagley S. 1987. Lessons from biodegradation. *Annu Rev Microbiol* 41:1–23. <https://doi.org/10.1146/annurev.mi.41.100187.000245>.
- Meighen EA. 1993. Bacterial bioluminescence: organization, regulation, and application of the lux genes. *FASEB J* 7:1016–1022. <https://doi.org/10.1096/fasebj.7.11.8370470>.
- Briggs WR. 2007. The LOV domain: a chromophore module servicing multiple photoreceptors. *J Biomed Sci* 14:499–504. <https://doi.org/10.1007/s11373-007-9162-6>.
- Hall M. 2020. Flavoenzymes for biocatalysis. *Enzymes* 47:37–62. <https://doi.org/10.1016/bs.enz.2020.05.001>.
- Cremades N, Velázquez-Campoy A, Martínez-Júlviz M, Neira JL, Pérez-Dorado I, Hermoso J, Jiménez P, Lanás A, Hoffman PS, Sancho J. 2009. Discovery of specific flavodoxin inhibitors as potential therapeutic agents against *Helicobacter pylori* infection. *ACS Chem Biol* 4:928–938. <https://doi.org/10.1021/cb900166q>.
- Sebastián M, Anoz-Carbonell E, Gracia B, Cossio P, Ainsa JA, Lans I, Medina M. 2018. Discovery of antimicrobial compounds targeting bacterial type FAD synthetases. *J Enzyme Inhib Med Chem* 33:241–254. <https://doi.org/10.1080/14756366.2017.1411910>.
- Lans I, Anoz-Carbonell E, Palacio-Rodríguez K, Ainsa JA, Medina M, Cossio P. 2020. In silico discovery and biological validation of ligands of FAD synthase, a promising new antimicrobial target. *PLoS Comput Biol* 16:e1007898. <https://doi.org/10.1371/journal.pcbi.1007898>.
- Anoz-Carbonell E, Rivero M, Polo V, Velázquez-Campoy A, Medina M. 2020. Human riboflavin kinase: species-specific traits in the biosynthesis of the FMN cofactor. *FASEB J* 34:10871–10886. <https://doi.org/10.1096/fj.202005566R>.
- Anoz-Carbonell E, Timson DJ, Pey AL, Medina M. 2020. The catalytic cycle of the antioxidant and cancer-associated human NQO1 enzyme: hydride transfer, conformational dynamics and functional cooperativity. *Antioxidants (Basel)* 9:772. <https://doi.org/10.3390/antiox9090772>.
- Martínez-Júlviz M, Goñi G, Pérez-Amigot D, Laplaza R, Ionescu I, Petrocelli S, Tondo M, Sancho J, Orellano E, Medina M. 2017. Identification of inhibitors targeting ferredoxin-NADP(+) reductase from the *Xanthomonas citri* subsp. *citri* phytopathogenic bacteria. *Molecules* 23:29. <https://doi.org/10.3390/molecules23010029>.
- Leone P, Galluccio M, Barbiroli A, Eberini I, Tolomeo M, Vrenna F, Gianazza E, Iametti S, Bonomi F, Indiveri C, Barile M. 2018. Bacterial production, characterization and protein modeling of a novel monofunctional isoform of FAD synthase in humans: an emergency protein? *Molecules* 23:116. <https://doi.org/10.3390/molecules23010116>.
- Leone P, Galluccio M, Quarta S, Anoz-Carbonell E, Medina M, Indiveri C, Barile M. 2019. Mutation of aspartate 238 in FAD synthase isoform 6 increases the specific activity by weakening the FAD binding. *Int J Mol Sci* 20:6203. <https://doi.org/10.3390/ijms20246203>.
- Villanueva R, Romero-Tamayo S, Laplaza R, Martínez-Olivan J, Velázquez-Campoy A, Sancho J, Ferreira P, Medina M. 2019. Redox- and ligand binding-dependent conformational ensembles in the human apoptosis-inducing factor regulate its pro-life and cell death functions. *Antioxid Redox Signal* 30:2013–2029.
- Jortzik E, Wang L, Ma J, Becker K. 2014. Flavins and flavoproteins: applications in medicine. *Methods Mol Biol* 1146:113–157. [https://doi.org/10.1007/978-1-4939-0452-5\\_7](https://doi.org/10.1007/978-1-4939-0452-5_7).
- Gudipati V, Koch K, Lienhart WD, Macheroux P. 2014. The flavoproteome of the yeast *Saccharomyces cerevisiae*. *Biochim Biophys Acta* 1844:535–544. <https://doi.org/10.1016/j.bbapap.2013.12.015>.
- Eggers R, Jammer A, Jha S, Kerschbaumer B, Lahham M, Strandback E, Toplak M, Wallner S, Winkler A, Macheroux P. 2021. The scope of flavin-dependent reactions and processes in the model plant *Arabidopsis thaliana*. *Phytochemistry* 189:112822. <https://doi.org/10.1016/j.phytochem.2021.112822>.
- Picard-Hagen N, Berthelot X, Champion JL, Eon L, Lyazrhi F, Marois M, Peglion M, Schuster A, Trouche C, Garin-Bastuji B. 2015. Contagious epididymitis due to *Brucella ovis*: relationship between sexual function, serology and bacterial shedding in semen. *BMC Vet Res* 11:125. <https://doi.org/10.1186/s12917-015-0440-7>.
- Mirnejad R, Jazi FM, Mostafaei S, Sedighi M. 2017. Molecular investigation of virulence factors of *Brucella melitensis* and *Brucella abortus* strains isolated from clinical and non-clinical samples. *Microb Pathog* 109:8–14. <https://doi.org/10.1016/j.micpath.2017.05.019>.
- Wegrzyn AB, Stolle S, Rienksma RA, Martins Dos Santos VAP, Bakker BM, Suarez-Diez M. 2019. Cofactors revisited—predicting the impact of flavoprotein-related diseases on a genome scale. *Biochim Biophys Acta Mol Basis Dis* 1865:360–370. <https://doi.org/10.1016/j.bbadis.2018.10.021>.
- Pérez-Amigot D, Taleb V, Boneta S, Anoz-Carbonell E, Sebastián M, Velázquez-Campoy A, Polo V, Martínez-Júlviz M, Medina M. 2019. Towards the competent conformation for catalysis in the ferredoxin-NADP+



- reductase from the *Brucella ovis* pathogen. *Biochim Biophys Acta Bioenerg* 1860:148058. <https://doi.org/10.1016/j.bbabi.2019.148058>.
40. Yoo M, Bestel-Corre G, Croux C, Riviere A, Meynial-Salles I, Soucaille P. 2015. A quantitative system-scale characterization of the metabolism of *Clostridium acetobutylicum*. *mBio* 6:e01808-15. <https://doi.org/10.1128/mBio.01808-15>.
  41. Stietz MS, Lopez C, Osifo O, Tolmasky ME, Cardona ST. 2017. Evaluation of the electron transfer flavoprotein as an antibacterial target in *Burkholderia cenocepacia*. *Can J Microbiol* 63:857–863. <https://doi.org/10.1139/cjm-2017-0350>.
  42. Sützl L, Foley G, Gillam EMJ, Bodén M, Haltrich D. 2019. The GMC superfamily of oxidoreductases revisited: analysis and evolution of fungal GMC oxidoreductases. *Biotechnol Biofuels* 12:118. <https://doi.org/10.1186/s13068-019-1457-0>.
  43. Yeh JI, Chinte U, Du S. 2008. Structure of glycerol-3-phosphate dehydrogenase, an essential monotopic membrane enzyme involved in respiration and metabolism. *Proc Natl Acad Sci U S A* 105:3280–3285. <https://doi.org/10.1073/pnas.0712331105>.
  44. Anderson JD, Smith H. 1965. The metabolism of erythritol by *Brucella abortus*. *J Gen Microbiol* 38:109–124. <https://doi.org/10.1099/00221287-38-1-109>.
  45. Meyer ME. 1967. Metabolic characterization of the genus *Brucella*: VI. Growth stimulation by *i*-erythritol compared with strain virulence for guinea pigs. *J Bacteriol* 93:996–1000. <https://doi.org/10.1128/jb.93.3.996-1000.1967>.
  46. Trefzer C, Škovierová H, Buroni S, Bobovská A, Nenci S, Molteni E, Pojer F, Pasca MR, Makarov V, Cole ST, Riccardi G, Mikušová K, Johnsson K. 2012. Benzothiazinones are suicide inhibitors of mycobacterial decaprenyl-phosphoryl- $\beta$ -D-ribofuranose 2'-oxidase DprE1. *J Am Chem Soc* 134:912–915. <https://doi.org/10.1021/ja211042r>.
  47. Dotreppe D, Mullier C, Letesson JJ, De Bolle X. 2011. The alkylation response protein AidB is localized at the new poles and constriction sites in *Brucella abortus*. *BMC Microbiol* 11:257. <https://doi.org/10.1186/1471-2180-11-257>.
  48. Kuo C-J, Wang S-T, Lin C-M, Chiu H-C, Huang C-R, Lee D-Y, Chang G-D, Chou T-C, Chen J-W, Chen C-S. 2018. A multi-omic analysis reveals the role of fumarate in regulating the virulence of enterohemorrhagic *Escherichia coli*. *Cell Death Dis* 9:381. <https://doi.org/10.1038/s41419-018-0423-2>.
  49. Eniyan K, Dharavath S, Vijayan R, Bajpai U, Gourinath S. 2018. Crystal structure of UDP-N-acetylglucosamine-enolpyruvate reductase (MurB) from *Mycobacterium tuberculosis*. *Biochim Biophys Acta Proteins Proteom* 1866:397–406. <https://doi.org/10.1016/j.bbapap.2017.11.013>.
  50. Di Salvo M, Yang E, Zhao G, Winkler ME, Schirch V. 1998. Expression, purification, and characterization of recombinant *Escherichia coli* pyridoxine 5'-phosphate oxidase. *Protein Expr Purif* 13:349–356. <https://doi.org/10.1006/prep.1998.0904>.
  51. Hong PC, Tzolis RM, Ficht TA. 2000. Identification of genes required for chronic persistence of *Brucella abortus* in mice. *Infect Immun* 68:4102–4107. <https://doi.org/10.1128/IAI.68.7.4102-4107.2000>.
  52. Lee MN, Takawira D, Nikolova AP, Ballou DP, Furtado VC, Phung NL, Still BR, Thorstad MK, Tanner JJ, Trimmer EE. 2009. Functional role for the conformationally mobile phenylalanine 223 in the reaction of methylenetetrahydrofolate reductase from *Escherichia coli*. *Biochemistry* 48:7673–7685. <https://doi.org/10.1021/bi9007325>.
  53. Herrou J, Czyż DM, Willett JW, Kim H-S, Chhor G, Babnigg G, Kim Y, Crosson S. 2016. WrpA is an atypical flavodoxin family protein under regulatory control of the *Brucella abortus* general stress response system. *J Bacteriol* 198:1281–1293. <https://doi.org/10.1128/JB.00982-15>.
  54. Tang G, Li N, Liu Y, Yu L, Yan J, Luo L. 2018. *Sinorhizobium meliloti* glutathione reductase is required for both redox homeostasis and symbiosis. *Appl Environ Microbiol* 84:e01937-17. <https://doi.org/10.1128/AEM.01937-17>.
  55. Varesio LM, Fiebig A, Crosson S. 2021. *Brucella ovis* cysteine biosynthesis contributes to peroxide stress survival and fitness in the intracellular niche. *Infect Immun* 89:e00808-20. <https://doi.org/10.1128/IAI.00808-20>.
  56. Rajashankar KR, Bryk R, Kniewel R, Buglino JA, Nathan CF, Lima CD. 2005. Crystal structure and functional analysis of lipamide dehydrogenase from *Mycobacterium tuberculosis*. *J Biol Chem* 280:33977–33983. <https://doi.org/10.1074/jbc.M507466200>.
  57. Abdou E, Jiménez de Bagüés MP, Martínez-Abadía I, Ouahrani-Bettache S, Pantescio V, Occhialini A, Al Dahouk S, Köhler S, Jubier-Maurin V. 2017. RegA plays a key role in oxygen-dependent establishment of persistence and in isocitrate lyase activity, a critical determinant of *in vivo* *Brucella suis* pathogenicity. *Front Cell Infect Microbiol* 7:186. <https://doi.org/10.3389/fcimb.2017.00186>.
  58. Viadas C, Rodríguez MC, Sangari FJ, Gorvel J-P, García-Lobo JM, López-Goñi I. 2010. Transcriptome analysis of the *Brucella abortus* BvrR/BvrS two-component regulatory system. *PLoS One* 5:e10216. <https://doi.org/10.1371/journal.pone.0010216>.
  59. Gennaris A, Ezraty B, Henry C, Agrebi R, Vergnes A, Oheix E, Bos J, Leverrier P, Espinosa L, Szewczyk J, Vertommen D, Iranzo O, Collet J-F, Barras F. 2015. Repairing oxidized proteins in the bacterial envelope using respiratory chain electrons. *Nature* 528:409–412. <https://doi.org/10.1038/nature15764>.
  60. Hajj Chehade M, Loiseau L, Lombard M, Pecqueur L, Ismail A, Smadja M, Gollinelli-Pimpaneau B, Mellot-Draznieks C, Hamelin O, Aussel L, Kieffer-Jaquinod S, Labessan N, Barras F, Fontecave M, Pierrat F. 2013. *ubil*, a new gene in *Escherichia coli* coenzyme Q biosynthesis, is involved in aerobic C5-hydroxylation. *J Biol Chem* 288:20085–20092. <https://doi.org/10.1074/jbc.M113.480368>.
  61. Young IG, Stroobant P, Macdonald CG, Gibson F. 1973. Pathway for ubiquinone biosynthesis in *Escherichia coli* K-12: gene-enzyme relationships and intermediates. *J Bacteriol* 114:42–52. <https://doi.org/10.1128/jb.114.1.42-52.1973>.
  62. Liu L-K, Abdelwahab H, Martin Del Campo JS, Mehra-Chaudhary R, Sobrado P, Tanner JJ. 2016. The structure of the antibiotic deactivating, N-hydroxylating rifampicin monooxygenase. *J Biol Chem* 291:21553–21562. <https://doi.org/10.1074/jbc.M116.745315>.
  63. Wong CM, Dilworth MJ, Glenn AR. 1994. Cloning and sequencing show that 4-hydroxybenzoate hydroxylase (PobA) is required for uptake of 4-hydroxybenzoate in *Rhizobium leguminosarum*. *Microbiology (Reading)* 140:2775–2786. <https://doi.org/10.1099/00221287-140-10-2775>.
  64. Westphal AH, Tischler D, Heinke F, Hofmann S, Grönig JAD, Labudde D, van Berkel WJH. 2018. Pyridine nucleotide coenzyme specificity of *p*-hydroxybenzoate hydroxylase and related flavoprotein monooxygenases. *Front Microbiol* 9:3050. <https://doi.org/10.3389/fmicb.2018.03050>.
  65. Lawrence AD, Deery E, McLean KJ, Munro AW, Pickersgill RW, Rigby SEJ, Warren MJ. 2008. Identification, characterization, and structure/function analysis of a corrin reductase involved in adenosylcobalamin biosynthesis. *J Biol Chem* 283:10813–10821. <https://doi.org/10.1074/jbc.M710431200>.
  66. Krapp AR, Rodriguez RE, Poli HO, Paladini DH, Palatnik JF, Carrillo N. 2002. The flavoenzyme ferredoxin (flavodoxin)-NAD(P)H reductase modulates NAD(P)H homeostasis during the soxRS response of *Escherichia coli*. *J Bacteriol* 184:1474–1480. <https://doi.org/10.1128/JB.184.5.1474-1480.2002>.
  67. Taga ME, Larsen NA, Howard-Jones AR, Walsh CT, Walker GC. 2007. BluB cannibalizes flavin to form the lower ligand of vitamin B12. *Nature* 446:449–453. <https://doi.org/10.1038/nature05611>.
  68. Campbell GRO, Taga ME, Mistry K, Lloret J, Anderson PJ, Roth JR, Walker GC. 2006. *Sinorhizobium meliloti* bluB is necessary for production of 5,6-dimethylbenzimidazole, the lower ligand of B12. *Proc Natl Acad Sci U S A* 103:4634–4639. <https://doi.org/10.1073/pnas.0509384103>.
  69. Hritz J, Zoldák G, Sedláč E. 2006. Cofactor assisted gating mechanism in the active site of NADH oxidase from *Thermus thermophilus*. *Proteins* 64:465–476. <https://doi.org/10.1002/prot.20990>.
  70. Moreno A, Taleb V, Sebastián M, Anoz-Carbonell E, Martínez-Júlvez M, Medina M. 2021. Cofactors and pathogens: FMN and FAD biosynthesis by the FAD synthase from *Brucella ovis*. *IUBMB Life*. <https://doi.org/10.1002/iub.2576>.
  71. Audic S, Lescot M, Claverie JM, Scholz HC. 2009. *Brucella microti*: the genome sequence of an emerging pathogen. *BMC Genomics* 10:352. <https://doi.org/10.1186/1471-2164-10-352>.
  72. Swartz TE, Tseng T-S, Frederickson MA, Paris G, Commerci DJ, Rajashekara G, Kim J-G, Mudgett MB, Splitter GA, Ugalde RA, Goldbaum FA, Briggs WR, Bogomolni RA. 2007. Blue-light-activated histidine kinases: two-component sensors in bacteria. *Science* 317:1090–1093. <https://doi.org/10.1126/science.1144306>.
  73. Foulongne V, Walravens K, Bourg G, Boschirolu ML, Godfroid J, Ramuz M, O'Callaghan D. 2001. Aromatic compound-dependent *Brucella suis* is attenuated in both cultured cells and mouse models. *Infect Immun* 69:547–550. <https://doi.org/10.1128/IAI.69.1.547-550.2001>.
  74. Strauss E, Kinsland C, Ge Y, McLafferty FW, Begley TP. 2001. Phosphopantothienoylcysteine synthetase from *Escherichia coli*: identification and characterization of the last unidentified coenzyme A biosynthetic enzyme in bacteria. *J Biol Chem* 276:13513–13516. <https://doi.org/10.1074/jbc.C100033200>.

75. Schulte M, Frick K, Gndt E, Jurkovic S, Burschel S, Labatzke R, Aierstock K, Fiegen D, Wohlwend D, Gerhardt S, Einsle O, Friedrich T. 2019. A mechanism to prevent production of reactive oxygen species by *Escherichia coli* respiratory complex I. *Nat Commun* 10:2551. <https://doi.org/10.1038/s41467-019-10429-0>.
76. López-Goñi I, Moriyón I. 2004. *Brucella*: molecular and cellular biology. Routledge, Abingdon-on-Thames, England.
77. Ronneau S, Moussa S, Barbier T, Conde-Álvarez R, Zuniga-Ripa A, Moriyon I, Letesson J-J. 2016. *Brucella*, nitrogen and virulence. *Crit Rev Microbiol* 42:507–525. <https://doi.org/10.3109/1040841X.2014.962480>.
78. Sangari FJ, Agüero J, García-Lobo JM. 2000. The genes for erythritol catabolism are organized as an inducible operon in *Brucella abortus*. *Microbiology (Reading)* 146:487–495. <https://doi.org/10.1099/00221287-146-2-487>.
79. Petersen E, Rajashekara G, Sanakkayala N, Eskra L, Harms J, Splitter G. 2013. Erythritol triggers expression of virulence traits in *Brucella melitensis*. *Microbes Infect* 15:440–449. <https://doi.org/10.1016/j.micinf.2013.02.002>.
80. Rodríguez MC, Viadas C, Seoane A, Sangari FJ, López-Goñi I, García-Lobo JM. 2012. Evaluation of the effects of erythritol on gene expression in *Brucella abortus*. *PLoS One* 7:e50876. <https://doi.org/10.1371/journal.pone.0050876>.
81. Barbier T, Collard F, Zúñiga-Ripa A, Moriyón I, Godard T, Becker J, Wittmann C, Van Schaftingen E, Letesson J-J. 2014. Erythritol feeds the pentose phosphate pathway via three new isomerases leading to D-erythrose-4-phosphate in *Brucella*. *Proc Natl Acad Sci U S A* 111:17815–17820. <https://doi.org/10.1073/pnas.1414622111>.
82. Ragunathan A, Malathi K, Anbarasu A. 2018. MurB as a target in an alternative approach to tackle the *Vibrio cholerae* resistance using molecular docking and simulation study. *J Cell Biochem* 119:1726–1732. <https://doi.org/10.1002/jcb.26333>.
83. Creasey EA, Isberg RR. 2012. The protein SdhA maintains the integrity of the Legionella-containing vacuole. *Proc Natl Acad Sci U S A* 109:3481–3486. <https://doi.org/10.1073/pnas.1121286109>.
84. Lamontagne J, Forest A, Marazzo E, Denis F, Butler H, Michaud J-F, Boucher L, Pedro I, Villeneuve A, Sitnikov D, Trudel K, Nassif N, Boudjelti D, Tomaki F, Chaves-Orlante E, Guzmán-Verri C, Brunet S, Côté-Martin A, Hunter J, Moreno E, Paramithiotis E. 2009. Intracellular adaptation of *Brucella abortus*. *J Proteome Res* 1594–1609. <https://doi.org/10.1021/pr800978p>.
85. Brüggemann H, Hagman A, Jules M, Sismeiro O, Dillies M-A, Gouyette C, Kunst F, Steinert M, Heuner K, Coppée J-Y, Buchrieser C. 2006. Virulence strategies for infecting phagocytes deduced from the *in vivo* transcriptional program of *Legionella pneumophila*. *Cell Microbiol* 8:1228–1240. <https://doi.org/10.1111/j.1462-5822.2006.00703.x>.
86. Bi S, Lv Q-Z, Wang T-T, Fuchs BB, Hu D-D, Anastassopoulou CG, Desalermos A, Muhammed M, Wu C-L, Jiang Y-Y, Mylonakis E, Wang Y. 2018. SDH2 is involved in proper hypha formation and virulence in *Candida albicans*. *Future Microbiol* 13:1141–1156. <https://doi.org/10.2217/fmb-2018-0033>.
87. Zhang YJ, Li J, Zhao W, Zhou MG. 2010. A single amino acid substitution in the SdhB protein of succinate dehydrogenase determines resistance to amicarbazol in *Xanthomonas oryzae* pv. *oryzae*. *Pest Manag Sci* 66: 627–633. <https://doi.org/10.1002/pes.1919>.
88. Karsi A, Gülsöy N, Corb E, Dumpala PR, Lawrence ML. 2009. High-throughput bioluminescence-based mutant screening strategy for identification of bacterial virulence genes. *Appl Environ Microbiol* 75: 2166–2175. <https://doi.org/10.1128/AEM.02449-08>.
89. Sharma A, Sharma D, Verma SK. 2017. Proteome wide identification of iron binding proteins of *Xanthomonas translucens* pv. *undulosa*: focus on secretory virulent proteins. *Biometals* 30:127–141. <https://doi.org/10.1007/s10534-017-9991-3>.
90. Oliver KE, Silo-Suh L. 2013. Impact of D-amino acid dehydrogenase on virulence factor production by a pseudomonas aeruginosa. *Can J Microbiol* 59:598–603. <https://doi.org/10.1139/cjm-2013-0289>.
91. Chang YC, Khanal Lamichhane A, Bradley J, Rodgers L, Ngamskulrungraj P, Kwon-Chung KJ. 2015. Differences between *Cryptococcus neoformans* and *Cryptococcus gattii* in the molecular mechanisms governing utilization of D-amino acids as the sole nitrogen source. *PLoS ONE* 10: e0131865. <https://doi.org/10.1371/journal.pone.0131865>.
92. Li Y, Zheng X, Zhu M, Chen M, Zhang S, He F, Chen X, Lv J, Pei M, Zhang Y, Zhang Y, Wang Y, Zhang J, Wang M, Wang X, Li G, Lu G. 2019. Mediated leucine catabolism is required for vegetative growth, conidiation and full virulence of the rice blast fungus. *Front Microbiol* 10:444. <https://doi.org/10.3389/fmicb.2019.00444>.
93. Shim S, Im YB, Jung M, Park WB, Yoo HS. 2018. Genes related to intracellular survival of *Brucella abortus* in THP-1 macrophage cells. *J Microbiol Biotechnol* 28:1736–1748. <https://doi.org/10.4014/jmb.1805.05068>.
94. Farfán-López M, Espinoza-Culupú A, García-de-la-Guarda R, Serral F, Sosa E, Palomino MM, Fernández Do Porto DA. 2020. Prioritisation of potential drug targets against *Bartonella bacilliformis* by an integrative *in-silico* approach. *Mem Inst Oswaldo Cruz* 115:e200184. <https://doi.org/10.1590/0074-02760200184>.
95. Rhodes DV, Crump KE, Makhlynets O, Snyder M, Ge X, Xu P, Stubbe J, Kitten T. 2014. Genetic characterization and role in virulence of the ribonucleotide reductases of *Streptococcus sanguinis*. *J Biol Chem* 289: 6273–6287. <https://doi.org/10.1074/jbc.M113.533620>.
96. Rinaldi J, Fernández I, Shin H, Sycz G, Gunawardana S, Kumarapperuma I, Paz JM, Otero LH, Cerutti ML, Zorreguieta Á, Ren Z, Klinke S, Yang X, Goldbaum FA. 2021. Dimer asymmetry and light activation mechanism in *Brucella* blue-light sensor histidine kinase. *mBio* 12:e00264-21. <https://doi.org/10.1128/mBio.00264-21>.
97. Rinaldi J, Arrar M, Sycz G, Cerutti ML, Berguer PM, Paris G, Estrín DA, Martí MA, Klinke S, Goldbaum FA. 2016. Structural insights into the HWE histidine kinase family: the *Brucella* blue light-activated histidine kinase domain. *J Mol Biol* 428:1165–1179. <https://doi.org/10.1016/j.jmb.2016.01.026>.
98. Bartosik D, Sochacka M, Baj J. 2003. Identification and characterization of transposable elements of *Paracoccus pantotrophus*. *J Bacteriol* 185: 3753–3763. <https://doi.org/10.1128/JB.185.13.3753-3763.2003>.
99. Gallagher LA, Manoil C. 2001. *Pseudomonas aeruginosa* PAO1 kills *Caenorhabditis elegans* by cyanide poisoning. *J Bacteriol* 183:6207–6214. <https://doi.org/10.1128/JB.183.21.6207-6214.2001>.
100. Zhang L, Trncik C, Andrade SLA, Einsle O. 2017. The flavinyl transferase ApbE of *Pseudomonas stutzeri* matures the NosR protein required for nitrous oxide reduction. *Biochim Biophys Acta Bioenerg* 1858:95–102. <https://doi.org/10.1016/j.bbabi.2016.11.008>.
101. Wunsch P, Zumft WG. 2005. Functional domains of NosR, a novel transmembrane iron-sulfur flavoprotein necessary for nitrous oxide respiration. *J Bacteriol* 187:1992–2001. <https://doi.org/10.1128/JB.187.6.1992-2001.2005>.
102. Hrast M, Jukić M, Patin D, Tod J, Dowson CG, Roper DI, Barreteau H, Gobec S. 2018. In silico identification, synthesis and biological evaluation of novel tetrazole inhibitors of MurB. *Chem Biol Drug Des* 91: 1101–1112. <https://doi.org/10.1111/cbdd.13172>.
103. Amera GM, Khan RJ, Pathak A, Jha RK, Jain M, Muthukumaran J, Singh AK. 2020. Structure based drug designing and discovery of promising lead molecules against UDP-N-acetylenolpyruvoylglucosamine reductase (MurB): a potential drug target in multi-drug resistant *Acinetobacter baumannii*. *J Mol Graph Model* 100:107675. <https://doi.org/10.1016/j.jmgm.2020.107675>.
104. Serer MI, Carica MDC, Trappe J, López Romero S, Bonomi HR, Klinke S, Cerutti ML, Goldbaum FA. 2019. A high-throughput screening for inhibitors of riboflavin synthase identifies novel antimicrobial compounds to treat brucellosis. *FEBS J* 286:2522–2535. <https://doi.org/10.1111/febs.14829>.
105. Bueno PS, Rodrigues-Vendramini FA, Toplak M, Macheroux P, Kioshima ES, Seixas FA. 2019. New inhibitors of chorismate synthase present anti-plagial activity against. *Future Microbiol* 14:969–980. <https://doi.org/10.2217/fmb-2019-0052>.
106. Maskar A, Meshram R. 2013. Homology modeling of chorismate synthase from *Brucella melitensis*: a novel target molecule. *Res and Rev: J Microbiology and Biotechnology* 2:7–18.
107. O'Loughlin J, Napolitano S, Alkhathami F, O'Beirne C, Marhöfer D, O'Shaughnessy M, Howe O, Tacke M, Rubini M. 2021. The antibacterial drug candidate SBC3 is a potent inhibitor of bacterial thioredoxin reductase. *Chembiochem* 22:1093–1098. <https://doi.org/10.1002/cbic.202000707>.
108. Deka RK, Deka A, Liu WZ, Norgard MV, Brautigam CA. 2022. Inhibition of bacterial FMN transferase: A potential avenue for countering antimicrobial resistance. *Protein Sci* 31:545–551. <https://doi.org/10.1002/pro.4241>.
109. Fitzpatrick TB, Amrhein N, Macheroux P. 2003. Characterization of YqjM, an Old Yellow Enzyme homolog from *Bacillus subtilis* involved in the oxidative stress response. *J Biol Chem* 278:19891–19897. <https://doi.org/10.1074/jbc.M211778200>.
110. Torres-Guzman JC, Padilla-Guerrero IE, Cervantes-Quintero KY, Martinez-Vazquez A, Ibarra-Guzman M, Gonzalez-Hernandez GA. 2021. Peculiarities of nitronate monooxygenases and perspectives for in vivo and in vitro applications. *Appl Microbiol Biotechnol* 105:8019–8032. <https://doi.org/10.1007/s00253-021-11623-1>.

111. Parsons HG, Dias VC. 1991. Intramitochondrial fatty acid metabolism: riboflavin deficiency and energy production. *Biochem Cell Biol* 69:490–497. <https://doi.org/10.1139/o91-073>.
112. Bornemann S. 2002. Flavoenzymes that catalyse reactions with no net redox change. *Nat Prod Rep* 19:761–772. <https://doi.org/10.1039/b108916c>.
113. Mansoorabadi SO, Thibodeaux CJ, Liu HW. 2007. The diverse roles of flavin coenzymes—nature's most versatile thespians. *J Org Chem* 72:6329–6342. <https://doi.org/10.1021/jo0703092>.
114. Lipton SA, Bossy-Wetzel E. 2002. Dueling activities of AIF in cell death versus survival: DNA binding and redox activity. *Cell* 111:147–150. [https://doi.org/10.1016/s0092-8674\(02\)01046-2](https://doi.org/10.1016/s0092-8674(02)01046-2).
115. Pinto JT, Cooper AJ. 2014. From cholesterologenesis to steroidogenesis: role of riboflavin and flavoenzymes in the biosynthesis of vitamin D. *Adv Nutr* 5:144–163. <https://doi.org/10.3945/an.113.005181>.
116. Romero E, Gómez Castellanos JR, Gadda G, Fraaije MW, Mattevi A. 2018. Same substrate, many reactions: oxygen activation in flavoenzymes. *Chem Rev* 118:1742–1769. <https://doi.org/10.1021/acs.chemrev.7b00650>.
117. Benson DA, Cavanaugh M, Clark K, Karsch-Mizrachi I, Lipman DJ, Ostell J, Sayers EW. 2013. GenBank. *Nucleic Acids Res* 41:D36–D42. <https://doi.org/10.1093/nar/gks1195>.
118. UniProt Consortium. 2021. UniProt: the universal protein knowledge-base in 2021. *Nucleic Acids Res* 49:D480–D489.
119. Gish W, States DJ. 1993. Identification of protein coding regions by database similarity search. *Nat Genet* 3:266–272. <https://doi.org/10.1038/ng0393-266>.
120. Dym O, Eisenberg D. 2001. Sequence-structure analysis of FAD-containing proteins. *Protein Sci* 10:1712–1728. <https://doi.org/10.1110/ps.12801>.
121. El-Gebali S, Mistry J, Bateman A, Eddy SR, Luciani A, Potter SC, Qureshi M, Richardson LJ, Salazar GA, Smart A, Sonhammer ELL, Hirsh L, Paladin L, Piovesan D, Tosatto SCE, Finn RD. 2019. The Pfam protein families database in 2019. *Nucleic Acids Res* 47:D427–D432. <https://doi.org/10.1093/nar/gky995>.
122. Jeske L, Placzek S, Schomburg I, Chang A, Schomburg D. 2019. BRENDA in 2019: a European ELIXIR core data resource. *Nucleic Acids Res* 47:D542–D549. <https://doi.org/10.1093/nar/gky1048>.
123. Xiang Z, Tian Y, He Y. 2007. PHIDIAS: a pathogen-host interaction data integration and analysis system. *Genome Biol* 8:R150. <https://doi.org/10.1186/gb-2007-8-7-r150>.
124. Xiang Z, Zheng W, He Y. 2006. BBP: *Brucella* genome annotation with literature mining and curation. *BMC Bioinformatics* 7:347. <https://doi.org/10.1186/1471-2105-7-347>.
125. Garg A, Gupta D. 2008. VirulentPred: a SVM based prediction method for virulent proteins in bacterial pathogens. *BMC Bioinformatics* 9:62. <https://doi.org/10.1186/1471-2105-9-62>.
126. Waseem H, Williams MR, Stedtfeld T, Chai B, Stedtfeld RD, Cole JR, Tiedje JM, Hashsham SA. 2017. Virulence factor activity relationships (VFARs): a bioinformatics perspective. *Environ Sci Process Impacts* 19:247–260. <https://doi.org/10.1039/c6em00689b>.
127. Biasini M, Bienert S, Waterhouse A, Arnold K, Studer G, Schmidt T, Kiefer F, Gallo Cassarino T, Bertoni M, Bordoli L, Schwede T. 2014. SWISS-MODEL: modelling protein tertiary and quaternary structure using evolutionary information. *Nucleic Acids Res* 42:W252–W258. <https://doi.org/10.1093/nar/gku340>.
128. Källberg M, Wang H, Wang S, Peng J, Wang Z, Lu H, Xu J. 2012. Template-based protein structure modeling using the RaptorX web server. *Nat Protoc* 7:1511–1522. <https://doi.org/10.1038/nprot.2012.085>.
129. Becana M, Yruea I, Sarath G, Catalán P, Hargrove MS. 2020. Plant hemoglobins: a journey from unicellular green algae to vascular plants. *New Phytol* 227:1618–1635. <https://doi.org/10.1111/nph.16444>.
130. Yruea I, Moreno-Yruea C, Olsen CA. 2021. Zn<sup>2+</sup>-dependent histone deacetylases in plants: structure and evolution. *Trends Plant Sci* 26:741–757. <https://doi.org/10.1016/j.tplants.2020.12.011>.
131. Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H, Remmert M, Söding J, Thompson JD, Higgins DG. 2011. Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Mol Syst Biol* 7:539. <https://doi.org/10.1038/msb.2011.75>.
132. Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. 2009. trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* 25:1972–1973. <https://doi.org/10.1093/bioinformatics/btp348>.
133. Lemoine F, Correia D, Lefort V, Doppelt-Azeroual O, Mareuil F, Cohen-Boulakia S, Gascuel O. 2019. NGPhylogeny.fr: new generation phylogenetic services for non-specialists. *Nucleic Acids Res* 47:W260–W265. <https://doi.org/10.1093/nar/gkz303>.
134. Delano WL. 2002. PyMOL: an open-source molecular graphics tool. *CCP4 Newsletter on Protein Crystallography* 40:82–92.