

Title: Assessment of serotonin in serum, plasma and platelets of aggressive dogs

Running title: Blood serotonin in aggressive dogs

Marta León, DVM, PhD^a, Belén Rosado, DVM, PhD^{b*}, Sylvia García-Belenguer, DVM, PhD,
ECVBM-CA^b, Gema Chacón, DVM, PhD^b, Ainara Villegas, MSc^b, Jorge Palacio, DVM, PhD,
ECVBM-CA^b

^a Laboratorios Merial S.A., Servicio Técnico-Animales de Compañía, Tarragona 161 D/E,
08014 Barcelona, Spain.

^b Departamento de Patología Animal. Facultad de Veterinaria, Universidad de Zaragoza, Miguel
Servet, 177, 50013 Zaragoza, Spain.

*Corresponding author (Belén Rosado): belen@unizar.es

Tel.: +0034 976 76 16 00 Ext. 4108

Fax: +0034 976 76 16 12

Other e-mail addresses:

Marta León: marta.leon@merial.com

Sylvia García-Belenguer: sgarcia@unizar.es

Gema Chacón gchacon@unizar.es

Ainara Villegas villegas@unizar.es

Jorge Palacio: jpalacio@unizar.es

Abstract

Canine aggression is the most common reason for the referral of dogs to behavior practices. In addition, dog bites represent an important problem for public health and animal welfare. The serotonergic system is believed to play an important role in modulating aggression. The aim of the present study was (1) to assess the suitability of different types of blood samples for measuring circulating serotonin in canine clinical studies, and (2) to investigate the relationship between the serotonergic system and canine aggression. The assessment of serotonin was simultaneously carried out in serum, plasma and platelets of 28 aggressive and 10 non-aggressive dogs with an enzymeimmunoassay technique. The mean serotonin concentration in aggressive dogs was significantly lower than in non-aggressive dogs in all the assayed samples. These findings suggest an inverse relationship between the activity of the serotonergic system and canine aggression. Considering the simplicity of the methodology, the authors propose sampling serum as the most suitable method for measuring circulating serotonin in dogs.

Keywords: serotonin, serum, plasma, platelet, dog, aggression

Introduction

Canine aggression is the most common reason for the referral of dogs to animal behaviorists (Bamberger and Houpt, 2006; Fatjó et al., 2007). Dog bites represent an important problem for public health (Overall and Love, 2001; Palacio et al., 2005). They also pose a problem in the field of animal welfare since a significant proportion of aggressive dogs is euthanized or abandoned (Hunthausen, 1997; Mikkelsen and Lund, 2000). If these problems are to be prevented and effectively managed, it is necessary to deepen in the understanding of the biological basis of canine aggression.

Serotonergic neurotransmission is believed to play a critical role in the control of aggression in several species (Ferrari et al., 2005). Low concentrations of the main serotonin (5-HT) metabolite 5-hydroxyindoleacetic acid (5-HIAA) in cerebrospinal fluid (CSF) have been associated with low thresholds for aggression or impulsivity in human and non-human primates (Mehlman et al., 1994; Stanley et al., 2000; Soderstrom et al., 2001; Howell et al., 2007) as well as in dogs (Reisner et al., 1996).

Technical and ethical restrictions linked to the study of the serotonergic system in psychiatric disorders through CSF and neurohistochemical studies have led scientists to search for more easily accessible biological samples. The platelet where blood 5-HT is mainly stored (Kema et al., 2000), has been proposed as a peripheral marker of the serotonergic neuron (Plein and Berk, 2001). Albeit blood 5-HT does not cross the hematoencephalic barrier, a correlation between blood and CSF serotonergic parameters have been found in humans (Sarrias et al., 1990). In the realm of human aggression, the measurement of blood 5-HT has offered inconclusive results since both low (Goveas et al., 2004) and high (Golubchik et al., 2009) concentrations have been found to be linked to aggressive behavior in different populations. In canine species, two recent studies have reported low serum serotonin concentrations in aggressive individuals (Çakiroglu et al., 2007; Rosado et al., 2010a).

The aim of the present study was two-fold. First, to assess the suitability of different types of blood samples for measuring circulating 5-HT in clinical studies in canine species. Second, to further investigate the relationship between the serotonergic system and canine aggression. To

this end, the assessment of 5-HT was simultaneously carried out in serum, plasma and platelets of aggressive and non-aggressive dogs.

Materials and methods

Aggressive animals

Two Spanish veterinary teaching hospitals (Universidad de Zaragoza and Cardenal Herrera-CEU, Valencia) contributed to the collection of cases. Dogs included in the present study were referred to the Companion Animal Behavior Services within the respective hospitals owing to problems of aggression towards people and/or other dogs. Displaying any other type of behavioral problem did not constitute an exclusion criterion. Animals showing an underlying causative medical condition to the problem of aggression were excluded from the study.

Twenty males (17 entire and 3 castrated) and 8 females (7 entire and one spayed) were included in the aggressive group. The mean age was 4.2 years old (ranging from 7 months to 13 years). The group consisted of 17 dogs of different breeds (four English cocker spaniels, two German shepherds, two Rottweilers, one Alaskan malamute, one American pit bull terrier, one English bull terrier, one Catalan sheepdog, one Lhasa Apso, one Poodle, one Teckel, one West Highland white terrier and one Yorkshire terrier) and 11 cross-breeds and mongrels.

Diagnosis of aggression was carried out by means of a detailed standard questionnaire on the dogs' behavior and daily routine. Clinical classification of aggression was established in accordance with three main diagnostic criteria: target, context and dog's communicative signals (based on Fatjó et al., 2007). In order to detect any underlying causative or contributory medical condition to the aggression problem, all dogs were screened through physical examination, complete blood count, serum biochemistry and thyroid hormone measurement at the time of admission.

According to the main diagnosis, two animals showed intraspecific aggression whereas the rest (n=26, 93%) showed aggression towards humans, mainly directed to the family members during conflict or competitive situations (69.2%). Several forms of aggression were present in 39.3% of the animals.

Control animals

The control group was made up of 10 beagles (5 entire males, 3 entire females and 2 spayed females) owned by the Universidad Cardenal Herrera-CEU for research purposes. The mean age was 3.1 years old (ranging from 1 to 6 years). All these dogs were healthy and lacked any history of aggression towards people and/or other dogs. The care and use of animals followed the European guidelines (European Union Directive 86/609/EEC, 1986).

Sample collection and preparation

Blood drawn (3 ml) into EDTA tubes was centrifuged at room temperature for 12 min at 152xg to obtain platelet-rich plasma (PRP). The PRP-supernatant was then transferred to another tube for platelet pellet preparation (according to the protocol by IBL, 1998; see below). The EDTA tubes were then centrifuged at 4500xg for 10 min at 4°C to obtain plasma. Blood drawn (3 ml) into anticoagulant-free tubes was centrifuged at 4500xg for 10 min at 4°C to obtain serum. Aliquots of plasma and serum were frozen and stored at –80°C until the day of the analysis.

Platelet count was carried out in PRP (20 µL) using a Neubauer-counting chamber. To obtain the platelet pellet, an aliquot of 200 µL of PRP was added to 800 µL of physiological saline and centrifuged at 4500xg for 10 min at 4°C. The supernatant was discarded and 200 µL of double-distilled water was then added to the pellet. Aliquots were stored at –80°C. The day of the analysis the samples were centrifuged at 10,000xg for 2 min at room temperature after thawing of the frozen aliquots.

Serotonin analysis

Serotonin was measured in duplicate with a commercial EIA technique (Serotonin-ELISA, DLD Diagnostika GmbH, Hamburg, Germany). The intra and interassay coefficients of variations were 3.9-5.4% and 6%, respectively. Concentrations of 5-HT were expressed in ng/mL (serum and plasma) and ng/10⁸ platelets (platelets).

Statistical analysis

A unifactorial multivariate analysis of variance was carried out to assess the effect of the factor “aggression” on the concentrations of 5-HT in the three types of samples, namely serum, plasma and platelets. Correlations were analyzed using the Pearson test.

Calculations were carried out using the statistical program SPSS 14.0. for Windows (SPSS, Inc, Chicago, USA). $P < 0.05$ denoted statistical significance.

Results

The unifactorial multivariate analysis of variance showed a significant effect of the factor “aggression” on 5-HT concentrations ($P < 0.01$). Aggressive animals showed lower 5-HT concentrations in serum, plasma and platelets than non-aggressive animals. The higher mean 5-HT concentration was detected in serum samples. Serotonin results according to the type of sample are summarized in Table 1.

Serotonin concentration in platelets was positively correlated with that in serum ($r = 0.465$; $P < 0.01$) and plasma ($r = 0.468$; $P < 0.01$). In addition, 5-HT concentrations in serum and plasma were positively correlated ($r = 0.489$; $P < 0.01$).

Discussion

In the present study, the assessment of 5-HT in serum, plasma and platelets was carried out both in aggressive and non-aggressive dogs. Regardless of the type of sample, aggressive dogs showed significantly lower concentrations of circulating 5-HT than non-aggressive dogs. This finding agrees with previous studies that found an association between high aggression or impulsivity and low concentrations of CSF 5-HIAA and circulating 5-HT in primates (CSF: Melhman et al., 1994; Stanley et al., 2000; Soderstrom et al., 2001; Howell et al., 2007; platelets: Goveas et al., 2004) as well as in dogs (CSF: Reisner et al., 1996; serum: Çakiroglu et al., 2007; Rosado et al., 2010a).

Several mechanisms regulate the functioning of the 5-HT system in the brain: (i) 5-HT synthesis from tryptophan and degradation, (ii) 5-HT re-uptake from synaptic cleft, and (iii) the density and sensitivity of 5-HT receptors (Popova, 2006). An alteration in any of these mechanisms may be the underlying reason for the reduced activity of the serotonergic system in aggression. Regarding canine aggression, an elevated plasma concentration of tryptophan was detected in a group of aggressive dogs in comparison to a non-aggressive group (Juhr et al., 2005). In addition, the administration of tryptophan-supplemented low-protein diets has shown to be effective in reducing dominance aggression in dogs (DeNapoli et al., 2000). Moreover, a recent

study by the authors found a higher platelet 5-HT uptake in aggressive dogs than in non-aggressive ones (Rosado et al., 2010b). Finally, a series of brain-imaging studies showed an increase in the uptake of the 5-HT_{2A} radioligand in all cortical areas of impulsive-aggressive dogs compared with normal dogs (Peremans et al., 2002, 2003 and 2005). All of these findings point towards an altered function of the serotonergic system in canine aggression.

In the present study, the mean concentration of 5-HT in the assayed samples reflected the normal distribution of blood 5-HT, with the higher concentration in serum, followed by platelets and finally, plasma. Circulating plasma 5-HT is taken up by platelets, and is accumulated during the platelet's life cycle in dense granules. Thus, under normal conditions, 5-HT is almost entirely confined in platelets whereas its concentration in plasma and other blood cells would be very low (Ortiz et al., 1988; Kema et al., 2000). Mück-Seler et al. (2002) have suggested that serum 5-HT concentration might indirectly represent a sum of platelet and plasma 5-HT concentration. In the present study, the sum of the mean (estimated) platelet and plasma 5-HT concentrations was similar, although not exact, to the mean serum 5-HT concentrations in each group of dogs.

Platelet and plasma 5-HT concentrations in the control group were similar to those reported by several studies on canine species (platelets: Meyers et al., 1982; Aranda et al., 1994; plasma: Haga et al., 1996). Likewise, serum 5-HT concentration was also similar to previously published data from whole blood in dogs (Ferrara et al., 1987; LaRosa et al., 1989; Chen et al., 1993). It is worth mentioning, however, the striking difference in serum 5-HT concentrations in the present study as compared with those in the study by Çakiroglu et al. (2007), where relatively lower concentrations were detected both in non-aggressive and aggressive dogs. This difference may be explained in terms of methodological differences in measuring serum 5-HT (e.g., the use of different ELISA-kits).

A statistically significant difference in 5-HT concentration between aggressive and non-aggressive dogs was noted both in serum and platelets as well as in plasma samples. This indicates that any of them may represent a valid blood sample for measuring circulating 5-HT in order to detect differences between both groups. The positive correlation found among the three

types of samples may also support this suggestion. The choice of either one of these different types of blood samples for clinical applications, however, deserves some comments. Plasma is easy to obtain but as previously mentioned 5-HT concentrations in this pool are low, and therefore, more sensitive techniques are required to accurately detect the amine (e.g., high performance liquid chromatography). In addition, plasma may be potentially contaminated with platelet 5-HT (Anderson et al., 1987). Lastly, platelet preparation is an arduous task as an additional platelet count is necessary to express the concentration of 5-HT apart from blood centrifugation to obtain PRP. Since serum 5-HT concentration is considered an indirect measure of platelet 5-HT content, it could be suggested that serum is the easiest sample type to assess circulating 5-HT in dogs. Nevertheless, the assessment of plasma 5-HT may be of great interest also in the field of research on the serotonergic system in canine aggression. It is suggested that plasma 5-HT is a rapid turnover pool, affected by short-term changes in the 5-HT physiology (e.g., synthesis, inactivation, uptake, etc.) while platelet 5-HT is a reserve, slow turnover pool influenced by sustained changes. Thus, the measurement of plasma or platelet (serum) 5-HT concentrations would be useful to study different aspects of 5-HT physiology (Ortiz et al., 1988).

One limitation to the present work is the use of an experimental cohort of beagles as control subjects for a group of aggressive pet dogs belonging to different breeds. In spite of their regular contact with humans, the rearing and environmental conditions in the beagles group clearly differed from those dogs in the pet group, and this may have affected the activity of the 5-HT system even early in life. In addition, it is possible that breed-related differences may occur in the 5-HT system functioning. It is important to note however that the use of experimental beagles as a control group might provide some advantages such as low individual variability (high group homogeneity) and, therefore, high reproducibility of the experiment in other laboratory settings. Another limitation of the study is that the two groups of study were not matched for sex and reproductive status, with proportionally more entire males in the aggressive than the control group, and there is evidence that testosterone (but also estradiol) interacts with the 5-HT system to influence the likelihood of aggression, probably by modulating 5-HT

receptors (Nelson and Chiavegatto, 2001; Soma et al., 2008). In another published study with a larger sample of dogs, however, we did not find a significant effect of sex on serum 5-HT levels (Rosado et al., 2010a). Finally, it is important to note that the above mentioned limitations would have not affected the first aim of the present study, that is: to assess and compare the suitability of different types of blood samples for measuring circulating 5-HT in dogs. Anyway, future studies on this field should consider breed, sex (and reproductive status) as well as age-matched groups in order to assess these possible influencing factors.

In conclusion, the present results may suggest an inverse relationship between canine aggression and the activity of the serotonergic system. To the authors' best of knowledge, this is the first study that simultaneously assesses the concentration of 5-HT in serum, plasma and platelets in aggressive and non-aggressive dogs. The three assayed blood samples showed differences between both groups, but considering the simplicity of the methodology, the authors propose serum as the more suitable sample for measuring circulating 5-HT in clinical studies in dogs. The determination of circulating 5-HT may have important applications for behavior medicine in the future. For example, it could be used for deciding which aggressive dogs might benefit from drug therapy and for monitoring the response to these therapies. In addition, it could be used for research purposes with the aim of creating new tools for diagnosis and treatment of canine aggression.

Acknowledgement

The authors are grateful to María Yetano for language support.

References

Anderson, G.M., Feibel, F.C., Cohen, D.J., 1987. Determination of serotonin in whole blood, platelet-rich plasma, platelet-poor plasma and plasma ultrafiltrate. *Life Sci.* 40, 1063-1070.

228 Aranda, E., Pizarro, M., Pereira, J., Mezzano, D., 1994. Accumulation of 5-hydroxytryptamine
 229 by aging platelets: studies in a model of suppressed thrombopoiesis in dogs. *Thrombosis and*
 230 *Haemostasis* 71, 488-492.

231 Bamberger, M., Houpt, K.A., 2006. Signalment factors, comorbidity, and trends in behavior
 232 diagnoses in dogs: 1,644 cases (1991-2001). *J Am. Vet. Med. Assoc.* 229, 1591-1601.

233 Çakiroglu, D., Meral, Y., Sancak, A.A., Cifti, G., 2007. Relationship between the serum
 234 concentrations of serotonin and lipids and aggression in dogs. *Vet. Rec.* 161, 59-61.

235 Chen, Y.K., Richter, H.M. 3rd, Go, V.L., Tyce, G.M., 1993. Free and conjugated catecholamines
 236 and serotonin in canine thoracic duct lymph: effects of feeding. *Am. J. Physiol.* 265, 84-89.

237 DeNapoli, J.S., Dodman, N.H., Shuster, L., Rand, W.M., Gross, K.L., 2000. Effect of dietary
 238 protein content and tryptophan supplementation on dominance aggression, territorial aggression,
 239 and hyperactivity in dogs. *J. Am. Vet. Med. Assoc.* 217, 504-508.

240 European Union Directive 86/609/EEC (1986) Council Directive of 24 November 1986 on the
 241 approximation of laws, regulations and administrative provisions of the Member States
 242 regarding the protection of animals used for experimental and other scientific purposes. *Official*
 243 *Journal of the European Communities Serie L358*, 132.

244 Fatjó, J., Amat, M., Mariotti, V., Ruiz de la Torre, J.L., Manteca, X., 2007. Analysis of 1040
 245 cases of canine aggression in a referral practice in Spain. *JVB:CAR* 2, 158-165.

246 Ferrara, A., Zinner, M.J., Jaffe, B.M., 1987. Intraluminal release of serotonin, substance P, and
 247 gastrin in the canine small intestines. *Dig. Dis. Sci.* 32, 289-294.

248 Ferrari, P.F., Palanza, P., Parmigiani, S., de Almeida, R.M.M., Miczek, K.A., 2005. Serotonin
 249 and aggressive behavior in rodents and nonhuman primates: Predispositions and plasticity. *Eur.*
 250 *J. Pharmacol.* 526, 259-273.

251 Golubchik, P., Mozes, T., Vered, Y., Weizman, A., 2009. Platelet poor plasma serotonin level
 252 in delinquent adolescents diagnosed with conduct disorder. *Prog Neuropsychopharmacol*
 253 *Biol Psychiatry* 33, 1223-1225

254 Goveas, J.S., Csernansky, J.G., Coccaro, E.F., 2004. Platelet serotonin content correlates
 255 inversely with life history of aggression in personality-disordered subjects. *Psychiatry Res.* 126,
 256 23-32.

257 Haga, N., Mizumoto, A., Satoh, M., Mochiki, E., Mizusawa, F., Ohshima, K., et al., 1996. Role
 258 of endogenous 5-hydroxytryptamine in the regulation of gastric contractions by motilin in dogs.
 259 *Am. J. Physiol.* 270, G20-28.

260 Howell, S., Westergaard, G., Hoos, B., Chavanne, T.J., Shoaf, S.E., Clevelan, A., et al., 2007.
 261 Serotonergic influences on life-history outcomes in free-ranging mal rhesus macaques. *Am. J.*
 262 *Primatol.* 69, 851-865.

263 Hunthausen, W., 1997. Effects of aggressive behavior on canine welfare. *J. Am. Vet. Med.*
 264 *Assoc.* 210, 1134-1136.

265 IBL GmbH (hamburg, Germany), 1998. Instructions for use Serotonin-ELISA (RE59121).

266 Juhr, N.C., Brand, U., Riedel, E., 2005. Plasma amino acid concentrations in aggressive dogs.
 267 *Berl. Munch Tierarztl. Wochenschr.* 118, 95-100.

268 Kema, I.P., de Vries, E.G., Muskiet, F.A., 2000. Clinical chemistry of serotonin and
 269 metabolites. *J. Chromatogr. B. Biomed. Sci. Appl.* 747, 33-48.

270 LaRosa, C.A., Sherlock, D., Kimura, K., Pimpl, W., Money, S.R., Jaffe, B.M., 1989. The role of
 271 serotonin in the canine secretory response to cholera toxin in vivo. *J. Pharmacol. Exp. Ther.*
 272 251, 71-76.

273 Mehlman, P.T., Higley, J.D., Fauscher, I., Lilly, A.A., Taub, D.M., Vickers, J., et al., 1994. Low
 274 CSF 5-HIAA concentrations and severe aggression and impaired impulse control in nonhuman
 275 primates. *Am. J. Psychiatry* 151, 1485-1491.

276 Meyers, K.M., Homsen, H., Seachord, C.L., 1982. Comparative study of platelet dense granule
 277 constituents. *Am. J. Physiol.* 243, R454-461.

278 Mikkelsen, J., Lund, J.D., 2000. Euthanasia of dogs due to behavioral problems: an
 279 epidemiological study of euthanasia of dogs in Denmark, with a special focus on problems of
 280 aggression. *Eur. J. Comp. Anim. Pract.* 10, 143-150.

281 Muck-Seler, D., Pivac, N., Sagud, M., Jakovljevic, M., Mihaljevic-Peles, A., 2002. The effects
 282 of paroxetine and tianeptine on peripheral biochemical markers in major depression. *Prog.*
 283 *Neuropsychopharmacol. Biol. Psychiatry* 26, 1235-1243.

284 Nelson, R.J., Chiavegatto, S., 2001. Molecular basis of aggression. *Trends Neurosci.* 24, 731-
 285 739.

286 Ortiz, J., Artigas, F., Gelpí, E., 1988. Serotonergic status in human blood. *Life Sci.* 43, 983-990.

287 Overall, K.L., Love, M., 2001. Dog bites to humans—demography, epidemiology, injury, and
 288 risk. *J. Am. Vet. Med. Assoc.* 218, 1923-1934.

289 Palacio, J., León, M., García-Belenguer, S., 2005. Aspectos epidemiológicos de las mordeduras
 290 caninas. *Gac. Sanit.* 19, 50-58.

291 Peremans, K., Audenaert, K., Coopman, F., Hoybergs, Y., Slegers, G., Van Bree, H., et al.,
 292 2002. Functional brain imaging of serotonin-2A receptors in impulsive dogs: a pilot study.
 293 *Vlaams Diergeneeskundig Tijdschrift.* 71, 340-347.

294 Peremans, K., Audenaert, K., Coopman, F., Blanckaert, P., Jacobs, F., Otte, A., et al., 2003.
 295 Estimates of regional cerebral blood flow and 5-HT_{2A} receptor density in impulsive, aggressive
 296 dogs with ^{99m}Tc-ECD and ¹²³I-5-I-R91150. *Eur. J. Nucl. Med. Mol. Imaging.* 30, 1538-1546.

297 Peremans, K., Audenaert, K., Hoybergs, Y., Otte, A., Goethals, I., Gielen, I., et al., 2005. The
 298 effect of citalopram hydrobromide on 5-HT_{2A} receptors in the impulsive-aggressive dog, as
 299 measured with ¹²³I-5-I-R91150 SPECT. *Eur. J. Nucl. Med. Mol. Imaging.* 32, 708-716.

300 Plein, H., Berk, M., 2001. The platelet as a peripheral marker in psychiatric illness. *Hum.*
 301 *Psychopharmacol.* 16, 229-236.

302 Popova, N.K., 2006. From genes to aggressive behavior: the role of serotonergic system.
 303 *Bioessays* 28, 495-503.

304 Reisner, I.R., Mann, J.J. Stanley, M., Huang, Y.Y., Houpt, K.A., 1996. Comparison of
 305 cerebrospinal fluid monoamine metabolite levels in dominant-aggressive and non-aggressive
 306 dogs. *Brain Res.* 714, 57-64.

307 Rosado, B., García-Belenguer, S., León, M., Chacón, G., Villegas, A., Palacio, J., 2010a. Blood
 308 concentrations of serotonin, cortisol and dehydroepiandrosterone in aggressive dogs. *Appl.*
 309 *Anim. Behav. Sci.* 123, 124-130.

310 Rosado, B., García-Belenguer, S., Palacio, J., Chacón, G., Villegas, A., Alcalde, A.I., 2010b.
 311 Serotonin transporter activity in platelets and canine aggression. *Vet. J.* 186, 104-105.

312 Sarrias, M.J., Cabré, P., Martínez, E., Artigas, F., 1990. Relationship between serotonergic
 313 measures in blood and cerebrospinal fluid simultaneously obtained in humans. *J. Neurochem.*
 314 54, 783-786.

315 Soderstrom, H., Blennow, K., Manhem, A., Forsman, A., 2001. CSF studies in violent
 316 offenders. I. 5-HIAA as a negative and HVA as a positive predictor of psychopathy. *J. Neural*
 317 *Transm.* 108, 869-878.

318 Soma, K.K., Scotti, M.A., Newman, A.E., Charlier, T.D., Demas, G.E., 2008. Novel
 319 mechanisms for neuroendocrine regulation of aggression. *Front. Neuroendocrinol.* 29, 476-489.

320 Stanley, B., Molcho, A., Stanley, M., Winchel, R., Gameroff, M.J., Parsons, B., Mann, J.J.,
 321 2000. Association of aggressive behavior with altered serotonergic function in patients who are
 322 not suicidal. *Am. J. Psychiatry* 157, 609-614.

323

324 **Table 1.** Mean (standard error) concentrations of serotonin in serum, plasma and platelets of
325 aggressive (n=28) and control non-aggressive dogs (n=10).