






ORIGINAL RESEARCH

Pediatric
OBESITY

WILEY

Population trajectories and age-dependent associations of obesity risk factors with body mass index from childhood to adolescence across European regions: A two-cohort study

Anton Schreuder¹  | Claudia Börnhorst² | Maïke Wolters²  |
 Toomas Veidebaum³ | Michael Tornaritis⁴ | Elida Sina² | Paola Russo⁵ |
 Luis A. Moreno^{6,7}  | Denes Molnar⁸ | Lauren Lissner⁹  |
 Stefaan De Henauw¹⁰ | Wolfgang Ahrens^{2,11} | Tanja Vrijkotte¹  | GrowH!
 Consortium

¹Department of Public Health, Academic Medical Center, University of Amsterdam, Amsterdam, The Netherlands

²Leibniz Institute for Prevention Research and Epidemiology – BIPS, Bremen, Germany

³National Institute for Health Development, Estonian Centre of Behavioral and Health Sciences, Tallinn, Estonia

⁴Research and Education Institute of Child Health, Strovolos, Cyprus

⁵Institute of Food Sciences, National Research Council, Avellino, Italy

⁶GENUD (Growth, Exercise, Nutrition and Development) Research Group, Faculty of Health Sciences, Universidad de Zaragoza, Instituto Agroalimentario de Aragón (IA2), Instituto de Investigación Sanitaria Aragón (IIS Aragón), Zaragoza, Spain

⁷Centro de Investigación Biomédica en Red de Fisiopatología de la Obesidad y Nutrición (CIBEROBN), Madrid, Spain

⁸Department of Pediatrics, Medical School, University of Pécs, Pécs, Hungary

⁹School of Public Health and Community Medicine, Institute of Medicine, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden

¹⁰Department of Public Health, Ghent University, Ghent, Belgium

¹¹Institute of Statistics, Faculty of Mathematics and Computer Science, University of Bremen, Bremen, Germany

Correspondence

Tanja Vrijkotte, Department of Public Health,
Academic Medical Center, University of
Amsterdam, P.O. Box 22660, 1100 DD
Amsterdam, The Netherlands.
Email: t.vrijkotte@amsterdamumc.nl

Funding information

Agencia Estatal de Investigación;
Bundesministerium für Bildung und Forschung,
Grant/Award Number: 01EA2102A; Fonds
Wetenschappelijk Onderzoek; FP7 Ideas:
European Research Council, Grant/Award
Number: 266044; Horizon 2020 Framework
Programme, Grant/Award Number: 727565;
Joint Programming Initiative A healthy diet for
a healthy life; Netherlands Cardiovascular
Research Initiative; Sixth Framework
Programme, Grant/Award Number: 016181;
ZonMw; Sarphati Amsterdam

Summary

Objective: To investigate population trajectories of behavioural risk factors of obesity from childhood to adolescence and their associations with body mass index (BMI) in children across European regions.

Methods: Data were harmonised between the European multi-centre IDEFICS/I. Family and the Amsterdam Born Children and their Development Cohort. Participants were aged 2.0–9.9 and 5.0–7.5 years at baseline, respectively, and were followed until age 18 years. Behavioural risk factors of interest included diet, physical activity, media use and sleep. Mixed effects models were used for statistical analyses to account for repeated measurements taken from the same child.

Results: The study included a total of 14 328 individuals: 4114, 4582, 3220 and 2412 participants from Northern, Southern, Eastern Europe and Amsterdam, respectively. Risk factor means and prevalences changed with age, but the trajectories were mostly similar across regions. Almost no associations between

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2023 The Authors. *Pediatric Obesity* published by John Wiley & Sons Ltd on behalf of World Obesity Federation.

behavioural factors and BMI were found at the age of 6 years. At 11 years, daily sugar-sweetened foods consumption, use of active transport, sports club membership and longer nocturnal sleep duration were negatively associated with BMI in most regions; positive associations were found with media use. Most associations at 11 years of age persisted to 15 years.

Conclusions: Whilst population trajectories of media use and nocturnal sleep duration are similar across European regions, those of other behavioural risk factors like active transport and daily vegetable consumption differ. Also, associations between behavioural risk factors and BMI become stronger with age and show similar patterns across regions.

KEYWORDS

adolescent, child, epidemiology, multicenter, prevalence, risk factors

1 | INTRODUCTION

Whilst some countries have reached a plateau, many others continue to show a trend of increasing average body mass index (BMI) amongst children up to the age of 18 years since the 1970s.^{1,2} Children with overweight or obesity have an increased risk of physical illnesses, mental disorders and early death.^{2–4} Children are especially vulnerable to an obesogenic environment due to their limited understanding of what healthy behaviours are, and their dependence on their parents and environment.^{2,5,6} Whereas weight gain is a result of energy intake greater than energy expenditure, the environment plays a complex role on physiological pathways that influence satiety and behaviours to attain this balance.⁷

Regional characteristics such as genetics, culture, climate and infrastructure can explain differences in weight status between populations.^{2,8} Previous works found region and ethnicity to be associated with variations in, for example, diet,^{9–11} physical activity,^{12,13} maternal BMI,^{14–18} parental educational level,^{14,19} feeding patterns,^{14,18} weight perception¹⁷ and height.²⁰ Additionally, the influence of obesity risk factors on children is complicated by changes that occur with age and development.² This means that children may be more vulnerable to certain risk factors during specific transition periods.^{21–27} Despite the extensive list of risk factors for obesity,^{2,27} less is known about the ages at which risk factors show their most detrimental effects. This information is crucial for guiding the implementation of effective strategies in sensitive time windows during the early life course and in different environments.

We performed an explorative study on various risk factors of obesity in a multi-centre population cohort from medium-sized cities across European regions and, for comparison, a cohort from a large city in the Netherlands. More specifically, we investigated population trajectories of behavioural risk factors from childhood to adolescence across European regions, and investigated whether these risk factors are associated with BMI in an age-dependent manner.

2 | METHODS

2.1 | Participant data

The present study is based on data from two cohorts: the IDEFICS (Identification and Prevention of Dietary- and Lifestyle-Induced Health Effects in Children and Infants)/I.Family (Investigating the determinants of food choice, lifestyle and health in European children, adolescents and their parents) and ABCD (Amsterdam Born Children and their Development) studies. The IDEFICS/I.Family study is a multi-centre population-based cohort of children from 8 European countries: Belgium, Cyprus, Estonia, Germany, Hungary, Italy, Sweden and Spain.^{28,29} The focus is on investigating the aetiology of diseases related to diet and behaviour in children and their families. The baseline examination wave (W0), conducted from September 2007 to June 2008, included 16 229 children aged 2.0–9.9 years. Two years later, the first follow-up wave (W1) took place, enrolling 2544 new participants. Two subsequent follow-up waves were conducted 6 (W2) and 14 years (W3) after baseline. Questionnaire assessments and physical examinations were performed using standardised procedures across all countries. At W3, BMI was self-reported because no physical examinations were performed. Ethical approval was obtained from the institutional review boards of all 8 study centres. Written informed consent for examinations and sample collection was provided by the parents and children aged 12 years and older; younger children provided oral consent.

The ABCD study is a cohort of Dutch children who were followed up since pregnancy.³⁰ The aim of ABCD is to investigate the effects of perinatal factors on health outcomes up to young adulthood. W0, W1 and W2 took place at around the ages of 6 years (range = 5.0–7.5 years), 12 years (10.6–13.7 years) and 16 years (15.1–17.2 years), respectively. At each wave, each parent filled in questionnaires about their own behaviours and the health and development of their child (fathers were only sent questionnaires at W1 and W2). The children were asked to fill in questionnaires about their own behaviours from W1 onwards. Growth data was measured at W0 and W1 when children underwent physical examinations by the ABCD health check. This was

complemented by youth healthcare registration measurements until the age of approximately 13 years.³¹ Growth data from W2 was self-reported. A total of 8266 pregnant women residing in Amsterdam filled out the pregnancy questionnaire between January 2003 and March 2004, of whom 7050 (85%) gave permission for follow-up.

2.2 | Variables and harmonisation

All variables shared between the IDEFICS/I.Family and ABCD were considered for harmonisation, excluding questions from the Strength and Difficulties Questionnaire because some questions were not assessed in IDEFICS/I.Family. Based on the expert opinion of three researchers (A.S., C.B. and T.V.), variables with potential for harmonisation were selected and converted into common formats. In most cases, this required a re-categorisation combining certain categories or assignment of continuous values to certain categories. Variables that were considered comparable between studies (with or without transformations in one or both cohorts) and which were available at three time points in one study and no fewer than two time points in the other were included in the present study. A description of all variables is given in the supplement.

2.2.1 | Outcome

The primary outcome of this study is BMI, a common measure for weight status in epidemiological studies. This was measured in W0–W2 in IDEFICS/I.Family, and self-reported in W3. BMI was also converted into age- and sex-specific z-scores and was categorised into children with underweight, normal weight, overweight and obesity according to the extended International Obesity Task Force criteria.³²

2.2.2 | Risk factors

The exposures of interest were known and suspected behavioural risk factors of obesity-related to (i) diet, namely water consumption (times/day), sweetened drinks consumption (times/day, including artificially sweetened drinks and sweetened milk), daily vegetable consumption (yes vs. no), daily fruit consumption (yes vs. no), daily sugar-sweetened foods consumption (yes vs. no) and vegetarian diet (yes vs. no); (ii) physical activity, namely active transport to school (yes vs. no; walking or cycling at least 4 days per week) and sports club membership (yes vs. no); (iii) media use, namely television (TV) and personal computer (PC) media use (h/day) and (iv) nocturnal sleep duration (h/night). Data on sugar-sweetened food consumption and sports club membership were not available for W1 of the ABCD cohort.

2.2.3 | Covariates

The following variables were considered as potential confounders: (i) parental characteristics, namely mother's BMI (kg/m^2), father's BMI

(kg/m^2), highest parental ISCED (International Standard Classification of Education, high vs. low/medium)³³ and migrant background (both parents vs. one parent vs. neither parent); (ii) family structure/vulnerabilities, namely number of children in household, only child (yes vs. no), single parent (yes vs. no) and parental divorce (yes vs. no) and (iii) peri-/post-natal factors, namely smoking during pregnancy (yes vs. no), alcohol consumption during pregnancy (yes vs. no), maternal age at birth (years), child's sex (male vs. female), birthweight (g), preterm birth (yes vs. no), C-section delivery (yes vs. no), exclusive breastfeeding duration (no exclusive breastfeeding vs. <3 months vs. 3–<6 months vs. ≥ 6 months) and age at introduction to new foods (<4 months vs. 4 to <6 months vs. ≥ 6 months before consuming foods other than breast milk and formula milk). A more detailed description of the variables used is provided in the supplement.

IDEFICS was designed as a controlled intervention trial from W0 to W1. To account for possible effects the intervention may have had in the long run, all models are based on the IDEFICS/I.Family data were adjusted for children living in the control or intervention region (binary variable). However, as the IDEFICS intervention did not show any effects on body fatness or behavioural factors in the total study group, we consider it unlikely that the IDEFICS intervention will have affected our results.^{34–36}

2.3 | Data set preparation

Examination waves were excluded from our analysis if one of the following criteria applied: age less than 2 years, age greater than 18 years, missing participant's BMI, missing all dietary variables (i.e., vegetables, fruits, added sugar foods, water and sweetened drinks consumption frequencies), and fewer than two observations per participant (after applying the other criteria). The final study samples are subsequently referred to as 'ABCD cohort' and 'IDEFICS/I.Family cohort'. Figure 1 depicts the selection process leading to the final study samples.

Multiple imputations with 10 iterations per imputation were used to handle missing data using the 'mice' package's default method for each data type.³⁷ This was performed independently for the IDEFICS/I.Family and ABCD cohorts. A wide data set format (based on each wave) of the data set was used for this to improve estimates, including all variables in the algorithm.³⁸ This procedure was repeated 100 times to create 100 complete data sets.

2.4 | Statistical analysis

All analyses were performed using statistical program R version 4.1.2.³⁹ For the IDEFICS/I.Family cohort, the subsequent analyses were stratified by three regions: Northern (Belgium, Sweden, Germany), Southern (Cyprus, Italy, Spain) and Eastern Europe (Estonia, Hungary). Though located in Northern Europe, the ABCD cohort was treated as a separate fourth region—Amsterdam—to enable comparisons between cohorts as well as between medium-sized cities (approximately 18 000–400 000 inhabitants) and a large city




FIGURE 1 Participant selection flowchart. BMI, body mass index.

(>700 000 inhabitants). Continuous variables were reported as means and standard deviations, and categorical variables were described using frequencies and percentages. Histograms of continuous variables were visually inspected and deemed to be normally distributed.

Mixed effects models were used to account for the repeated measurements taken within the same child. In the main analysis, the mean (continuous variables) or prevalence (binary variables) of each risk factor across age were estimated, including subject-specific random intercepts. The dependent variable for each model was a risk factor, and the independent variables were three polynomials of age (i.e., age, age² and age³), sex and the three interaction terms between each pair (i.e., age-sex, age²-sex and age³-sex). Linear mixed models were used when the risk factor was continuous, and binomial generalised linear mixed models were used for binary risk factors. A BOBYQA (Bound Optimisation BY Quadratic Approximation) optimiser was used for the latter to handle convergence issues.⁴⁰ Results are displayed in plots stratified by sex with 99% confidence intervals.

In a second analysis, we estimated the age-specific associations between each behavioural risk factor and BMI, as well as associations between the well-established risk factors parental

ISCED and parental BMI and children's BMI. Linear mixed models were used with BMI as the dependent variable. The independent variables included the three polynomials of age, sex, their three interaction terms, a risk factor and three interaction terms between each polynomial of age and risk factor (i.e., age-risk factor, age²-risk factor and age³-risk factor) as a minimal adjustment set. All additional confounders derived from a directed acyclic graph (see Figure S1) were added as independent variables for full model adjustment.⁴¹ Table S1 lists all corresponding adjustment sets. The coefficients with 99% confidence intervals for the association of each risk factor with BMI at the ages of 6, 11 and 15 years were reported; this was done based on reparametrised analyses with age centred at 6, 11 and 15 years, respectively.

All analyses were performed on all 100 imputed data sets, the results of which were pooled. Sensitivity analyses were also performed on a cohort where, in addition to the other exclusion criteria, observations with underweight status were excluded. The purpose was to assess whether associations between risk factors and BMI differ depending on whether most changes in BMI occur in children with underweight compared to children without.

3 | RESULTS

The descriptive statistics at baseline before and after imputations are summarised in Table S3 and Table 1, respectively. The IDEFICS/I.Family cohort consisted of 11 916 participants with 27 831 observations over

all examination waves, and the ABCD cohort consisted of 2412 participants with 5884 observations (Figure 1). The frequency of missing data in the final IDEFICS/I.Family (max. 34.2%) and ABCD cohorts (max. 39.7%) are reported in Table S2. Several differences between cohorts were noted (Table 1). ABCD had a greater proportion of children whose

TABLE 1 Characteristics of the study population at baseline (one imputed data set).

Variable	Amsterdam, n = 2412	Northern Europe, n = 4114	Southern Europe, n = 4582	Eastern Europe, n = 3220
Continuous variables, mean (SD)				
Age (years)	6.1 (2.3)	5.9 (1.9)	6.2 (1.7)	6.2 (2.1)
Birthweight (g)	3504 (536)	3397 (596)	3183 (552)	3451 (579)
Mother's BMI (kg/m ²)	23.5 (3.9)	24.2 (4.3)	23.9 (4.3)	23.6 (4.4)
Father's BMI (kg/m ²)	24.8 (3.0)	26.0 (3.7)	27.2 (4.1)	26.9 (4.5)
Number of children in household	2.2 (0.9)	2.2 (0.9)	2.2 (0.9)	2.0 (0.9)
Water consumption (times/day)	1.5 (1.1)	2.4 (1.4)	3.6 (0.9)	2.7 (1.4)
Sweetened drinks consumption (times/day)	2.1 (1.2)	1.8 (1.8)	1.8 (1.5)	2.0 (1.7)
Media use (h/day)	1.4 (0.9)	1.5 (0.9)	1.7 (1.1)	1.8 (1.1)
Nocturnal sleep duration (h/night)	10.5 (0.9)	10.9 (0.9)	10.0 (1.0)	9.6 (1.0)
BMI z-score	−0.2 (1.1)	0.1 (1.0)	0.7 (1.2)	0.1 (1.1)
Categorical variables, frequency (%)				
Female sex (yes)	1175 (48.7)	2060 (50.1)	2325 (50.7)	1570 (48.8)
High parental ISCED (yes)	1836 (76.1)	2429 (59.0)	1867 (40.7)	1703 (52.9)
Migrant background	-	-	-	-
Neither parent	1724 (71.5)	3380 (82.2)	3706 (80.9)	3091 (96.0)
One parent	462 (19.2)	321 (7.8)	637 (13.9)	105 (3.3)
Both parents	226 (9.4)	413 (10.0)	239 (5.2)	24 (0.7)
Smoking during pregnancy	-	-	-	-
No smoking	2240 (92.9)	3518 (85.5)	3898 (85.1)	2909 (90.3)
<1 cigarette per day	59 (2.4)	299 (7.3)	381 (8.3)	226 (7.0)
≥1 cigarette per day	113 (4.7)	297 (7.2)	303 (6.6)	85 (2.6)
Alcohol during pregnancy (yes)	729 (30.2)	1073 (26.1)	1167 (25.5)	944 (29.3)
Preterm birth (yes)	212 (8.8)	464 (11.3)	539 (11.8)	335 (10.4)
C-section delivery (yes)	316 (13.1)	816 (19.8)	1683 (36.7)	600 (18.6)
Exclusive breastfeeding duration	-	-	-	-
No exclusive breastfeeding	379 (15.7)	1364 (33.2)	1621 (35.4)	551 (17.1)
<3 months	773 (32.0)	893 (21.7)	978 (21.3)	831 (25.8)
3 to <6 months	899 (37.3)	1213 (29.5)	1228 (26.8)	827 (25.7)
≥6 months	361 (15.0)	644 (15.7)	755 (16.5)	1011 (31.4)
Age at introduction to new foods ^a	-	-	-	-
<4 months	123 (5.1)	639 (15.5)	226 (4.9)	494 (15.3)
4 to <6 months	892 (37.0)	2093 (50.9)	1855 (40.5)	1275 (39.6)
≥6 months	1397 (57.9)	1382 (33.6)	2501 (54.6)	1451 (45.1)
Being an only child (yes)	383 (15.9)	646 (15.7)	809 (17.7)	884 (27.5)
Single parent (yes)	175 (7.3)	324 (7.9)	948 (20.7)	399 (12.4)
Parents divorced (yes)	176 (7.3)	498 (12.1)	184 (4.0)	529 (16.4)
Daily vegetable consumption (yes)	1619 (67.1)	3261 (79.3)	2090 (45.6)	2050 (63.7)
Daily fruit consumption (yes)	1571 (65.1)	2816 (68.4)	3113 (67.9)	2027 (63.0)

(Continues)

TABLE 1 (Continued)

Variable	Amsterdam, n = 2412	Northern Europe, n = 4114	Southern Europe, n = 4582	Eastern Europe, n = 3220
Daily sugar-sweetened foods consumption (yes)	1764 (85.8)	2148 (52.2)	2146 (46.8)	1611 (50.0)
Vegetarian diet (yes)	80 (3.3)	34 (0.8)	20 (0.4)	13 (0.4)
Active transport to school (yes) ^b	1102 (45.7)	1673 (40.7)	1204 (26.3)	1240 (38.5)
Sports club member (yes)	1147 (55.8)	2207 (53.6)	1967 (42.9)	1301 (40.4)
With overweight or obesity (yes)	186 (7.7)	452 (11.0)	1351 (29.5)	452 (14.0)

Note: Continuous variables are reported as means with SDs in brackets. Categorical variables are reported as frequencies with percentages in brackets.

Abbreviations: BMI, body mass index; SD, standard deviation.

^aFoods other than breast milk and formula milk.

^bWalking or cycling at least 4 days per week.




FIGURE 2 Mean television (TV) and personal computer (PC) media use across age and European regions, stratified by sex. Shaded areas represent 99% confidence intervals, where darker shades indicate an overlap of confidence intervals from the two lines.

parents had high ISCED (76.1% in ABCD vs. 50.3% in IDEFICS/I. Family) and migrant backgrounds (28.6% vs. 14.6%), and a lower mean paternal BMI (24.8 vs. 26.7 kg/m²), fewer C-section deliveries (13.1% vs. 26.0%) and fewer single-parent households (7.6% vs. 14.0%).

3.1 | Population trajectories

The age-dependent means and prevalences of behavioural risk factors are given in Figures 2 and 3, and S2–S9. Compared to the IDEFICS/I. Family cohort, the ABCD participants reported less water consumption (mean = 2.9 vs. 1.5 times/day, respectively) and more often sugar-sweetened foods on a daily basis at baseline (49.2% vs. 85.8%) (Figures S2 and S6). Children in Amsterdam also had a higher proportion of vegetarians across all ages (0.5% vs. 3.3% at baseline), increasing beyond 10% amongst female teenagers (Figure S7). Except for sports club membership, males more often engaged in unhealthier behaviours than females (i.e., more frequent intake of sweetened drinks, less frequent intake of fruits and vegetables, less likely to be vegetarian and more media use).

Most behaviours showed similar trends across the four regions. For example, media usage, sweetened drinks consumption frequency and the proportion of sports club members increased with age (Figures 2,

S3 and S8). In contrast, the proportion of daily fruit consumers, the proportion of daily sugar-sweetened foods consumers, and nocturnal sleep duration decreased with age (Figures S5, S6 and S9).

Other risk factors displayed more inter-regional variation, the most notable being the proportion of active transport users (Figure 3). Active transport usage was highest in Amsterdam—approaching 100% amongst teenagers—and lowest in Southern Europe—remaining stable at 10%. Whilst this was similar between the sexes, an exception was that active transport was used more amongst male teenagers in Eastern Europe.

The prevalence of daily vegetable consumers also varied across regions (Figure S4), with the smallest fluctuation with age in Northern Europe (between 81% and 86% across all ages), the steepest increase with age in Southern Europe (mean at age 5 years = 40% vs. mean at age 15 years = 78%), a more gradual increase with age in Eastern Europe (mean at age 5 years = 64% vs. mean at age 15 years = 74%) and a decrease with age in Amsterdam (mean at age 5 years = 70% vs. mean at age 15 years = 59%).

3.2 | Associations with BMI

The average BMI, average BMI z-score and the proportion of children with overweight or obesity with age in each region are shown in




FIGURE 3 Proportion of active transport users across age and European regions, stratified by sex. Shaded areas represent 99% confidence intervals, where darker shades indicate an overlap of confidence intervals from the two lines.

TABLE 2 Fully adjusted associations between risk factors and body mass index (BMI) at the age of 6 years.

Risk factor	Amsterdam Coefficient (LCI–UCI)	Northern Europe Coefficient (LCI–UCI)	Southern Europe Coefficient (LCI–UCI)	Eastern Europe Coefficient (LCI–UCI)
Highest parental ISCED ^a (high)	−0.41 (−1.00 to 0.18)	−0.07 (−0.22 to 0.08)	−0.45 (−0.66 to −0.23)	−0.10 (−0.33 to 0.13)
Mother's BMI ^b (kg/m ²)	0.04 (−0.02 to 0.10)	0.05 (0.03–0.07)	0.07 (0.05–0.09)	0.07 (0.05–0.10)
Father's BMI ^b (kg/m ²)	0.03 (−0.06 to 0.11)	0.00 (−0.02 to 0.01)	−0.01 (−0.03 to 0.01)	−0.01 (−0.03 to 0.02)
Water consumption ^c (times/day)	−0.10 (−0.30 to 0.10)	−0.01 (−0.05 to 0.03)	0.06 (−0.02 to 0.14)	0.01 (−0.06 to 0.07)
Sweetened drinks consumption ^c (times/day)	−0.05 (−0.22 to 0.11)	−0.04 (−0.08 to 0.00)	−0.05 (−0.10 to 0.00)	0.02 (−0.03 to 0.07)
Daily vegetable consumption ^c (yes)	−0.09 (−0.62 to 0.43)	0.13 (−0.01 to 0.28)	0.04 (−0.11 to 0.19)	−0.04 (−0.22 to 0.14)
Daily fruit consumption ^c (yes)	0.34 (−0.17 to 0.86)	0.02 (−0.11 to 0.15)	0.04 (−0.12 to 0.20)	−0.01 (−0.19 to 0.17)
Daily sugar-sweetened foods consumption ^c (yes)	−0.97 (−2.36 to 0.41)	−0.02 (−0.14 to 0.10)	−0.17 (−0.32 to −0.02)	0.07 (−0.10 to 0.24)
Vegetarian diet ^c (yes)	0.07 (−0.95 to 1.09)	−0.02 (−0.80 to 0.76)	0.43 (−0.63 to 1.49)	−0.33 (−1.69 to 1.02)
Active transport to school ^d (yes)	−0.71 (−1.29 to −0.12)	−0.03 (−0.16 to 0.09)	−0.11 (−0.29 to 0.08)	−0.06 (−0.26 to 0.13)
Sports club member ^d (yes)	0.25 (−0.65 to 1.14)	0.11 (−0.01 to 0.23)	0.11 (−0.04 to 0.27)	0.01 (−0.17 to 0.19)
Media use ^e (h/day)	−0.17 (−0.37 to 0.03)	−0.04 (−0.11 to 0.03)	−0.03 (−0.10 to 0.04)	0.02 (−0.07 to 0.10)
Nocturnal sleep duration ^f (h/night)	0.24 (0.03–0.46)	−0.04 (−0.11 to 0.03)	−0.07 (−0.14 to 0.01)	−0.03 (−0.13 to 0.07)

Note: Results from linear mixed models. The IDEFICS/I.Family cohort was divided into Northern (Belgium, Sweden and Germany), Southern (Cyprus, Italy and Spain) and Eastern Europe (Estonia and Hungary). Coefficients whose 99% confidence intervals do not include zero are indicated in bold.

Abbreviations: BMI, body mass index; LCI, lower 99% confidence interval; UCI, upper 99% confidence interval.

^aAdjusted for: age, sex and migrant background.

^bAdjusted for: age, sex, migrant background, highest parental ISCED, number of children in household, only child, single parent and parental divorce.

^cAdjusted for: age, sex, migrant background, highest parental ISCED, mother's BMI, father's BMI, smoking during pregnancy, alcohol during pregnancy, birthweight, maternal age at birth, preterm birth, C-section delivery, exclusive breastfeeding duration and age at introduction to new food items, number of children in household, only child, single parent, parental divorce, nocturnal sleep duration, media use and active transport to school.

^dAdjusted for: age, sex, migrant background, highest parental ISCED, mother's BMI, father's BMI, number of children in household, only child, single parent, parental divorce and media use.

^eAdjusted for: age, sex, migrant background, highest parental ISCED, mother's BMI, father's BMI, number of children in household, only child, single parent and parental divorce.

^fAdjusted for: age, sex, migrant background, highest parental ISCED, number of children in household, only child, single parent, parental divorce, media use and active transport to school.

Figures S10–S12. The fully adjusted effect estimates obtained for the associations between the behavioural risk factors and BMI at the ages of 6, 11 and 15 years are reported in Tables 2–4, respectively. Across all regions and ages, each increase of the mother's

BMI by 1 kg/m² increased the child's BMI by 0.04–0.21 kg/m² on average. At the ages of 11 and 15 years, the child's BMI was also positively associated with father's BMI and negatively associated with high parental ISCED.

TABLE 3 Fully adjusted associations between risk factors and body mass index (BMI) at the age of 11 years.

Risk factor	Amsterdam Coefficient (LCI-UCI)	Northern Europe Coefficient (LCI-UCI)	Southern Europe Coefficient (LCI-UCI)	Eastern Europe Coefficient (LCI-UCI)
Highest parental ISCED ^a (high)	-1.13 (-1.49 to -0.76)	-0.80 (-0.98 to -0.63)	-1.23 (-1.46 to -0.99)	-0.51 (-0.76 to -0.27)
Mother's BMI ^b (kg/m ²)	0.17 (0.14-0.21)	0.18 (0.16-0.20)	0.20 (0.17-0.22)	0.18 (0.15-0.20)
Father's BMI ^b (kg/m ²)	0.14 (0.10-0.19)	0.11 (0.10-0.13)	0.11 (0.09-0.13)	0.09 (0.07-0.12)
Water consumption ^c (times/day)	0.01 (-0.09 to 0.12)	0.05 (0.00-0.11)	0.00 (-0.08 to 0.08)	0.08 (0.01-0.16)
Sweetened drinks consumption ^c (times/day)	0.01 (-0.08 to 0.10)	-0.01 (-0.05 to 0.03)	0.01 (-0.04 to 0.05)	-0.04 (-0.08 to 0.01)
Daily vegetable consumption ^c (yes)	-0.28 (-0.55 to 0.00)	-0.12 (-0.30 to 0.06)	-0.30 (-0.48 to -0.12)	-0.05 (-0.25 to 0.15)
Daily fruit consumption ^c (yes)	0.01 (-0.25 to 0.28)	-0.07 (-0.23 to 0.08)	-0.21 (-0.39 to -0.03)	0.02 (-0.17 to 0.22)
Daily sugar-sweetened foods consumption ^c (yes)	-1.06 (-3.45 to 1.33)	-0.22 (-0.37 to -0.07)	-0.22 (-0.40 to -0.04)	-0.36 (-0.55 to -0.17)
Vegetarian diet ^c (yes)	0.05 (-0.49 to 0.59)	0.41 (-0.23 to 1.06)	0.42 (-0.76 to 1.60)	-1.12 (-2.45 to 0.21)
Active transport to school ^d (yes)	-0.56 (-0.99 to -0.12)	-0.05 (-0.20 to 0.10)	-0.38 (-0.59 to -0.17)	0.29 (0.08-0.50)
Sports club member ^d (yes)	0.43 (-1.31 to 2.17)	-0.30 (-0.47 to -0.12)	-0.32 (-0.50 to -0.14)	-0.26 (-0.47 to -0.05)
Media use ^e (h/day)	0.15 (0.04-0.27)	0.12 (0.06-0.18)	0.18 (0.11-0.25)	0.13 (0.06-0.21)
Nocturnal sleep duration ^f (h/night)	0.08 (-0.09 to 0.24)	-0.10 (-0.18 to -0.02)	-0.15 (-0.25 to -0.05)	-0.16 (-0.29 to -0.03)

Note: Results from linear mixed models. The IDEFICS/I.Family cohort was divided into Northern (Belgium, Sweden and Germany), Southern (Cyprus, Italy and Spain) and Eastern Europe (Estonia and Hungary). Coefficients whose 99% confidence intervals do not include zero are indicated in bold.

Abbreviations: BMI, body mass index; LCI, lower 99% confidence interval; UCI, upper 99% confidence interval.

^aAdjusted for: age, sex and migrant background.

^bAdjusted for: age, sex, migrant background, highest parental ISCED, number of children in household, only child, single parent and parental divorce.

^cAdjusted for: age, sex, migrant background, highest parental ISCED, mother's BMI, father's BMI, smoking during pregnancy, alcohol during pregnancy, birthweight, maternal age at birth, preterm birth, C-section delivery, exclusive breastfeeding duration and age at introduction to new food items, number of children in household, only child, single parent, parental divorce, nocturnal sleep duration, media use and active transport to school.

^dAdjusted for: age, sex, migrant background, highest parental ISCED, mother's BMI, father's BMI, number of children in household, only child, single parent, parental divorce and media use.

^eAdjusted for: age, sex, migrant background, highest parental ISCED, mother's BMI, father's BMI, number of children in household, only child, single parent and parental divorce.

^fAdjusted for: age, sex, migrant background, highest parental ISCED, number of children in household, only child, single parent, parental divorce, media use and active transport to school.

Regarding behavioural risk factors at the age of 6 years, active transport in Amsterdam and daily sugar-sweetened food consumption in Southern Europe were found to be negatively associated with BMI, and nocturnal sleep duration was positively associated with BMI in Amsterdam (Table 2). At 11 years, positive associations were found for water consumption and media use whereas negative associations were found for daily vegetable consumption, daily sugar-sweetened foods consumption, sports club membership and nocturnal sleep duration (Table 3). For active transport, negative associations were found in Amsterdam and Southern Europe and a positive association was found in Eastern Europe. Only a few associations persisted until the age of 15 years, namely water consumption, daily sugar-sweetened foods consumption, active transport, sports club membership, media use and nocturnal sleep duration (Table 4).

Fully adjusted associations did not differ markedly from the minimally adjusted associations (Tables S4-S6). Associations found from the sensitivity analyses—excluding 1655/14 328 (11.6%) children with underweight at W0—were markedly similar to their respective results from the main analyses (Figure S13, Tables S7-S14).

4 | DISCUSSION

4.1 | Population trajectories

We performed an explorative analysis on 10 known and suspected risk factors of obesity in children from four European regions within two cohorts. Whilst most regions showed similar trends in means and prevalences with age, considerable regional differences were found for the proportion of active transport users and daily vegetable consumers. Media use, nocturnal sleep duration and the proportion of sports club members varied the least across regions. Except for being more likely to be sports club members, males more often engaged in unhealthy behaviours than females.⁴²⁻⁴⁷ In line with the literature, males consumed more sweetened drinks,⁴⁶ had more screen time,^{42,44,45} were less likely to eat vegetables and fruits every day^{44,45} and were less often vegetarian.⁴⁰⁻⁴⁵

Some risk factors from Amsterdam stand out as compared to the IDEFICS/I.Family cohort regions. The high proportion of ABCD participants using an active form of transport can be attributable to the Netherlands having the world's best cycling infrastructure.⁴⁸

TABLE 4 Fully adjusted associations between risk factors and body mass index (BMI) at the age of 15 years.

Risk factor	Amsterdam Coefficient (LCI-UCI)	Northern Europe Coefficient (LCI-UCI)	Southern Europe Coefficient (LCI-UCI)	Eastern Europe Coefficient (LCI-UCI)
Highest parental ISCED ^a (high)	-0.69 (-1.16 to -0.22)	-1.67 (-2.14 to -1.20)	-0.91 (-1.53 to -0.29)	-0.68 (-1.19 to -0.16)
Mother's BMI ^b (kg/m ²)	0.21 (0.17-0.26)	0.21 (0.16-0.25)	0.15 (0.09-0.21)	0.20 (0.14-0.25)
Father's BMI ^b (kg/m ²)	0.14 (0.07-0.20)	0.14 (0.10-0.19)	0.09 (0.02-0.15)	0.13 (0.08-0.19)
Water consumption ^c (times/day)	0.19 (0.04-0.34)	-0.06 (-0.22 to 0.10)	0.07 (-0.21 to 0.36)	0.06 (-0.11 to 0.23)
Sweetened drinks consumption ^c (times/day)	0.03 (-0.09 to 0.15)	0.09 (0.00-0.19)	-0.06 (-0.20 to 0.07)	0.14 (0.05-0.23)
Daily vegetable consumption ^c (yes)	-0.01 (-0.40 to 0.38)	-0.37 (-0.88 to 0.15)	-0.09 (-0.76 to 0.58)	0.40 (-0.16 to 0.95)
Daily fruit consumption ^c (yes)	0.01 (-0.38 to 0.39)	-0.47 (-0.95 to 0.01)	-0.39 (-1.03 to 0.26)	0.24 (-0.28 to 0.77)
Daily sugar-sweetened foods consumption ^c (yes)	-0.10 (-0.67 to 0.48)	0.15 (-0.35 to 0.65)	-0.99 (-1.64 to -0.35)	-0.23 (-0.74 to 0.29)
Vegetarian diet ^c (yes)	-0.11 (-0.78 to 0.56)	0.21 (-1.15 to 1.56)	3.03 (-2.11 to 8.17)	-0.58 (-3.33 to 2.16)
Active transport to school ^d (yes)	-0.11 (-0.70 to 0.48)	-0.60 (-1.09 to -0.10)	-0.89 (-1.60 to -0.18)	0.10 (-0.43 to 0.62)
Sports club member ^d (yes)	0.41 (-0.24 to 1.07)	-0.83 (-1.33 to -0.33)	-0.88 (-1.52 to -0.24)	-0.37 (-0.90 to 0.16)
Media use ^e (h/day)	0.16 (0.03-0.28)	0.29 (0.15-0.43)	0.03 (-0.19 to 0.25)	0.17 (0.03-0.31)
Nocturnal sleep duration ^f (h/night)	-0.15 (-0.30 to 0.01)	-0.08 (-0.31 to 0.15)	0.09 (-0.22 to 0.40)	-0.36 (-0.65 to -0.08)

Note: Results from linear mixed models. The IDEFICS/I.Family cohort was divided into Northern (Belgium, Sweden and Germany), Southern (Cyprus, Italy and Spain) and Eastern Europe (Estonia and Hungary). Coefficients whose 99% confidence intervals do not include zero are indicated in bold.

Abbreviations: BMI, body mass index; LCI, lower 99% confidence interval; UCI, upper 99% confidence interval.

^aAdjusted for: age, sex and migrant background.

^bAdjusted for: age, sex, migrant background, highest parental ISCED, number of children in household, only child, single parent and parental divorce.

^cAdjusted for: age, sex, migrant background, highest parental ISCED, mother's BMI, father's BMI, smoking during pregnancy, alcohol during pregnancy, birthweight, maternal age at birth, preterm birth, C-section delivery, exclusive breastfeeding duration and age at introduction to new food items, number of children in household, only child, single parent, parental divorce, nocturnal sleep duration, media use and active transport to school.

^dAdjusted for: age, sex, migrant background, highest parental ISCED, mother's BMI, father's BMI, number of children in household, only child, single parent, parental divorce and media use.

^eAdjusted for: age, sex, migrant background, highest parental ISCED, mother's BMI, father's BMI, number of children in household, only child, single parent and parental divorce.

^fAdjusted for: age, sex, migrant background, highest parental ISCED, number of children in household, only child, single parent, parental divorce, media use and active transport to school.

Regarding fluids consumption, sweetened drinks consumption frequency was similar across regions but not water consumption frequency. Whereas drinking less water is likely to be compensated by drinking more sweetened beverages,^{49,50} this pattern was not clear in our findings. The higher water consumption in Southern Europe may be attributable to the warmer climate and the observed larger body sizes. Less water consumption before the age of 11 in Amsterdam is likely an underestimation compared to the other European regions. This is because a different questionnaire was used for W0 as opposed to later waves: water consumption was measured as times per week instead of glasses per day, respectively.

The population trajectories of most behaviours shifted in the transition from childhood to adolescence. A possible causal explanation is that children obtain more social contacts around that age, thus their behaviours become more likely to be shaped by their peers than by their parents or siblings.^{51,52} However, differences may also be partly related to children starting to fill in the questionnaires by themselves at approximately 12 years of age. Despite both self-reports and proxy reports being prone to social desirability bias, a discrepancy usually remains between the two.⁵³⁻⁵⁵

4.2 | Associations with BMI

Consistent with previous literature,^{2,5-7,22,31,56} parental ISCED and parental BMI were strongly associated with child's BMI regardless of the adjustment set. The associations of maternal BMI with the child's BMI were stronger than for the paternal BMI across the reported ages (i.e., 6, 11 and 15 years). This may be because most children spend more time with their mothers and imitate behaviours from them,⁵ and because women are more often responsible for meal preparation.^{57,58} Moreover, physiological adaptations might already occur in the intra-uterine period (foetal programming) as a consequence of the adverse metabolic or inflammatory environment.⁵⁹ Regardless, members within the same household tend to have similar weight status given shared genes and behaviours.

Amongst behavioural risk factors, almost no associations were found with BMI at the age of 6 years. At 11 and 15 years of age, negative associations were found between BMI and sweetened drinks consumption frequency, daily sugar-sweetened foods consumption, active transport, sports club membership, less media use and longer nocturnal sleep duration in most regions. We note that nocturnal sleep duration changed from being positively associated with BMI

(in Amsterdam) at the age of 6 years to becoming negatively associated at 11 and 15 years of age. This suggests that too much sleep as well as too little sleep is associated with BMI,^{25,60} though we did not consider other possible factors such as sleep quality, time of going to bed and time of getting up.

The classification of sugary foods and drinks as risk factors for obesity is well established,⁶¹ so the negative associations found require an explanation. One possibility is that sugar-sweetened foods are difficult to classify and may be unreliably reported. Moreover, foods considered unhealthy may be particularly prone to misreporting due to socially desirable answering behaviour. Additionally, total caloric intake was unavailable. Another possible explanation is that children who are considered to be overweight by their parents or themselves may have more (self-)imposed dietary restrictions than others.⁶²

A notable finding was that active transport was positively associated with BMI in Eastern Europe at age 11. A possible explanation is that, in Eastern Europe, active transport is a proxy for lower income and the inability to afford other modes of transport.⁴⁸ The observation that active transport becomes less popular amongst females beyond puberty whilst remaining viable for males may be related to the perception of the built environment.⁶³ For example, well-lit streets at night are associated with active transport amongst females.

The sensitivity analyses produced results equivalent to those from the main analyses. This suggests that the associations between behavioural risk factors and BMI do not differ when stratified into BMI categories. However, it is possible that the exclusion of observations with underweight status was insufficient to noticeably affect the results.

Overall, the effect sizes of the associations found between each behavioural risk factor and BMI can be interpreted as small.⁶⁴ Therefore, an implication of this exploratory study is that behavioural risk factors should not be considered as independent from other factors: targeting individual risk factors is expected to offer limited benefits. The causal pathway from upstream factors to behaviour and from behaviour to BMI is complex and unique to each individual, also dependent on genetic and environmental factors which were not available for this study.^{2,5-8} Consideration of the systemic conditions that shape health-related behaviours—socioeconomic status in particular—would likely improve the effectiveness of preventing children from gaining too much weight.⁵⁶

4.3 | Strengths and limitations

Strengths of this study are its large sample size, the inclusion of many risk factors, and the use of data from various European countries and from two independent studies. Whilst Amsterdam would typically be grouped with Northern Europe, the proportions of migrants and highly educated inhabitants, for example, are greater in Amsterdam; these differences may be associated with city size. The finding that most population risk factor trajectories were similar indicates that harmonisation was performed well and that many behaviours may be universally promising targets for interventions.

Some limitations should be taken into consideration. In this study, we investigated the age-specific associations between behavioural risk factors and BMI. Despite following a longitudinal modelling approach based on mixed effects models (accounting for correlations amongst repeated measurements taken from the same child), the associations between the behavioural risk factors and BMI at the different ages should be interpreted cross-sectionally because the exposures and outcomes were assessed at the same ages.

Data collection was not performed in a consistent manner throughout the ABCD examination waves. Unlike the IDEFICS/I.Family data, the ABCD data were not interspersed across all ages. In combination with the smaller sample size, the gaps between ages 7–10 and 12–14 years resulted in less smooth population trajectories as compared to the trajectories estimated based on the IDEFICS/I.Family data. However, to obtain stable estimates for the associations between behavioural risk factors and BMI, associations were only estimated at ages for which there was sufficient data amongst both cohorts.

Our measure of media use was only based on TV and PC screen time. Currently, the estimated percentages of 9- to 16-year-olds with internet access who used smartphones daily in the studied countries ranged from 76% in Spain to 87% in Estonia.⁶⁵ With the increase of mobile phones and other digital media being used amongst children, it may be considered a limitation that smartphone exposure was not available for this study. Whilst this may be negligible for the first and second assessment waves, our reported media use may be an underestimation of total media usage in the later examination waves.⁶⁶

5 | CONCLUSIONS

The population means and prevalences of most behavioural risk factors of obesity generally change with age. Some risk factors trajectories are consistent across European regions, such as media use and nocturnal sleep duration, whilst other trajectories differ considerably between regions, such as active transport and daily vegetable consumption. Overall, associations between behavioural risk factors and BMI become stronger with age and show similar patterns across regions. Though often reproduced across European regions and across cohorts with different data collection methods, the strength of association between each behavioural risk factor and BMI can be considered small. Associations that were inconsistent with the literature or varied across regions require further research to determine whether the collected data is an adequate proxy of the risk factor that is trying to be measured. Whilst all well-established healthy behaviours should be encouraged, regional and socioeconomic factors should be considered on an individual level when mitigating each risk factor.

AUTHOR CONTRIBUTIONS

Anton Schreuder, Claudia Börnhorst and Toomas Veidebaum contributed to conceptualisation. Anton Schreuder and Claudia Börnhorst contributed to the methodology. Anton Schreuder and Claudia Börnhorst contributed to the formal analysis. Claudia Börnhorst, Maïke Wolters, Toomas Veidebaum, Michael Tornaritis, Elida Sina, Paola

Russo, Luis A. Moreno, Denes Molnar, Lauren Lissner, Stefaan De Henauw, Wolfgang Ahrens and Toomas Veidebaum contributed to data acquisition. Anton Schreuder, Claudia Børnhorst and Toomas Veidebaum contributed to the first draft of the manuscript. All authors contributed to reviewing and editing the manuscript. All authors have read and agreed to the final version of the manuscript.

ACKNOWLEDGEMENTS

The authors warmly thank the participating parents and children, and all others who contributed to the IDEFICS, I.Family and ABCD studies. The authors also thank the Research Priority Area Yield, University of Amsterdam.

FUNDING INFORMATION

The GrowH! Project is funded by the Joint Programming Initiative 'A Healthy Diet for a Healthy Life' (JPI HDHL), a research and innovation initiative of EU member states and associated countries. The funding agencies supporting this work are (in alphabetical order of participating countries): Belgium: Research Foundation—Flanders (FWO); Germany: Federal Ministry of Education and Research (BMBF; grant no. 01EA2102A); Spain: Spanish State Research Agency (AEI); The Netherlands: The Netherlands Organisation for Health Research and Development (ZonMw). This project has received funding from the European Union's Horizon 2020 research and innovation programme under the ERA-NET Cofund action No. 727565. The data have been made available for this publication by IDEFICS (<http://www.idefics.eu>) and I.Family studies (<http://www.ifamilystudy.eu/>). The IDEFICS Study was funded by the European Community within the Sixth RTD Framework Programme Contract No. 016181 (FOOD). The I.Family Study was funded by the European Community within the Seventh RTD Framework Programme Contract No. 266044. The ABCD study was financially supported by the Lifestyle Innovations Based on Youth Knowledge and Experience (LIKE) programme, a grant from the Netherlands Cardiovascular Research Initiative, an initiative supported by the Dutch Heart Foundation, ZonMw, CVON2016-07 LIKE, and supported by Sarphati Amsterdam.

CONFLICT OF INTEREST STATEMENT

No conflict of interest was declared.

DATA AVAILABILITY STATEMENT

The IDEFICS and I.Family studies data that support the findings of this study are not publicly available because they contain information that could compromise the quality of the study, but the data sharing committee is willing to receive requests. Interested researchers can contact the IDEFICS and I.Family consortia (<http://www.ideficsstudy.eu/idefics/> and <http://www.ifamilystudy.eu/>) to discuss the possibilities for data access. The individual ABCD study data are not available for a public repository for ethical reasons but can be made available to other researchers for purposes of reproducing results or for collaboration. Researchers wishing to apply for the data can contact the project leader of the ABCD study (abcd@amsterdamumc.nl).

ORCID

Anton Schreuder  <https://orcid.org/0000-0002-6602-1977>
 Maïke Wolters  <https://orcid.org/0000-0002-4943-2141>
 Luis A. Moreno  <https://orcid.org/0000-0003-0454-653X>
 Lauren Lissner  <https://orcid.org/0000-0002-8296-2849>
 Tanja Vrijkotte  <https://orcid.org/0000-0003-3641-4048>

REFERENCES

1. Abarca-Gómez L, Abdeen ZA, Hamid ZA, et al. Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128.9 million children, adolescents, and adults. *Lancet*. 2017; 390(10113):2627-2642. doi:10.1016/S0140-6736(17)32129-3
2. Williams EP, Mesidor M, Winters K, Dubbert PM, Wyatt SB. Overweight and obesity: prevalence, consequences, and causes of a growing public health problem. *Curr Obes Rep*. 2015;4(3):363-370. doi:10.1007/s13679-015-0169-4
3. Lindberg L, Danielsson P, Persson M, Marcus C, Hagman E. Association of childhood obesity with risk of early all-cause and cause-specific mortality: a Swedish prospective cohort study. *PLoS Med*. 2020;17(3):e1003078. doi:10.1371/journal.pmed.1003078
4. Lindberg L, Hagman E, Danielsson P, Marcus C, Persson M. Anxiety and depression in children and adolescents with obesity: a nationwide study in Sweden. *BMC Med*. 2020;18(1):30. doi:10.1186/s12916-020-1498-z
5. van der Horst K, Oenema A, Ferreira I, et al. A systematic review of environmental correlates of obesity-related dietary behaviors in youth. *Health Educ Res*. 2006;22(2):203-226. doi:10.1093/her/cyl069
6. Blundell JE, Baker JL, Boyland E, et al. Variations in the prevalence of obesity among European countries, and a consideration of possible causes. *Obes Facts*. 2017;10(1):25-37. doi:10.1159/000455952
7. Romieu I, Dossus L, Barquera S, et al. Energy balance and obesity: what are the main drivers? *Cancer Causes Control*. 2017;28(3):247-258. doi:10.1007/s10552-017-0869-z
8. Silventoinen K, Jelenkovic A, Sund R, et al. Genetic and environmental variation in educational attainment: an individual-based analysis of 28 twin cohorts. *Sci Rep*. 2020;10(1):12681. doi:10.1038/s41598-020-69526-6
9. Tognon G, Moreno LA, Mouratidou T, et al. Adherence to a Mediterranean-like dietary pattern in children from eight European countries. The IDEFICS study. *Int J Obes (Lond)*. 2014;38(S2):S108-S114. doi:10.1038/ijo.2014.141
10. Tognon G, Hebestreit A, Lanfer A, et al. Mediterranean diet, overweight and body composition in children from eight European countries: cross-sectional and prospective results from the IDEFICS study. *Nutr Metab Cardiovasc Dis*. 2014;24(2):205-213. doi:10.1016/j.numecd.2013.04.013
11. Tran E, Dale HF, Jensen C, Lied GA. Effects of plant-based diets on weight status: a systematic review. *Diabetes Metab Syndr Obes*. 2020; 13:3433-3448. doi:10.2147/DMSO.S272802
12. Ridgers ND, Salmon J, Timperio A. Too hot to move? Objectively assessed seasonal changes in Australian children's physical activity. *Int J Behav Nutr Phys Act*. 2015;12(1):77. doi:10.1186/s12966-015-0245-x
13. Gracia-Marco L, Ortega FB, Ruiz JR, et al. Seasonal variation in physical activity and sedentary time in different European regions. The HELENA study. *J Sports Sci*. 2013;31(16):1831-1840. doi:10.1080/02640414.2013.803595
14. Børnhorst C, Siani A, Russo P, et al. Early life factors and inter-country heterogeneity in BMI growth trajectories of European children: the IDEFICS Study. *PLoS One*. 2016;11(2):e0149268. doi:10.1371/journal.pone.0149268
15. Vrijkotte TGM, Oostvogels AJJM, Stronks K, Roseboom TJ, Hof MHP. Growth patterns from birth to overweight at age 5-6 years

- of children with various backgrounds in socioeconomic status and country of origin: the ABCD study. *Pediatr Obes*. 2020;15(8):e12635. doi:[10.1111/ijpo.12635](https://doi.org/10.1111/ijpo.12635)
16. Gademan MGJ, Vermeulen M, Oostvogels AJM, et al. Maternal pre-pregnancy BMI and lipid profile during early pregnancy are independently associated with offspring's body composition at age 5–6 years: the ABCD study. *PLoS One*. 2014;9(4):e94594. doi:[10.1371/journal.pone.0094594](https://doi.org/10.1371/journal.pone.0094594)
 17. de Hoog MLA, Stronks K, van Eijnden M, Gemke RJB, Vrijkotte TGM. Ethnic differences in maternal underestimation of offspring's weight: the ABCD study. *Int J Obes (Lond)*. 2012;36(1):53–60. doi:[10.1038/ijo.2011.199](https://doi.org/10.1038/ijo.2011.199)
 18. Sirkka O, Hof MH, Vrijkotte T, et al. Feeding patterns and BMI trajectories during infancy: a multi-ethnic, prospective birth cohort. *BMC Pediatr*. 2021;21(1):34. doi:[10.1186/s12887-020-02456-4](https://doi.org/10.1186/s12887-020-02456-4)
 19. Börnhorst C, Russo P, Veidebaum T, et al. The role of lifestyle and non-modifiable risk factors in the development of metabolic disturbances from childhood to adolescence. *Int J Obes (Lond)*. 2020;44(11):2236–2245. doi:[10.1038/s41366-020-00671-8](https://doi.org/10.1038/s41366-020-00671-8)
 20. Turchin MC, Chiang CW, Palmer CD, Sankararaman S, Reich D, Hirschhorn JN. Evidence of widespread selection on standing variation in Europe at height-associated SNPs. *Nat Genet*. 2012;44(9):1015–1019. doi:[10.1038/ng.2368](https://doi.org/10.1038/ng.2368)
 21. Alderman H, Behrman JR, Glewwe P, Fernald L, Walker S. Evidence of impact of interventions on growth and development during early and middle childhood. *Child and Adolescent Health and Development. Disease Control Priorities*. Vol 8. Third ed. The World Bank; 2017:79–98. doi:[10.1596/978-1-4648-0423-6_ch7](https://doi.org/10.1596/978-1-4648-0423-6_ch7)
 22. Howe LD, Firestone R, Tilling K, Lawlor DA. *Trajectories and Transitions in Childhood and Adolescent Obesity*. Springer; 2015:19–37. doi:[10.1007/978-3-319-20484-0_2](https://doi.org/10.1007/978-3-319-20484-0_2)
 23. Hebestreit A, Barba G, de Henauw S, et al. Cross-sectional and longitudinal associations between energy intake and BMI z-score in European children. *Int J Behav Nutr Phys Act*. 2016;13(1):23. doi:[10.1186/s12966-016-0344-3](https://doi.org/10.1186/s12966-016-0344-3)
 24. Tahir MJ, Willett W, Forman MR. The association of television viewing in childhood with overweight and obesity throughout the life course. *Am J Epidemiol*. 2019;188(2):282–293. doi:[10.1093/aje/kwy236](https://doi.org/10.1093/aje/kwy236)
 25. Snell EK, Adam EK, Duncan GJ. Sleep and the body mass index and overweight status of children and adolescents. *Child Dev*. 2007;78(1):309–323. doi:[10.1111/j.1467-8624.2007.00999.x](https://doi.org/10.1111/j.1467-8624.2007.00999.x)
 26. Laverty AA, Hone T, Goodman A, Kelly Y, Millett C. Associations of active travel with adiposity among children and socioeconomic differentials: a longitudinal study. *BMJ Open*. 2021;11(1):e036041. doi:[10.1136/bmjopen-2019-036041](https://doi.org/10.1136/bmjopen-2019-036041)
 27. Hemmingsson E. Early childhood obesity risk factors: socioeconomic adversity, family dysfunction, offspring distress, and junk food self-medication. *Curr Obes Rep*. 2018;7(2):204–209. doi:[10.1007/s13679-018-0310-2](https://doi.org/10.1007/s13679-018-0310-2)
 28. Ahrens W, Bammann K, Siani A, et al. The IDEFICS cohort: design, characteristics and participation in the baseline survey. *Int J Obes (Lond)*. 2011;35(S1):S3–S15. doi:[10.1038/ijo.2011.30](https://doi.org/10.1038/ijo.2011.30)
 29. Ahrens W, Siani A, Adan R, et al. Cohort profile: the transition from childhood to adolescence in European children—how I.Family extends the IDEFICS cohort. *Int J Epidemiol*. 2016;46:dyw317. doi:[10.1093/ije/dyw317](https://doi.org/10.1093/ije/dyw317)
 30. van Eijnden M, Vrijkotte TG, Gemke RJ, van der Wal MF. Cohort profile: the Amsterdam born children and their development (ABCD) study. *Int J Epidemiol*. 2011;40(5):1176–1186. doi:[10.1093/ije/dyq128](https://doi.org/10.1093/ije/dyq128)
 31. Vrijkotte TGM, Varkevisser TMCK, van Schalkwijk DB, Hartman MA. Maternal underestimation of child's weight at pre-school age and weight development between age 5 and 12 years: the ABCD-study. *Int J Environ Res Public Health*. 2020;17(14):5197. doi:[10.3390/ijerph17145197](https://doi.org/10.3390/ijerph17145197)
 32. Cole TJ, Lobstein T. Extended international (IOTF) body mass index cut-offs for thinness, overweight and obesity. *Pediatr Obes*. 2012;7(4):284–294. doi:[10.1111/j.2047-6310.2012.00064.x](https://doi.org/10.1111/j.2047-6310.2012.00064.x)
 33. UNESCO Institute for Statistics. *International Standard Classification of Education: ISCED 2011*. UNESCO Institute for Statistics; 2012.
 34. Pigeot I, Baranowski T, De Henauw S. The IDEFICS intervention trial to prevent childhood obesity: design and study methods. *Obes Rev*. 2015;16:4–15. doi:[10.1111/obr.12345](https://doi.org/10.1111/obr.12345)
 35. De Henauw S, Huybrechts I, De Bourdeaudhuij I, et al. Effects of a community-oriented obesity prevention programme on indicators of body fatness in preschool and primary school children. Main results from the IDEFICS study. *Obes Rev*. 2015;16:16–29. doi:[10.1111/obr.12346](https://doi.org/10.1111/obr.12346)
 36. De Bourdeaudhuij I, Verbestel V, De Henauw S, et al. Behavioral effects of a community-oriented setting-based intervention for prevention of childhood obesity in eight European countries. Main results from the IDEFICS study. *Obes Rev*. 2015;16:30–40. doi:[10.1111/obr.12347](https://doi.org/10.1111/obr.12347)
 37. van Buuren S, Groothuis-Oudshoorn K. mice: Multivariate imputation by chained equations in R. *J Stat Softw*. 2011;45(3):1–67. doi:[10.18637/jss.v045.i03](https://doi.org/10.18637/jss.v045.i03)
 38. Huque MH, Moreno-Betancur M, Quartagno M, Simpson JA, Carlin JB, Lee KJ. Multiple imputation methods for handling incomplete longitudinal and clustered data where the target analysis is a linear mixed effects model. *Biom J*. 2020;62(2):444–466. doi:[10.1002/bimj.201900051](https://doi.org/10.1002/bimj.201900051)
 39. R Core Team. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria; 2020.
 40. Powell MJ. The BOBYQA algorithm for bound constrained optimization without derivatives. *Cambridge NA Report NA2009/06*. University of Cambridge; 2009.
 41. Tennant PWG, Murray EJ, Arnold KF, et al. Use of directed acyclic graphs (DAGs) to identify confounders in applied health research: review and recommendations. *Int J Epidemiol*. 2021;50(2):620–632. doi:[10.1093/ije/dyaa213](https://doi.org/10.1093/ije/dyaa213)
 42. Atkin AJ, Sharp SJ, Corder K, van Sluijs EMF. Prevalence and correlates of screen time in youth. *Am J Prev Med*. 2014;47(6):803–807. doi:[10.1016/j.amepre.2014.07.043](https://doi.org/10.1016/j.amepre.2014.07.043)
 43. Rittsteiger L, Hinz T, Oriwol D, Wäsche H, Santos-Hövenner C, Woll A. Sports participation of children and adolescents in Germany: disentangling the influence of parental socioeconomic status. *BMC Public Health*. 2021;21(1):1446. doi:[10.1186/s12889-021-11284-9](https://doi.org/10.1186/s12889-021-11284-9)
 44. Müller MJ, Koertzing I, Mast M, Langnase K, Grund A. Physical activity and diet in 5 to 7 years old children. *Public Health Nutr*. 1999;2(3a):443–444. doi:[10.1017/S1368980099000609](https://doi.org/10.1017/S1368980099000609)
 45. Pearson N, Salmon J, Campbell K, Crawford D, Timperio A. Tracking of children's body-mass index, television viewing and dietary intake over five-years. *Prev Med*. 2011;53(4–5):268–270. doi:[10.1016/j.yjmed.2011.07.014](https://doi.org/10.1016/j.yjmed.2011.07.014)
 46. Grimes CA, Riddell LJ, Campbell KJ, Nowson CA. Dietary salt intake, sugar-sweetened beverage consumption, and obesity risk. *Pediatrics*. 2013;131(1):14–21. doi:[10.1542/peds.2012-1628](https://doi.org/10.1542/peds.2012-1628)
 47. Patelakis E, Lage Barbosa C, Haftenberger M, et al. Prevalence of vegetarian diet among children and adolescents in Germany. *Ernahr Umsch*. 2019;66(5):85–91.
 48. González SA, Aubert S, Barnes JD, Larouche R, Tremblay MS. Profiles of active transportation among children and adolescents in the global matrix 3.0 initiative: a 49-country comparison. *Int J Environ Res Public Health*. 2020;17(16):5997. doi:[10.3390/ijerph17165997](https://doi.org/10.3390/ijerph17165997)
 49. Franse CB, Wang L, Constant F, Fries LR, Raat H. Factors associated with water consumption among children: a systematic review. *Int J Behav Nutr Phys Act*. 2019;16(1):64. doi:[10.1186/s12966-019-0827-0](https://doi.org/10.1186/s12966-019-0827-0)
 50. Mantziki K, Renders C, Seidell J. Water consumption in European children: associations with intake of fruit juices, soft drinks and related

- parenting practices. *Int J Environ Res Public Health*. 2017;14(6):583. doi:[10.3390/ijerph14060583](https://doi.org/10.3390/ijerph14060583)
51. Chung BW. Peers' parents and educational attainment: the exposure effect. *Labour Econ*. 2020;64:101812. doi:[10.1016/j.labeco.2020.101812](https://doi.org/10.1016/j.labeco.2020.101812)
 52. Bogl LH, Mehlig K, Ahrens W, et al. Like me, like you – relative importance of peers and siblings on children's fast food consumption and screen time but not sports club participation depends on age. *Int J Behav Nutr Phys Act*. 2020;17(1):50. doi:[10.1186/s12966-020-00953-4](https://doi.org/10.1186/s12966-020-00953-4)
 53. Börnhorst C, Huybrechts I, Ahrens W, et al. Prevalence and determinants of misreporting among European children in proxy-reported 24 h dietary recalls. *Br J Nutr*. 2013;109(7):1257-1265. doi:[10.1017/S0007114512003194](https://doi.org/10.1017/S0007114512003194)
 54. Koning M, de Jong A, de Jong E, Visscher TLS, Seidell JC, Renders CM. Agreement between parent and child report of physical activity, sedentary and dietary behaviours in 9-12-year-old children and associations with children's weight status. *BMC Psychol*. 2018; 6(1):14. doi:[10.1186/s40359-018-0227-2](https://doi.org/10.1186/s40359-018-0227-2)
 55. Forrestal SG. Energy intake misreporting among children and adolescents: a literature review. *Matern Child Nutr*. 2011;7(2):112-127. doi:[10.1111/j.1740-8709.2010.00270.x](https://doi.org/10.1111/j.1740-8709.2010.00270.x)
 56. Zhu W, Marchant R, Morris RW, Baur LA, Simpson SJ, Cripps S. Bayesian network modelling to identify on-ramps to childhood obesity. *BMC Med*. 2023;21(1):105. doi:[10.1186/s12916-023-02789-8](https://doi.org/10.1186/s12916-023-02789-8)
 57. Revenson TA, Abraído-Lanza AF, Majerovitz SD, Jordan C. Couples coping with chronic illness: what's gender got to do with it? *Couples Coping with Stress: Emerging Perspectives on Dyadic Coping*. American Psychological Association; 2005:137-156. doi:[10.1037/11031-007](https://doi.org/10.1037/11031-007)
 58. Bianchi SM, Sayer LC, Milkie MA, Robinson JP. Housework: who did, does or will do it, and how much does it matter? *Soc Forces*. 2012; 91(1):55-63. doi:[10.1093/sf/sos120](https://doi.org/10.1093/sf/sos120)
 59. Gluckman PD, Buklijas T, Hanson MA. The developmental origins of health and disease (DOHaD) concept. *The Epigenome and Developmental Origins of Health and Disease*. Elsevier; 2016:1-15. doi:[10.1016/B978-0-12-801383-0.00001-3](https://doi.org/10.1016/B978-0-12-801383-0.00001-3)
 60. Tan X, Chapman CD, Cedernaes J, Benedict C. Association between long sleep duration and increased risk of obesity and type 2 diabetes: a review of possible mechanisms. *Sleep Med Rev*. 2018;40:127-134. doi:[10.1016/j.smrv.2017.11.001](https://doi.org/10.1016/j.smrv.2017.11.001)
 61. Lucan SC, DiNicolantonio JJ. How calorie-focused thinking about obesity and related diseases may mislead and harm public health. An alternative. *Public Health Nutr*. 2015;18(4):571-581. doi:[10.1017/S1368980014002559](https://doi.org/10.1017/S1368980014002559)
 62. García-Blanco L, Berasaluce A, Romanos-Nanclares A, Martínez-González MÁ, Moreno-Galarraga L, Martín-Calvo N. Parental perception of child's weight, their attitudes towards child's dietary habits and the risk of obesity. *World J Pediatr*. 2022;18(7):482-489. doi:[10.1007/s12519-022-00540-6](https://doi.org/10.1007/s12519-022-00540-6)
 63. Pizarro AN, Santos MP, Ribeiro JC, Mota J. Physical activity and active transport are predicted by adolescents' different built environment perceptions. *J Public Health*. 2012;20(1):5-10. doi:[10.1007/s10389-011-0432-4](https://doi.org/10.1007/s10389-011-0432-4)
 64. Fritz CO, Morris PE, Richler JJ. Effect size estimates: current use, calculations, and interpretation. *J Exp Psychol Gen*. 2012;141(1):2-18. doi:[10.1037/a0024338](https://doi.org/10.1037/a0024338)
 65. Smahel D, Machackova H, Mascheroni G, et al. *EU Kids Online 2020: Survey Results from 19 Countries*. EU Kids Online; 2020.
 66. Andone I, Błaszczewicz K, Eibes M, Trendafilov B, Montag C, Markowetz A. How age and gender affect smartphone usage. *Proceedings of the 2016 ACM International Joint Conference on Pervasive and Ubiquitous Computing: Adjunct*. ACM; 2016:9-12. doi:[10.1145/2968219.2971451](https://doi.org/10.1145/2968219.2971451)

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Schreuder A, Börnhorst C, Wolters M, et al. Population trajectories and age-dependent associations of obesity risk factors with body mass index from childhood to adolescence across European regions: A two-cohort study. *Pediatric Obesity*. 2024;19(2):e13088. doi:[10.1111/ijpo.13088](https://doi.org/10.1111/ijpo.13088)