



Comparative genomics of *Staphylococcus aureus* strains from wild birds and pig farms elucidates levels of mobilomes, antibiotic pressure and host adaptation

Idris Nasir Abdullahi^a, Carmen Lozano^a, Myriam Zarazaga^a, Carmen Simón^b, Ursula Höfle^c, Raphael N. Sieber^d, Javier Latorre-Fernández^a, Marc Stegger^{d,e}, Carmen Torres^{a,*}

^a Area of Biochemistry and Molecular Biology, OneHealth-UR Research Group, University of La Rioja, Logroño, Spain

^b Faculty of Veterinary Medicine, University of Zaragoza, Zaragoza, Spain

^c Spanish Wildlife Research Institute IREC (CSIC–UCLM–JCCM), SaBio (Health and Biotechnology) Research Group, Ciudad Real, Spain

^d Department of Bacteria, Parasites and Fungi, Statens Serum Institut, Copenhagen, Denmark

^e Antimicrobial Resistance and Infectious Diseases Laboratory, Harry Butler Institute, Murdoch University, Murdoch, Western Australia, Australia

ARTICLE INFO

Article history:

Received 12 October 2023

Revised 1 December 2023

Accepted 8 December 2023

Available online 19 December 2023

Editor: Stefania Stefani

Keywords:

S. aureus evolution

Host adaptation

Antibiotic pressure

Mobilome

ABSTRACT

Objectives: This study characterized the resistome, mobilome and phylogenomic relatedness of *Staphylococcus aureus* strains previously obtained from healthy nestling storks (HNS), pigs (HP) and pig farmers (HPF) to analyse possible transmission pathways of *S. aureus* with implications for the spread of antimicrobial resistance.

Methods: The genomic contents of 52 *S. aureus* strains obtained from the nasal cavity of HNS, HP and HPF in Spain were sequenced using the Illumina NextSeq platform to characterize their resistome, virulome and mobile genetic elements. The relatedness of strains was assessed by core-genome single nucleotide polymorphisms (SNPs).

Results: The frequencies of multidrug-resistance phenotype and transposons were significantly lower in strains from HNS than in those from HP and HPF ($P < 0.005$). However, the presence of human immune evasion cluster genes in *S. aureus* strains from HNS was significantly higher than in those from HP and HPF ($P < 0.005$). Interestingly, the frequencies of plasmids and phages were not significantly associated with the host ($P > 0.05$). The phylogenetic analysis identified a cluster of all the MSSA-CC398 strains carrying ϕ Sa3 and *ermT* on *rep13* separately from the two MRSA-CC398 strains (carrying *ermT* on *repUS18*). Highly related MRSA-CC398 strains were detected in some pigs and related farmers (<10 SNPs).

Conclusion: This study confirms high-level antibiotic selection in *S. aureus* in HP and HPF in comparison to HNS. Furthermore, our findings highlight the continuous transmission of MRSA-CC398 in the pig-to-human interface and MSSA-CC398 with human adaptation markers in HNS. Molecular surveillance of *S. aureus* using the One Health model is required to establish appropriate control strategies.

© 2023 The Authors. Published by Elsevier Ltd on behalf of International Society for Antimicrobial Chemotherapy.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

1. Introduction

Antimicrobial resistance (AMR) constitutes one of the major global health challenges that need a holistic ‘OneHealth’ approach. In this regard, *Staphylococcus aureus* is one of the suitable bacteria, as certain genetic lineages can cross host species barriers and transfer AMR genes among humans, animals and their shared envi-

ronment [1]. The emergence and spread of methicillin-resistant *S. aureus* (MRSA) is often blamed on the overuse and overprescription of antibiotics in human and veterinary medicine and livestock production [2]. On the other side, wild animals generally carry low-level AMR in commensal bacteria [3,4], but some of them, such as migratory birds that forage in areas close to anthroponotic activities or livestock grazing areas, could be colonized by MRSA [5].

S. aureus is a multi-host bacterium and is generally a significant component of the nasal and skin microbiota of humans and animals (including wild animals) [6–9]. However, it can also causes various infections in humans and animals [10,11]. *S. aureus* is an

* Corresponding author. Mailing address: Area of Biochemistry and Molecular Biology, University of La Rioja, 26006 Logroño, Spain.

E-mail address: carmen.torres@unirioja.es (C. Torres).

asymptomatic colonizer in pigs, but it can cause disease in other livestock, such as mastitis in dairy bovines [12,13]. AMR, virulence and host adaptation in *S. aureus* are relevant features in its pathogenesis and evolution [14]. Molecular studies have shown that livestock can act as intermittent carriers of *S. aureus* and reservoirs for zoonosis and dissemination of high-level AMR in farmers, the communities close to the farms, and food derived from the animals [15–19]. In this regard, the major livestock-adapted lineage (MRSA-CC398) emerged from human-adapted MSSA-CC398 after the loss of ϕ Sa3 and the acquisition of *tet*(M) and *SCCmec* elements and spread in the European pig production [20,21], with subsequent spillover back into humans in the community and healthcare settings [19,22]. Years later, the human-adapted and community-associated MSSA-CC398 reemerged to cause invasive infections in some European countries and China [23–26]. Epidemiologically, the *scn* and *tet*(M) genes are considered molecular markers that can be used to track these lineages [21]. However, fundamental questions that need to be elucidated are the molecular markers of host-switch and virulence level of MRSA-CC398 when they are in non-livestock hosts such as humans and other animals living close to pig farms.

Recently, there has been much interest but scarce data about the global epidemiology of MRSA in wildlife [27]. MRSA could be transmitted by anthropogenic or livestock activities which have been hypothesized to be the primary link to wild animals [8]. As MRSA gets transferred to wild animals, it can be responsible for the spread of AMR genes through mobile genetic elements [27]. Consequently, these underscore the need for global and elaborate studies on wild animals, to direct the best strategies for the control against the spread of MRSA across ecosystems and to limit the global emergence of AMR traits.

Intra-species AMR and lineage diversity (*S. aureus* strains from a single host with more than one AMR profile or lineage at a given point in time) could provide an insight into the dynamics of AMR, inter-host transmission and inter-ecological transmission of *S. aureus* [28]. Few comparative genomic studies have determined the effect of hosts, habitat, ecology and human occupation on antibiotic selection pressure and associated mobile genetic elements. The hosts from the two ecological niches (wildlife and pig farms) were selected in this study due to their distinct differences concerning antimicrobial pressures.

Here, we conducted a comparative genome-based study of *S. aureus* obtained from nestling white storks, healthy pigs and pig farmers to elucidate levels of AMR, mobile genetic elements, and their host adaptation profile to get insight into possible transmission pathways of *S. aureus* with implications for the spread of AMR.

2. Materials and methods

2.1. *Staphylococcus aureus* strains in this study

Fifty-two *S. aureus* strains obtained in previous studies [29–31] from 87 nestlings of white stork parents foraging in natural areas ($n = 5$) and landfills ($n = 18$), and from healthy pigs ($n = 17$) and pig farmers ($n = 12$) from four pig farms (A–D), were subjected to whole genome sequencing (Table 1). One *S. aureus* strain from each of the nestling storks was selected, although more than one strain was included when they presented different clonal complexes (CC). Concerning the *S. aureus* strains from the pig farms (A to D), the selection was based on similarity in their sequence type and AMR genes in each farm and the similarity of genetic lineages and AMR genes from strains of pigs and pig farmers as previously determined [29]. Multidrug resistance (MDR) was defined when a strain carried genes that mediate resistance to ≥ 3 classes of antimicrobial agents [32]. Comprehensive information on the sample collection and processing procedures, bacterial isolation and identi-

fication, *S. aureus* antimicrobial susceptibility and AMR genotyping are presented in previous studies [29–31]. All the study protocols have been reviewed and approved by the ethical research committees of the University of Zaragoza, the University of La Rioja and the University of Castilla La Mancha (Spain).

2.2. Genome sequencing, assembly and phylogenetic analyses

Whole genome sequencing of the 52 selected *S. aureus* strains was carried out on the Illumina NextSeq platform. The MagNA Pure 96 DNA Multi-Sample Kit (Life Technologies, Carlsbad, CA, USA, 4413021) was used to extract genomic DNA according to instructions provided by the manufacturers. The Qubit 1X dsDNA HS Assay Kit (Thermo Fisher Scientific, Scoresby, VIC, Australia) was used for DNA quantification, and sequencing libraries were prepared using the Illumina Nextera XT DNA Library Preparation Kit (Illumina, San Diego, CA, USA, FC-131-1096) and sequenced on the NextSeq 500 platform (Illumina, San Diego, CA) using a 300-cycle kit to obtain paired-end 150 bp reads, as previously described [33].

All the genomes analysed in this study were *de novo* assembled using SPAdes (v.3.15.5). Core-genome single nucleotide polymorphisms (SNPs) between the 52 *S. aureus* strains in this study were detected with the NASP pipeline (v.1.0.0) [34] and used to reconstruct their phylogenetic relationship. Briefly, the raw sequencing data from all CC398 strains were mapped together with 88 previously published *S. aureus* CC398 genomes (Bioproject number: PRJNA514245) against the chromosome of ST398 strain S0385 (GenBank accession no. AM990992) as a reference to obtain a CC398 phylogeny. GATK (v.4.2.2) was used to call SNPs and excluded positions featuring $<90\%$ unambiguous variant calls and <10 depth. IQ-TREE (v.2.1.2) was used to construct the phylogenetic trees using ModelFinder with 100 bootstrap replicates. The graphical data were added to the phylogenies with iTOL (v.6.6) [35]. Supplementary Table S1 shows the SNPs difference between the *S. aureus* CC398 strains from this study (both MRSA and MSSA) and 89 publicly available genomes of *S. aureus* CC398 strains from Price et al. [21].

2.3. Genome annotation, typing and in silico analysis

The sequence types (STs) were determined with MLST v.2.16 [36]. Virulence factors, plasmid replicons and antimicrobial resistance genes were identified using ABRicate (v.0.9.0) (<https://github.com/tseemann/abricate>) and the respective databases VFDB, Plasmidfinder and Resfinder databases from the Center for Genomic Epidemiology (<http://www.genomicepidemiology.org>). Mutations associated with AMR were identified using ResFinder (v.4.1) [37] and PointFinder [38]. Biocide and heavy metal resistance genes were identified using BACMET [39]. PHASTER was used to identify all prophage elements [40]. Moreover, manual mapping of raw sequencing reads using the KMA algorithm [41] with a minimum of 90% coverage, 80% identity and minimum $6 \times$ depth was performed on sets of prophage integrases and IEC genes and their corresponding GenBank accession numbers: Sa12int (NC_010147), Sa9int (NC_007057), Sa8int (NC_007622), Sa7int (NC_007049), Sa6int (NC_005356), Sa6int (M27965), Sa5int (NC_004615), Sa4int (NC_002953), Sa3int (NC_009641), Sa3int (NC_004617), Sa3int (DQ530361), Sa2int (NC_004616), Sa2int (NC_002321), Sa1int (NC_003288), phiJB_int (NC_028669), Se-bagoInt (MK618716), IEC_chp (NC_009641), IEC_sak (NC_009641), IEC_scn (NC_009641), IEC_sea (NC_009641) and IEC_sep (BA000018). The *spa*Typer (v.1.0) tool was used to confirm the *spa* types [42]. The *SCCmec* types were assigned using *SCCmecFinder* (v.1.2) (<https://cge.food.dtu.dk/services/SCCmecFinder/>). The genetic environment of the *ermT* gene was illustrated in comparison with the reference strains; MRSA pUR1902 (GenBank accession

Table 1
Characteristics of *S. aureus* strains from the nestling storks, pigs and pig farmers.

Variables	Nestling storks from parents foraging in landfills (n = 18)	Nestling storks from parents foraging in natural areas (n = 5)	Pigs (n =17)	Pig farmers (n 12)
No. of strains with methicillin resistance trait	0	0	14	12
No. of strains with MDR phenotype	1	0	17	12
<i>spa</i> types	t015, t127, t209, t223, t227, t521, t335, t571, t774, t1094, t1451, t1654, t3380, t7778, t18009	t571, t1451, t2313, t3380 t6220	t011, t034, t1430, t1451	t011, t034, t1451
ST	ST1, ST5, ST15, ST22, ST25, ST26, ST30, ST45, ST97, ST109, ST398	ST97, ST291, ST130, ST398	ST9, ST398	ST398
CC	CC1, CC5, CC9, CC15, CC22, CC25, CC30, CC45, CC97, CC398	CC97, CC130, CC398	CC9, CC398	CC398

number: HF583291.1), MRSA pUR2941 (GenBank accession number: HF583290.1) and MRSA AV4_1 (GenBank accession number: SAMN00828682) using EasyFig software.

2.4. Genome availability

All the raw genome reads generated from this study have been deposited at the European Nucleotide Archive under study accession no. PRJEB66351.

3. Results and discussion

3.1. Genetic relatedness of the *S. aureus* strains

The genomes of the 52 *S. aureus* strains revealed that pig and pig farmer strains belonged mainly to MRSA-CC398, except for three MSSA-CC9, whereas stork nestling strains belonged to many different CCs and included MSSA-CC398 but not MRSA-CC398 (Table 1). All MSSA-CC9 from pigs clustered into the same cluster, which was different from the MSSA-CC9 strain from a nestling stork (Fig. 1). Furthermore, strains from nestlings of parents feeding in landfills were interspersed compared to those from nestlings of parents feeding in natural areas with no apparent difference in spillover patterns between the two groups.

The phylogenetic analysis of the CC398 strains furthermore showed that all MSSA-CC398 strains from storks (both from natural and landfill areas) clustered within the human clade, whereas the MRSA-CC398 strains from pigs and pig farmers clustered in the LA clade (Fig. 2). Furthermore, closely related MRSA-CC398 strains (<10 SNPs) carrying the same AMR profile were detected between pigs and pig farmers from the same farm (Supplementary Table S1), indicating within-farm transmission of these *S. aureus* strains. This suggests that pig farmers have contracted the LA MRSA-CC398 strains because of their occupation. However, nestling storks carried the MSSA-CC398-*scn*-positive subclade, which is known to frequently circulate among humans [25], and infrequently in livestock. In this regard, nestlings of parent storks that foraged in landfill areas with anthroponotic remains could be hypothesized to be the source of this subclade. However, some CC398- and non-CC398-*scn*-positive strains were obtained from nestlings of parent storks that foraged in natural areas where no or very limited traces of anthropogenic remains can be expected. The reason for this unexpected finding could be complex and related to multiple factors that need to be further studied.

3.2. Association of strains' origin with antibiotic pressure and mobilome levels

Bivariate logistic analysis showed that the frequencies of MDR phenotype and transposons were significantly lower in nesting

storks than in pigs and pig farmers ($P < 0.005$) (Supplementary Table S2). However, the presence of the *scn* gene (a marker of the IEC system) in *S. aureus* strains from nestling storks was significantly higher than in those of pigs and pig farmers ($P < 0.005$). The reason for the abundance of the *scn* gene among the storks' strains is not clear, as previously indicated, because it is considered a human adaptation marker. Some authors suggest that various prophages could be relevant determinants for *S. aureus* host switching, transmissibility, infection and adaptation [17]. Future studies focused on this issue could elucidate the mechanisms involved. On the other hand, the frequencies of plasmids and phages were not significantly associated with hosts ($P > 0.05$) (Supplementary Table S2).

3.3. Antimicrobial and metal resistance and associated mobile genetic elements

Generally, most of the *S. aureus* strains from nestling storks were either resistant to only one class of antibiotics (in most cases beta-lactam or macrolides) or entirely susceptible to all antibiotics. In this sense, the resistome profile of the strains was mainly *blaZ*, *ermT*, *lnuA* and *tet(K)* as previously detected by PCR [30]. In addition, we here further identified the *ant9'* gene in strain X3799, which was co-localized with *ermA* linked with a Tn554 transposon (Supplementary Table S3). This was similarly reported in a previous study on *Staphylococcus lugdunensis* [43,44]. To our knowledge, the linkage of the *ermA* gene in Tn554 has never been reported in MSSA, suggesting the expansion of the ecology of these genes outside methicillin-resistant *S. lugdunensis*. Moreover, *S. aureus* strain X4139 from a nestling of a parent stork foraging in landfills carried the *vga(A)V* gene, which we could not associate with any detectable mobile genetic element (Supplementary Table S3).

Supplementary Table S4 shows the complete resistome of the 35 *S. aureus* strains from pigs and pig farmers. All the *S. aureus* strains presented an MDR phenotype mediated by a large repertoire of resistance markers (often >5 different genes). In this sense, the resistome of these strains mainly consisted of genes that mediate resistance to beta-lactams, macrolide-lincomycin-streptogramin B (MLS_B), tetracyclines, aminoglycosides, sulfamethoxazole-trimethoprim or fluoroquinolones, as previously detected by PCR [29]. In addition, the aminoglycoside gene *str* that mediates resistance to streptomycin was detected in 12 strains. Also, *ant9'*, another aminoglycoside gene, was co-located with the *ermT* gene on plasmid *repUS18* in pig (X4905) and pig farmer (X5473) strains (Supplementary Table S4).

All the *S. aureus* strains from pigs (both MRSA and MSSA) presented a MDR phenotype, contrary to those from the nestling storks, where only one strain presented an MDR phenotype/genotype (X3799). This observation could be associated with the differences concerning antimicrobial pressures (high and low, respectively) between the two ecological niches. All the strains




Fig. 1. Phylogenetic tree based on core genome SNP analysis of 52 *S. aureus* strains to demonstrate the influence of antibiotic pressure based on the host.

from pigs and pig farmers presented the characteristic tetracycline resistance where 20 *S. aureus* strains from pigs and pig farmers carried *tet*(M), 12 carried *tet*(K) and 8 carried *tet*(L). It is important to highlight that all the *tet*(L)-carrying *S. aureus* strains were additionally *tet*(M)-positive. Interestingly, the *tet*(M) and *tet*(L) were located in plasmid *rep22* and *repUS43*, respectively, in all the MRSA-CC398 strains harbouring these genes. But *tet*(M)- and *tet*(L)-carrying MSSA-CC9 strains from the two pigs and a pig farmer did not harbour these plasmid replicons. Perhaps these plasmids are emerging mobilomes for *tet*(M)- and *tet*(L) genes in the MRSA-CC398 strains. The *tet*(K) gene in most of the *S. aureus* strains, including the one from a stork nestling (X4139), was located in *rep7a*, while only one strain (X5751) was not associated with this plasmid replicon. The *tet*(M) gene was localized in various transposons such as those detected in this study (i.e., Tn6009, Tn925 and Tn916). The Tn6009 is a member of the Tn916–Tn1545 family and is a conjugative (noncomposite) transposon of Tn916 [45]. The *tet*(M) has previously been reported to be carried by *repUS43* alongside Tn6009 in an *E. faecalis* strain [46], but to our knowledge not reported in *S. aureus*. This denotes the diversity of transposons associated with *tet*(M)-mediated tetracycline resistance and suggests their essential role in the evolution and horizontal spread of this resistance marker in *S. aureus* from pigs and pig farmers. However, the *tet*(L) is almost always linked with the *dfrK* gene on small plasmids [47].

Concerning all the strains carrying *tet*(L) and *dfrK* genes, the plasmid *rep22* was always co-located with these genes and is 99% similar to the plasmid pKKS2187 in an MRSA-CC398 strain from a pig (GenBank accession number: FM207105). Moreover, the *dfrK* gene was also found to be located in Tn558, which is often in-

tegrated into the *radC* gene in MRSA-CC398 strains [48]. Two *S. aureus* strains from nestling storks (X4135 and X4139) presented tetracycline resistance mediated by the *tet*(K) gene carried in circular plasmid *rep7a*, while the *tet*(M) and *tet*(L) genes were absent from all nestling stork strains. It appears that *tet*(M) (in both MRSA and MSSA) was strongly associated with tetracycline resistance in pigs and pig farmers, which is in line with previous findings by Price et al. [21].

Aside from tetracycline, resistance to other clinical antibiotics was also detected, such as to MLS_B antibiotics. AMR to this class of antibiotics was specifically mediated by *lnuB* and *lsaE* in nine *S. aureus* strains from the pigs and pig farmers. However, MLS_B resistance was mediated by *ermA* and *vga(A)*V among the two non-CC398 strains from nestling storks (X3799 and X4139). Also, the *vga(A)*LC gene is located on a *rep7b* in 10 strains (Supplementary Table S3). Furthermore, the *ermT* gene that mediates a peculiar MLS_B resistance (erythromycin-resistant/clindamycin-inducible resistance) was found in five MSSA-CC398 strains from nestling storks (Supplementary Table S3) and two MRSA-CC398 strains from a pig and a pig farmer. the *ermT*-positive-MRSA-CC398 strains are very unusual [49].

Another antibiotic of interest is chloramphenicol, which has long been prohibited for use in pig production, and neither is in use for chemotherapy of human infections in Spain. However, florfenicol (a closely related antibiotic) is approved by the USA Food and Drug Agency and the European Medicines Agencies for use in swine, cattle and sheep [50]. Thus, the detection of *fexA* encoding for chloramphenicol resistance illustrates its persistence due to the use of florfenicol in pig farms, as this was not present in any of the nestling stork strains.




Fig. 2. Circular phylogenetic tree to show the human- and animal-adapted subclades based on SNP analyses of CC398 strains obtained from this study in comparison with publicly available genomes of CC398 lineages from Price et al [21].

Finally, a striking difference in AMR between the strains from the two ecological niches was the ciprofloxacin resistance mediated by DNA topoisomerase IV point mutations at GyrA (S84L) and DNA gyrase at GrlA (S80F) or GrlA (S80Y). These common mutations were found in 16 (55.2%) of the *S. aureus* strains from pigs and pig farmers but not among those from the nestling storks. A similar observation was reported in a previous study on porcine MRSA strains in Germany and Denmark [51,52]. It has been speculated that the mutations in GrlA (S80F or S80Y) might affect recombination between the Φ Sa3 phage and *S. aureus* chromosome, which could contribute to the absence of this prophage in MRSA-CC398 from pigs and pig farmers, although an apparent link between fluoroquinolone resistance and integration or stability of Φ Sa3 phage could not be demonstrated [52].

Concerning heavy metal resistance genes, certain categorical differences in genes encoding resistance to copper and zinc were observed and widespread among the *S. aureus* strains from pigs and pig farmers, whereas they were absent in the nestling storks' strains. Zinc and copper are frequently used in animal feed formulations, and these metals may co-select for MRSA-CC398 strains that carry the *czrC* and *copA* genes [53,54]. Interestingly, only the *czrC* gene was detected among the MRSA-CC9 strains. The *czrC* gene is part of the SCCmec element of MRSA-CC398 [54], so probably the *czrC* in MRSA-CC9 has a different role as these strains do not contain the SCCmec elements. Moreover, the *copA* gene appears to be exclusively associated with MRSA strains. Apart from the *cadX* gene that was found only on the MRSA-CC398 subclade with

nestling stork strains, the *arsB* gene that encodes for arsenic pump membrane protein was found only in an MRSA-CC130 strain as previously described [55]. However, this gene has also been largely present in MRSA-CC130 strains of wild animals [27]. Moreover, the *cadD* (cadmium transport protein D) gene was carried by 15 (57.7%) MRSA strains (including CC398) from nestling storks. The presence of these genes suggests the existence of metal pollution in the environment where these strains originate from, which could be linked to human activities [56]. It is important to highlight that heavy metal resistance is a matter of public health concern due to its potential hazards in the food chain and the co-selection of AMR [56].

All the MRSA-CC398 strains of pigs and pig farmers carried the SCCmec type Vc (5C2&5). The vast majority of MRSA-CC398 strains carry this SCCmec type, which often contains the *czrC* and *copA* genes that detoxify cadmium, zinc and copper [57], of which the *copA* contributes to MRSA-CC398 survival in pig farm environment [58]. Moreover, the use of zinc oxide in livestock production has been found to be associated with the co-selection of MRSA-CC398 [54].

Concerning biocide resistances, only the *qacG* gene was detected in 20 (68.9%) *S. aureus* strains (MRSA-CC398 and MRSA-CC9), all from pigs and pig farmers. The acquisition of *qacG* carried on plasmid *rep21* could facilitate the persistence of these lineages in the pig/pig farmer environment, as they are resistant to quaternary ammonium disinfectants, thus making their eradication very difficult [59].



Fig. 3. Schematic comparison between the environment of the *ermT* gene in the five MSSA-CC398 and two MRSA-CC398 strains with the reference strain MRSA-pURX2941 (HF583290) and pURX1902 (Gene bank accession number: HF583291) and MRSA-AV_4 (Gene bank accession number: SAMN00828682). Colors and arrows indicate the represented genes and their orientation.

3.4. Genetic environment of the *ermT* gene in MSSA and MRSA strains

Despite the involvement of the *ermT* gene as a major mediator of inducible MLS_B resistance in MSSA-CC398 strains, its environment and its associated genes and mobilome have rarely been described. The *in silico* analysis of the *ermT* sequences of two MRSA strains from a pig (X4905) and a pig farmer (X5473) and five MSSA strains from nestling storks (X3906, X3913, X4603, X4630, X4703) revealed striking differences in their genetic environment (Figure 3). First, the five *ermT*-carrying MSSA strains were all associated with cadmium-resistance genes, *cadR* and *cadD*, which were absent from the two *ermT*-carrying MRSA strains. The IS257 was located upstream of the *ermT* gene in MSSA strains, except in one strain in which it was located downstream. Interestingly these

markers were absent in the *ermT*-carrying MRSA strains. The *ermT*-gene of the MSSA strains was associated with plasmid *rep13*. Often, the *ermT* gene in MSSA is carried by a plasmid and has recently been recognized as a biomarker of erythromycin-clindamycin inducible resistance in this lineage [25]. Contrary to this observation, the *ermT* gene in MRSA produces erythromycin-clindamycin constitutive resistance, a phenomenon previously reported in the literature [49].

3.5. Virulome profile of the *S. aureus* strains from nestling storks and pig farm hosts

Toxins constitute important virulence determinants of *S. aureus*, with enterotoxins being the most implicated in food safety, especially in meat and dairy products from livestock [60]. More-

over, other virulence factors could be responsible for a range of *S. aureus*-related infectious diseases [11]. The MRSA-CC398 strains were entirely negative for *tst*, *lukS/F-PV*, *eta*, *etb*, *etc*, *etd* and all genes encoding enterotoxins. This is in line with previous findings [25]. However, the *sem*, *seo*, *seu* and *sei* genes were identified in the three MSSA-CC9 strains from pigs. Moreover, *sen*, *sem*, *sei*, *seg*, *seu* and *seo* genes were also identified in four MSSA strains of the nestling storks belonging to CC5, CC9, CC25 and CC45 (Supplementary Table S3). From our findings, it appears that virulence genes are more predominant in MSSA than in MRSA strains.

Aside from the toxins, some *S. aureus* enzymes were commonly present in all the strains, such as the *adsA* that encodes adenosine synthase A, a cell-wall-anchored enzyme that converts adenosine monophosphate to adenosine and seems to be related to the evasion of host immune responses in *S. aureus* [61]. Furthermore, the *icaABCD* operon and its *icaR* regulatory gene were present in all the strains. This denotes that the *S. aureus* strains easily adhere to the mucosa and serve as a fundamental step in nasal colonization and persistence on environmental surfaces and fomites [62].

3.6. Host adaptation markers of *S. aureus*

Nineteen (82.6%) strains from the nestling storks carried the *scn* gene mediated by the prophage ϕ Sa3 (Supplementary Table S3), whereas all *S. aureus* (MRSA-CC398 and MSSA-CC9) from pigs and pig farmers lacked the prophage ϕ Sa3 marker (i.e., *scn*), indicating that they can be considered livestock-associated or originating from nonhuman hosts (Supplementary Table S4), since ϕ Sa3 plays a significant role in the host adaptation of *S. aureus* [14]. For CC398, previous studies divided the lineage into a human-adapted MSSA clade (*scn* positive) and the emerging livestock-associated (LA)-MRSA clade (*scn* negative) [21], even though LA-MRSA CC398, in rare cases, can readapt to the human host through the regaining of an IEC-harbouring ϕ Sa3 [63,64]. After a phylogenetic analysis of our CC398 strains, the *scn*-positive stork strains clustered in the human-adapted MSSA lineages, whereas the pig and pig farmer strains clustered in the livestock clade as expected by the absence of *scn* (Fig. 2). Besides prophage ϕ Sa3, integrases of other prophages including ϕ Sa2, ϕ Sa6 and ϕ Sa9 were also frequently identified in *S. aureus* strains of the nestling storks, pigs and pig farmers regardless of the methicillin susceptibility and genetic lineages, whereas only MSSA-CC398 strains harboured prophages carrying integrase genes *SebogInt* and *Sa9Int* (Supplementary Tables S3 and S4).

4. Conclusion

Using phylogenetic analysis and the comparison of patterns in host specificity and AMR markers, this study confirms spillover patterns of *S. aureus* between pigs and pig farmers, and from human-associated reservoirs to nestling storks. Furthermore, AMR genes from the pigs and pig farmers were highly associated with mobile genetic elements compared to those from the nestling storks, suggesting increased mobility of AMR genes in pig farms.

A larger sample size could provide more credence to data obtained from the statistical analysis, as some of the observed values were small. However, the idea behind the analyses was to present the relationship of the test variables with the origin or host of the *S. aureus* strains. Collectively, our findings underscore the need to strengthen molecular surveillance of *S. aureus* using the One Health model to direct appropriate control strategies globally.

Funding: This work was supported by the project PID2019-106158RB-I00 of the MCIN/AEI/10.13039/501100011033 of Spain and project SBPLY/19/180501/000325 of the regional government of Castilla – La Mancha co-financed by the European Union's funds

for regional development (Feder). Also, it received funding from the European Union's H2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement no. 801586.

Conflict of interest: None declared by the authors.

Ethical Approval: All the study protocols have been reviewed and approved by the ethical research committees of the University of Zaragoza, the University of La Rioja and the University of Castilla La Mancha (Spain).

Supplementary materials

Supplementary material associated with this article can be found, in the online version, at doi:10.1016/j.jgar.2023.12.003.

References

- [1] Abdullahi IN, Fernández-Fernández R, Juárez-Fernández G, Martínez-Álvarez S, Eguizabal P, Zarazaga M, et al. Wild animals are reservoirs and sentinels of *Staphylococcus aureus* and MRSA clones: a problem with 'One Health' concern. *Antibiotics* (Basel) 2021;10:1556. doi:10.3390/antibiotics10121556.
- [2] Manyi-Loh C, Mamphweli S, Meyer E, Okoh A. Antibiotic use in agriculture and its consequential resistance in environmental sources: potential public health implications. *Molecules* 2018;23:795. doi:10.3390/molecules23040795.
- [3] Silva V, Lopes AF, Soeiro V, Caniça M, Manageiro V, Pereira JE, et al. Nocturnal birds of prey as carriers of *Staphylococcus aureus* and other staphylococci: diversity, antimicrobial resistance and clonal lineages. *Antibiotics* (Basel) 2022;11:240. doi:10.3390/antibiotics11020240.
- [4] Mama OM, Ruiz-Ripa L, Fernández-Fernández R, González-Barrio D, Ruiz-Fons JF, Torres C. High frequency of coagulase-positive staphylococci carriage in healthy wild boar with detection of MRSA of lineage ST398-t011. *FEMS Microbiol Lett* 2019;366:fny292. doi:10.1093/femsle/fny292.
- [5] Gómez P, Lozano C, Camacho MC, Lima-Barbero JF, Hernández JM, Zarazaga M, Höfle U, Torres C. Detection of MRSA ST3061-t843-mecC and ST398-t011-mecA in white stork nestlings exposed to human residues. *J Antimicrob Chemother* 2016;71:53–7.
- [6] Trósciańczyk A, Nowakiewicz A, Kasela M, Malm A, Tracz AM, Hahaj-Siembida A, et al. Multi-host pathogen *Staphylococcus aureus*: epidemiology, drug resistance and occurrence in humans and animals in Poland. *Antibiotics* (Basel) 2023;12:1137. doi:10.3390/antibiotics12071137.
- [7] Martínez-Seijas C, Mascarós P, Lizana V, Martí-Marco A, Arnau-Bonachera A, Chillida-Martínez E, et al. Genomic characterization of *Staphylococcus aureus* in wildlife. *Animals* (Basel) 2023;13:1064. doi:10.3390/ani13061064.
- [8] Heaton CJ, Gerbig GR, Sensius LD, Patel V, Smith TC. *Staphylococcus aureus* epidemiology in wildlife: a systematic review. *Antibiotics* (Basel) 2020;9:89. doi:10.3390/antibiotics9020089.
- [9] Zarazaga M, Gómez P, Ceballos S, Torres C. Molecular epidemiology of *Staphylococcus aureus* lineages in the animal-human interface (Chapter 10). *Staphylococcus aureus*. Fetsch A, editor. Academic Press; 2020. doi:10.1016/B978-0-12-809671-0.00010-3.
- [10] Haag AF, Fitzgerald JR, Penadés JR. *Staphylococcus aureus* in animals. *Microbiol Spectr* 2019;7. doi:10.1128/microbiolspec.GPP3-0060-2019.
- [11] Howden BP, Giulieri SG, Wong Fok Lung T, Baines SL, Sharkey LK, Lee JYH, et al. *Staphylococcus aureus* host interactions and adaptation. *Nat Rev Microbiol* 2023;21:380–95. doi:10.1038/s41579-023-00852-y.
- [12] Campos B, Pickering AC, Rocha LS, Aguilar AP, Fabres-Klein MH, de Oliveira Mendes TA, et al. Diversity and pathogenesis of *Staphylococcus aureus* from bovine mastitis: current understanding and future perspectives. *BMC Vet Res* 2022;18:115. doi:10.1186/s12917-022-03197-5.
- [13] Costa SS, Ribeiro R, Serrano M, Oliveira K, Ferreira C, Leal M, Pomba C, Couto I. *Staphylococcus aureus* causing skin and soft tissue infections in companion animals: antimicrobial resistance profiles and clonal lineages. *Antibiotics* (Basel) 2022;11:599. doi:10.3390/antibiotics11050599.
- [14] Richardson EJ, Bacigalupe R, Harrison EM, Weinert LA, Lycett S, Vrieling M, et al. Gene exchange drives the ecological success of a multi-host bacterial pathogen. *Nat Ecol Evol* 2018;2:1468–78. doi:10.1038/s41559-018-0617-0.
- [15] Gelbířová T, Brodříková K, Karpíšková R. Livestock-associated methicillin-resistant *Staphylococcus aureus* in Czech retail ready-to-eat meat products. *Int J Food Microbiol* 2022;374:109727. doi:10.1016/j.jfoodmicro.2022.109727.
- [16] Golob M, Pate M, Kušar D, Zajc U, Papić B, Očepnik M, Zdobc I, Avberšek J. Antimicrobial resistance and molecular characterization of methicillin-resistant *Staphylococcus aureus* from two pig farms: longitudinal study of LA-MRSA. *Antibiotics* (Basel) 2022;11:1532. doi:10.3390/antibiotics11111532.
- [17] Chaguza C, Smith JT, Bruce SA, Gibson R, Martin IW, Andam CP. Prophage-encoded immune evasion factors are critical for *Staphylococcus aureus* host infection, switching, and adaptation. *Cell Genom* 2022;2:100194. doi:10.1016/j.xgen.2022.100194.

- [18] Avberšek J, Golob M, Papić B, Dermota U, Grmek Košnik I, Kušar D, et al. Live-stock-associated methicillin-resistant *Staphylococcus aureus*: Establishing links between animals and humans on livestock holdings. *Transbound Emerg Dis* 2021;68:789–801.
- [19] Anker JCH, Koch A, Ethelberg S, Mølbak K, Larsen J, Jepsen MR. Distance to pig farms as risk factor for community-onset livestock-associated MRSA CC398 infection in persons without known contact to pig farms: a nationwide study. *Zoonoses Public Health* 2018;65:352–60. doi:10.1111/zph.12441.
- [20] EFSA. Analysis of the baseline survey on the prevalence of methicillin-resistant *Staphylococcus aureus* (MRSA) in holdings with breeding pigs, in the EU, 2008 – Part A: MRSA prevalence estimates. <https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2009.1376> (accessed April 20, 2023).
- [21] Price LB, Stegger M, Hasman H, Aziz M, Larsen J, Andersen PS, et al. *Staphylococcus aureus* CC398: host adaptation and emergence of methicillin resistance in livestock. *mBio* 2012;3:e00305–11 Erratum in: *mBio*. 2013;4. doi:10.1128/mBio.00520-12. doi:10.1128/mBio.00305-11.
- [22] Sieber RN, Larsen AR, Urth TR, Iversen S, Møller CH, Skov RL, Larsen J, Stegger M. Genome investigations show host adaptation and transmission of LA-MRSA CC398 from pigs into Danish healthcare institutions. *Sci Rep* 2019;9:18655. doi:10.1038/s41598-019-55086-x.
- [23] Bouillier K, Zeggay A, Gbaguidi-Naore H, Hocquet D, Chirouze C, Bertrand X. Epidemiology and risk factors of nasal carriage of *Staphylococcus aureus* CC398 in two distinct cohorts in France. *Front Microbiol* 2022;13:1068420. doi:10.3389/fmicb.2022.1068420.
- [24] Bouillier K, Hocquet D, Sauget M, Bertrand X, Chirouze C. Epidemiology and risk factors of *Staphylococcus aureus* CC398 bone and joint infections. *BMC Infect Dis* 2020;20:384. doi:10.1186/s12879-020-05098-0.
- [25] Mama OM, Aspiroz C, Ruiz-Ripa L, Ceballos S, Iñiguez-Barrio M, Cercenado E, et al. Study Group of Clinical *S. aureus* CC398. Prevalence and genetic characteristics of *Staphylococcus aureus* CC398 strains from invasive infections in Spanish hospitals, focusing on the livestock-independent CC398-MSSA clade. *Front Microbiol* 2021;12:623108. doi:10.3389/fmicb.2021.623108.
- [26] He W, Chen H, Zhao C, Zhang F, Li H, Wang Q, Wang X, Wang H. Population structure and characterisation of *Staphylococcus aureus* from bacteraemia at multiple hospitals in China: association between antimicrobial resistance, toxin genes and genotypes. *Int J Antimicrob Agents* 2013;42:211–19. doi:10.1016/j.ijantimicag.2013.04.031.
- [27] Gómez P, Ruiz-Ripa L, Fernández-Fernández R, Gharsa H, Ben Slama K, Höfle U, et al. Genomic analysis of *Staphylococcus aureus* of the lineage CC130, including *mecC*-carrying MRSA and MSSA strains recovered of animal, human, and environmental origins. *Front Microbiol* 2021;12:655994. doi:10.3389/fmicb.2021.655994.
- [28] Randad PR, Larsen J, Kaya H, Pisanic N, Ordak C, Price LB, et al. Transmission of antimicrobial-resistant *Staphylococcus aureus* clonal complex 9 between pigs and humans, United States. *Emerg Infect Dis* 2021;27:740–8.
- [29] Abdullahi IN, Lozano C, Simon C, Latorre-Fernandez J, Zarazaga M, Torres CNasal staphylococci community of healthy pigs and pig-farmers in Aragon (Spain). Predominance and within-host resistome diversity in MRSA-CC398 and MSSA-CC9 lineages. *One Health* 2023;16:100505. doi:10.1016/j.onehlt.2023.100505.
- [30] Abdullahi IN, Juárez-Fernández G, Höfle U, Latorre-Fernández J, Cardona-Cabrera T, Mínguez-Romero D, et al. *Staphylococcus aureus* carriage in the nasotracheal cavities of white stork nestlings (*Ciconia ciconia*) in Spain: genetic diversity, resistomes and virulence factors. *Microb Ecol* 2023;86:1993–2002.
- [31] Abdullahi IN, Juárez-Fernández G, Höfle U, Cardona-Cabrera T, Mínguez D, Pineda-Pampliega J, et al. Nasotracheal microbiota of nestlings of parent white storks with different foraging habits in Spain. *Ecohealth* 2023;20:105–121.
- [32] Magiorakos AP, Srinivasan A, Carey RB, Carmeli Y, Falagas ME, Giske CG, et al. Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired resistance. *Clin Microbiol Infect* 2012;18:268–81.
- [33] Abdullahi IN, Lozano C, Zarazaga M, Saidenberg ABS, Stegger M, Torres C. Clonal relatedness of coagulase-positive staphylococci among healthy dogs and dog-owners in Spain: detection of multidrug-resistant-MSSA-CC398 and novel linezolid-resistant-MRSA-CC5. *Front Microbiol* 2023;14:1121564. doi:10.3389/fmicb.2023.1121564.
- [34] Sahli JW, Lemmer D, Travis J, Schupp JM, Gillece JD, Aziz M, et al. NASP: an accurate, rapid method for the identification of SNPs in WGS datasets that supports flexible input and output formats. *Microb Genom* 2016;2:e000074. doi:10.1099/mgen.0.000074.
- [35] Letunic I, Bork P. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree display and annotation. *Nucleic Acids Res* 2021;49:W293–6.
- [36] Jolley KA, Bray JE, Maiden MCJ. Open-access bacterial population genomics: BIGSdb software, the PubMLST.org website and their applications. *Wellcome Open Res* 2018;3:124. doi:10.12688/wellcomeopenres.14826.1.
- [37] Bortolaia V, Kaas RS, Ruppe E, Roberts MC, Schwarz S, Cattoir V, et al. ResFinder 4.0 for predictions of phenotypes from genotypes. *J Antimicrob Chemother* 2020;75:3491–500.
- [38] Zankari E, Allesøe R, Joensen KG, Cavaco LM, Lund O, Aarestrup FM. PointFinder: a novel web tool for WGS-based detection of antimicrobial resistance associated with chromosomal point mutations in bacterial pathogens. *J Antimicrob Chemother* 2017;72:2764–8. doi:10.1093/jac/dkx217.
- [39] Pal C, Bengtsson-Palme J, Rensing C, Kristiansson E, Larsson DG. BacMet: antibacterial biocide and metal resistance genes database. *Nucleic Acids Res* 2014;42:D737–43. doi:10.1093/nar/gkt1252.
- [40] Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, Wishart DS. PHASTER: a better, faster version of the PHAST phage search tool. *Nucleic Acids Res* 2016;44:W16–21.
- [41] Clausen PTLC, Aarestrup FM, Lund O. Rapid and precise alignment of raw reads against redundant databases with KMA. *BMC Bioinformatics* 2018;19:307. doi:10.1186/s12859-018-2336-6.
- [42] Bartels MD, Petersen A, Worning P, Nielsen JB, Larner-Svensson H, Johansen HK, et al. Comparing whole-genome sequencing with Sanger sequencing for spa typing of methicillin-resistant *Staphylococcus aureus*. *J Clin Microbiol* 2014;52:4305–8.
- [43] Chang SC, Lin LC, Lu JJ. Comparative genomic analyses reveal potential factors responsible for the ST6 oxacillin-resistant *Staphylococcus lugdunensis* endemic in a hospital. *Front Microbiol* 2021;12:765437. doi:10.3389/fmicb.2021.765437.
- [44] Chang SC, Lin LC, Ge MC, Liu TP, Lu JJ. Characterization of a novel, type II staphylococcal cassette chromosome *mec* element from an endemic oxacillin-resistant *Staphylococcus lugdunensis* clone in a hospital setting. *J Antimicrob Chemother* 2019;74:2162–5. doi:10.1093/jac/dkz189.
- [45] Soge OO, Beck NK, White TM, No DB, Roberts MC. A novel transposon, Tn6009, composed of a Tn916 element linked with a *Staphylococcus aureus* mer operon. *J Antimicrob Chemother* 2008;62:674–80. doi:10.1093/jac/dkn255.
- [46] Founou RC, Founou LL, Allam M, Ismail A, Essack SY. *Enterococcus faecalis* ST21 harbouring Tn6009 strain from a carriage sample in South Africa. *S Afr Med J* 2021;111:98–9. doi:10.7196/SAMJ.2021.v111i2.15454.
- [47] Feßler A, Kadlec K, Wang Y, Zhang WJ, Wu C, Shen J, Schwarz S. Small antimicrobial resistance plasmids in livestock-associated methicillin-resistant *Staphylococcus aureus* CC398. *Front Microbiol* 2018;9:2063. doi:10.3389/fmicb.2018.02063.
- [48] Kadlec K, Schwarz S. Identification of the novel *dfrK*-carrying transposon Tn559 in a porcine methicillin-susceptible *Staphylococcus aureus* ST398 strain. *Antimicrob Agents Chemother* 2010;54:3475–7. doi:10.1128/AAC.00464-10.
- [49] Gómez-Sanz E, Kadlec K, Feßler AT, Zarazaga M, Torres C, Schwarz S. Novel *erm(T)*-carrying multiresistance plasmids from porcine and human strains of methicillin-resistant *Staphylococcus aureus* ST398 that also harbor cadmium and copper resistance determinants. *Antimicrob Agents Chemother* 2013;57:3275–82. doi:10.1128/AAC.00171-13.
- [50] Trif E, Cerbu C, Olah D, Zăblău SD, Spînu M, Potârniche AV, Pall E, Brudăscă F. Old antibiotics can learn new ways: a systematic review of florfenicol use in veterinary medicine and future perspectives using nanotechnology. *Animals (Basel)* 2023;13:1695. doi:10.3390/ani13101695.
- [51] Krüger-Haker H, Ji X, Bartel A, Feßler AT, Hanke D, Jiang N, et al. Metabolic characteristics of porcine LA-MRSA CC398 and CC9 strains from Germany and China via Biolog Phenotype MicroArray™. *Microorganisms* 2022;10:2116. doi:10.3390/microorganisms10112116.
- [52] Leinweber H, Sieber RN, Bojer MS, Larsen J, Ingmer H. Fluoroquinolone resistance does not facilitate phage Φ13 integration or excision in *Staphylococcus aureus*. *Access Microbiol* 2023;5:acmi000583.v4. doi:10.1099/acmi.0.000583.v4.
- [53] Huang L, Ahmed S, Gu Y, Huang J, An B, Wu C, Zhou Y, Cheng G. The effects of natural products and environmental conditions on antimicrobial resistance. *Molecules* 2021;26:4277. doi:10.3390/molecules26144277.
- [54] Cavaco LM, Hasman H, Aarestrup FM. Zinc resistance of *Staphylococcus aureus* of animal origin is strongly associated with methicillin resistance. *Vet Microbiol* 2011;150:344–8. doi:10.1016/j.vetmic.2011.02.014.
- [55] Monecke S, Roberts MC, Braun SD, Diezel K, Müller E, Reinicke M, et al. Sequence analysis of novel *Staphylococcus aureus* lineages from wild and captive macaques. *Int J Mol Sci* 2022;23:11225. doi:10.3390/ijms231911225.
- [56] Rebelo A, Mourão J, Freitas AR, Duarte B, Silveira E, Sanchez-Valenzuela A, et al. Diversity of metal and antibiotic resistance genes in *Enterococcus* spp. from the last century reflects multiple pollution and genetic exchange among phyla from overlapping ecosystems. *Sci Total Environ* 2021;787:147548. doi:10.1016/j.scitotenv.2021.147548.
- [57] Karamatakis T, Papadopoulos P, Tsergouli K, Angelidis AS, Melidou A, Sergelidis D, Papa A. Genetic characterization of livestock-associated methicillin-resistant *Staphylococcus aureus* strain in Greece. *Braz J Microbiol* 2021;52:2091–6.
- [58] Schijffelen MJ, Boel CH, van Strijp JA, Fluit AC. Whole genome analysis of a livestock-associated methicillin-resistant *Staphylococcus aureus* ST398 strain from a case of human endocarditis. *BMC Genomics* 2010;11:376. doi:10.1186/1471-2164-11-376.
- [59] Seier-Petersen MA, Nielsen LN, Ingmer H, Aarestrup FM, Agersø Y. Biocide susceptibility of *Staphylococcus aureus* CC398 and CC30 strains from pigs and identification of the biocide resistance genes, *qacG* and *qacC*. *Microb Drug Resist* 2015;21:527–36.
- [60] Grispoli L, Karama M, Armani A, Hadjicharalambous C, Cenci-Goga BT. *Staphylococcus aureus* enterotoxin in food of animal origin and staphylococcal food poisoning risk assessment from farm to table. *Italian J Ani Sci* 2021;20:1677–90.
- [61] Darisipudi MN, Bröker BM. How *S. aureus* blinds the inflammasome to escape immune control. *EBioMedicine* 2021;71:103549. doi:10.1016/j.ebiom.2021.103549.

- [62] Idrees M, Sawant S, Karodia N, Rahman A. *Staphylococcus aureus* biofilm: morphology, genetics, pathogenesis and treatment strategies. *Int J Environ Res Public Health* 2021;18:7602. doi:[10.3390/ijerph18147602](https://doi.org/10.3390/ijerph18147602).
- [63] Sieber RN, Urth TR, Petersen A, Møller CH, Price LB, Skov RL, et al. Phage-mediated immune evasion and transmission of livestock-associated methicillin-resistant *Staphylococcus aureus* in humans. *Emerg Infect Dis* 2020;26:2578–85. doi:[10.3201/eid2611.201442](https://doi.org/10.3201/eid2611.201442).
- [64] Pérez-Moreno MO, Centelles-Serrano MJ, Nogales-López J, Domenech-Spanedda MF, Lozano C, Torres C. Unusual presence of the immune evasion gene cluster in livestock-associated MRSA of lineage CC398 causing peridural and psoas abscesses in a poultry farmer. *Enferm Infecc Microbiol Clin* 2017;35:651–4.