

Earth's Future

RESEARCH ARTICLE

10.1029/2023EF003629

Special Section:

CMIP6: Trends, Interactions,
Evaluation, and Impacts

Key Points:

- Important differences in drought projections as a function of drought metrics
- The temporal relationship between the precipitation-based climatic metrics is high worldwide
- A weak relationship is found between climatic and ecological drought indices

Supporting Information:

Supporting Information may be found in the online version of this article.

Correspondence to:

L. Gimeno-Sotelo,
luis.gimeno-sotelo@uvigo.es

Citation:

Gimeno-Sotelo, L., El Kenawy, A., Franquesa, M., Noguera, I., Fernández-Duque, B., Domínguez-Castro, F., et al. (2024). Assessment of the global relationship of different types of droughts in model simulations under high anthropogenic emissions. *Earth's Future*, 12, e2023EF003629. <https://doi.org/10.1029/2023EF003629>

Received 25 FEB 2023

Accepted 19 JAN 2024

Author Contributions:

Conceptualization: Luis Gimeno-Sotelo, Sergio M. Vicente-Serrano

Formal analysis: Luis Gimeno-Sotelo, Fergus Reig

Funding acquisition: Sergio M. Vicente-Serrano







Investigation: Luis Gimeno-Sotelo, Sergio M. Vicente-Serrano

Methodology: Luis Gimeno-Sotelo, Sergio M. Vicente-Serrano

Project administration: Sergio M. Vicente-Serrano

© 2024 The Authors. Earth's Future published by Wiley Periodicals LLC on behalf of American Geophysical Union. This is an open access article under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Assessment of the Global Relationship of Different Types of Droughts in Model Simulations Under High Anthropogenic Emissions

Luis Gimeno-Sotelo¹ , Ahmed El Kenawy², Magí Franquesa² , Iván Noguera², Beatriz Fernández-Duque², Fernando Domínguez-Castro², Dhais Peña-Angulo³ , Fergus Reig², Rogert Sorí¹ , Luis Gimeno¹, Raquel Nieto¹ , and Sergio M. Vicente-Serrano² 

¹Centro de Investigación Mariña, Universidade de Vigo, Environmental Physics Laboratory (EPhysLab), Ourense, Spain,

²Instituto Pirenaico de Ecología, Consejo Superior de Investigaciones Científicas (IPE-CSIC), Zaragoza, Spain,

³Department of Geography, University of Zaragoza, Zaragoza, Spain

Abstract This study provides a global analysis of the relationship between drought metrics obtained from several climatic, hydrologic and ecological variables in a climate change framework using CMIP6 model data. A comprehensive analysis of the evolution of drought severity on a global scale is carried out for the historical experiment (1850–2014) and for future simulations under a high emissions scenario (SSP5-8.5). This study focuses on comparing trends in the magnitude and duration of drought events according to different standardized indices over the world land-surface area. The spatial and temporal relationship between the different drought indices on a global scale was also evaluated. Overall, there is a fairly large consensus among models and drought metrics in pointing to drought increase in southern North America, Central America, the Amazon region, the Mediterranean, southern Africa and southern Australia. Our results show important spatial differences in drought projections, which are highly dependent on the drought metric employed. While a strong relationship between climatic indices was evident, climatic and ecological drought metrics showed less dependency over both space and time. Importantly, our study demonstrates uncertainties in future projections of drought trends and their interannual variability related to the relationship among indices, stressing the importance of coherent climatic, hydrological and plant physiological patterns when analyzing CMIP6 model simulations of droughts under a warming climate scenario.

Plain Language Summary Using climate change models, we perform a drought analysis in terms of climatic, hydrologic and ecological variables on a global scale, studying the projections under a high emission scenario. We analyze how drought events will evolve in the future with respect to their magnitude and duration, and if the different drought metrics agree in space and time. In general, models and metrics agree that there will be drought increase in southern North America, Central America, the Amazon region, the Mediterranean, southern Africa and southern Australia. However, results differ across the world and really depend on the metric used. We show that climatic indices are strongly connected with each other, but no so related to ecological ones. We also find that there are uncertainties in future projections of drought trends, highlighting that we should always take into account the spatial and temporal agreement between climatic, hydrological and plant physiological patterns when studying drought projections.

1. Introduction

Assessment of future drought projections is at the forefront of scientific debate in the current research on climate, hydrology, agriculture, and ecology. This is simply due to the multiple dimensions of droughts, which cause strong complexity for drought assessment and quantification (Douville et al., 2021; Lloyd-Hughes, 2014). In addition, the strong environmental and socioeconomic implications of drought changes in future climate scenarios adds more complexity to this debate (Naumann et al., 2021; Van Loon et al., 2016; Xu et al., 2019).

In order to robustly assess future changes in drought severity, we must refer to different types of drought. This is fundamental to properly evaluate the impacts associated with drought in future climates. Generally, the concepts of meteorological drought (precipitation deficits), agricultural droughts (crop failure or yield decrease), ecological droughts (damages in natural vegetation, reduced photosynthesis activity, and carbon uptake and increased plant mortality), and hydrological droughts (reductions in the availability of water in different sources

Resources: Sergio M. Vicente-Serrano

Supervision: Luis Gimeno, Sergio M. Vicente-Serrano

Visualization: Luis Gimeno-Sotelo

Writing – original draft: Luis Gimeno-Sotelo, Ahmed El Kenawy, Magí Franesca, Iván Noguera, Beatriz Fernández-Duque,

Fernando Domínguez-Castro, Dhais Peña-Angulo, Rogert Sorí, Luis Gimeno, Raquel Nieto, Sergio M. Vicente-Serrano

Writing – review & editing:

Luis Gimeno-Sotelo, Ahmed El Kenawy, Magí Franesca, Iván Noguera, Beatriz Fernández-Duque, Fernando Domínguez-Castro, Dhais Peña-Angulo, Rogert Sorí, Luis Gimeno, Raquel Nieto, Sergio M. Vicente-Serrano

such as reservoir storages, streamflow and groundwater) are used commonly to refer to drought types. These types are largely impacted by different physical and ecological processes (Douville et al., 2021; Lobell, 2014; Vicente-Serrano, Quiring, et al., 2020; Wilhite & Buchanan-Smith, 2005).

In the literature, a wide spectrum of studies characterized drought projections on the global scale using model simulations of various climatic, hydrological, and vegetation variables under different future climates scenarios (e.g., Cook et al., 2014, 2020; Lu et al., 2019; Martin, 2018; Papalexiou et al., 2021; Ridder et al., 2022; Ukkola et al., 2020; Vicente-Serrano, Domínguez-Castro, et al., 2020; Zeng et al., 2022; Zhao & Dai, 2021; Zhu & Yang, 2021). Nonetheless, most of these studies focused on metrics directly simulated by different Coupled Model Intercomparison Projects (CMIP) since they allow to directly evaluate drought impacts on a variety of agricultural, ecological, and hydrological systems (Bachmair et al., 2016, 2018; Hlavinka et al., 2009; O'Connor et al., 2022; Quiring & Papakryiakou, 2003; Stagge, Kohn, et al., 2015; Vicente-Serrano et al., 2012).

In the literature, the most widely used drought metrics for drought monitoring and impact assessment are synthetic indices that combine precipitation and atmospheric evaporative demand (AED), allowing for a direct quantification of drought severity and drought extent (Dai, 2021; Tomas-Burguera et al., 2020; van der Schrier et al., 2013; Vicente-Serrano et al., 2010), as well as their impacts on ecosystems (Bachmair et al., 2015). For future simulations, different studies analyzed drought projections based on these indices, employing Earth System Models (ESMs) outputs under different future climate scenarios (Dai, 2012; Naumann et al., 2018; Spinoni et al., 2020; Vicente-Serrano, Domínguez-Castro, et al., 2020; Zhao & Dai, 2022). According to these scenarios, drought severity would increase, mainly as a consequence of the enhanced AED in a warming climate. Nonetheless, some studies suggest uncertainty of using these metrics (e.g., Berg & Sheffield, 2018; McColl et al., 2022). Specifically, the criticisms argue that these indices are not necessarily representative of the metrics based on water storage (i.e., soil moisture), surface water generation (i.e., runoff) or vegetation activity (i.e., leaf area and net primary production (NPP)). These arguments would be supported by the notion that hydrological and ecological systems might show different dynamics and responses under future climates (Berg & Sheffield, 2018; Scheff, 2018). Furthermore, CMIP models generate simulations of hydrological and plant metrics, which would make it unnecessary to focus on climate metrics as proxies of drought impacts (McColl et al., 2022). Moreover, drought indices that include AED in their calculations might overestimate drought severity under high-emissions future climate scenarios. This is simply because future increase in AED is likely to be higher than the expected increase in land evapotranspiration (Et) (Milly & Dunne, 2016; Roderick et al., 2015; Scheff, 2018; Yang et al., 2019), which is also determined by water availability.

As such, a more complete spatio-temporal comparison of different drought metrics is necessary to provide a more robust picture of how drought responds to future climate and better assessment of the advantages and limitations of using drought indices based on simulations of climate variables versus the use of indices based on simulated hydrological and ecological variables. Although recent studies have analyzed global drought projections based on the latest model outputs from the CMIP6 using different drought metrics (e.g., Cook et al., 2020; Li et al., 2021; Meng et al., 2022; Papalexiou et al., 2021; Ukkola et al., 2020; Zeng et al., 2022; Zhao & Dai, 2022; Zhu and Yang, 2021), these studies lacked the opportunity to investigate some drought metrics that are important for assessing agricultural and ecological droughts. As such, a focus on these gaps may provide new evidence that provides more certainties about the use of different drought metrics to assess future trends in drought severity. On the other hand, it is necessary to test the spatial and temporal relationship among the different drought metrics, which can give indications on the reliability of drought projections. In the pursuit of this background, the objectives of this study are to:

1. know the differences in the future drought projections based on a wide set of climatic, ecological and hydrological drought metrics, providing an assessment of the regional and global coherence and uncertainty in the projections.
2. determine the spatial and temporal relationship among the different drought metrics in replicating drought severity based on the comparison of the spatial and temporal agreement of drought conditions at the annual scale.

Accordingly, the current global assessment can contribute to the arising debate on the robustness of the different drought metrics, and the existing uncertainties for agricultural, ecological, and hydrological drought projections under a high-emission climate scenario.

2. Data and Methods

We employed monthly data of a set of hydroclimatic variables from the CMIP6 experiment (Eyring et al., 2016). These variables included precipitation, runoff, total column soil moisture, leaf area index (LAI) and NPP. Data were provided for the historical period (1850–2014) and for the Shared Socioeconomic Pathway (SSP; 5–8.5) from 2015 to 2100. All CMIP6 individuals that secure data for the necessary variables, as well as the period 1850–2100, were considered in our analysis (see Table S1 in Supporting Information S1). Recalling that the CMIP6 outputs are provided in different native spatial resolutions, we interpolated data to a common resolution of $2.5^{\circ} \times 2.5^{\circ}$ by means of a bilinear transformation. We used this low spatial resolution in order to be conservative to avoid that the models are resampled to a higher spatial resolution than the original simulations. In general, bilinear transformation provides excellent results to convert CMIP models to a comparable spatial resolution, and it has been used for the data analysis in the IPCC reports (Gutiérrez et al., 2021; Stocker et al., 2013), being a usual procedure in climate studies (Almazroui et al., 2020; Rettie et al., 2023; Thorarinsdottir et al., 2020; Wu et al., 2022). To assess future projections in drought severity, our decision was made to consider the SSP5-8.5 scenario, which represents the worst possible scenario compared to the historical experiment, thereby accentuating the most pronounced differences among various climatic, hydrological, and ecological metrics.

The standardized drought indices were computed based on the common data inputs (e.g., precipitation, runoff, total column soil moisture, LAI and NPP). Nonetheless, other indices were computed using a combination of new variables. For example, maximum and minimum air temperatures, relative humidity, wind speed and solar radiation, were used to calculate AED following the Penman-Monteith FAO-56 equation (Pereira et al., 2015). Overall, based on these data and data of Evapotranspiration (Et), we calculated different indices using: (a) the difference between precipitation and AED (P-AED), which is a metric that has been widely used for drought assessment since it summarizes the balance between the water available in the form of precipitation and the existing AED (Tomas-Burguera et al., 2020; Vicente-Serrano et al., 2010), (b) precipitation minus land evapotranspiration (P-Et), which is considered a long-term water budget and has been accordingly used to assess drought severity in several works (e.g., Padrón et al., 2020), and (c) the difference between Et and AED (Et-AED), which compares the difference between the available water to evaporate and the water demand by the atmosphere (Kim & Rhee, 2016; Vicente-Serrano et al., 2018) and is highly related to plant water stress (Stephenson, 1990). All these drought metrics were transformed into the same standardized units to make robust spatial and temporal comparisons. To fit data distribution, a log-logistic distribution was used, which is capable of standardizing different climate and hydrological records under different climate conditions, as being evidenced in earlier works (e.g., Vicente-Serrano & Beguería, 2016; Vicente-Serrano, Domínguez-Castro, et al., 2020). The only exception was for precipitation, which was fitted to a Gamma distribution (Stagge, Tallaksen, et al., 2015). We tested the goodness of fit of the standardized indices using the Kolmogorov-Smirnov test (Kolmogorov, 1933; Smirnov, 1948), which was used to assess if they followed a normal distribution. It compares the cumulative distribution function of a data set to the theoretical normal distribution, quantifying the maximum difference (the KS statistic). The p-value represents the probability of obtaining a KS statistic as extreme as the one observed, assuming that the data follows a normal distribution (null hypothesis). A smaller p-value reveals stronger evidence against the normality assumption, while a larger p-value indicates weaker evidence. The obtained results suggest that, for the majority of world regions, the multimodel median p-value is above 0.01 and therefore we can assume that the indices that were constructed fit well to a normal distribution (Figure S1 in Supporting Information S1). Afterward, a second standardization procedure was carried out independently for each of the 12 monthly series of the indices. To make this standardization, both the mean and the standard deviation were computed for the reference period 1850–2014. This procedure minimizes the possible impacts of strong trends presented in the analyzed variables for future scenarios in the possibility of calculating the drought indices (Vicente-Serrano, Domínguez-Castro, et al., 2020). Furthermore, this standardization allows for a robust spatial and temporal comparability between the different metrics. Accordingly, drought duration and magnitude can be quantified for each time series and for the different indices. Drought events were identified using the run theory (Fleig et al., 2006; Tallaksen et al., 1997), considering a threshold of $z = -1.28$, which corresponds to a 10% probability of a standard normal observation being below that value. For drought event identification, all indices were computed at the 3-month time scale. To analyze the trends in the duration and magnitude of drought events, we used the annual values summarizing the drought conditions recorded during the different months of the year. A linear regression model was fitted as a function of time, and the estimated slope was used to quantify the amount




Figure 1. Workflow chart that summarizes the methodology followed in this study.

of change over time. The significance of these changes was assessed using the Mann–Kendall test (Kendall, 1948; Mann, 1945), at a significance level of $\alpha = 0.05$.

We also analyzed the relationship between the annual indices (computed at 12-month time scale) using partial correlations, which enabled us to understand the unique association between each pair of variables, while controlling for the effects of the other variables (Kim, 2015). They were computed using the Kendall's rank correlation coefficient, that is, Kendall's τ coefficient (Kendall, 1938). This coefficient is a nonparametric measure of rank correlation that is more suitable than parametric statistics (e.g., Pearson's linear correlation coefficient) because it accounts for the non-linear relationships between variables.

For each grid point, the temporal agreement between the indices (computed at 12-month scale) was assessed by obtaining the percentage of simultaneous occurrence of years in which a pair of indices were below $z = -1.28$, thus producing a 2-dimensional representation of the results. Also, we computed the percentage of grid points where each pair of indices showed z -value below -1.28 , resulting in a time series.

A workflow chart that summarizes the methodology followed in this study can be found in Figure 1.




Figure 2. Evolution of the annual average percentage of global land area affected by extreme dry conditions (5%) from 1850 to 2100. Gray lines represent the value for the different independent models and red lines refer to the median.

3. Results

3.1. Evolution of Drought Severity Based on Different Metrics

Figure 2 shows the evolution of the world land surface affected by drought between 1850 and 2100. It is computed as the percentage of land grid points below the fifth percentile of each raw (non-standardized) variable. This percentile is computed independently for each month, considering the 1850–2014 reference period. For all the variables, we found an increase in the world land surface impacted by dry conditions from 1850 to 2100, albeit with some considerable spatial differences. Results demonstrate that precipitation, leaf area, and runoff would likely show a small increase of drought severity in the future. For precipitation-Et and NPP, the increase is mostly intermediate, although a sharp increase in NPP is noted between 2010 and 2030, followed by a constant behavior to the end of the twenty-first century. The possible mechanisms of this sharp increase are not well determined but probably this would be caused by discontinuities in the radiative forcing of the historical and the SSP 5–8.5 scenarios. It can be a feature of certain problem for the temporal comparability of the model outputs considering periods covered by different forcing experiments. For precipitation-AED, Et-AED and soil moisture, a remarkable increase is noted at the end of the twenty-first century. As illustrated in Figures S2 and S3 in Supporting Information S1, some variables exhibited important seasonal and regional differences. For example, during the boreal winter season, drought based on NPP, soil moisture, and Et-AED increase. Rather, for precipitation and runoff, irrelevant drought increase is noted from 1850 to 2100. On the contrary, in the boreal summer season, the main drought increase is recorded for precipitation-AED, Et-AED, and soil moisture, with little increase for other variables (e.g., precipitation, runoff, and precipitation-Et).

Overall, we noted an increase in the magnitude of drought events that affects large areas of the world in terms of precipitation-AED, Et-AED, and soil moisture, albeit with significant spatial differences (Figure 3). Interestingly, these three drought metrics showed a high agreement in terms of the areas that are likely to exhibit the highest increase in the magnitude of drought periods, including the Mediterranean region, Central America, northern South America and western South America, West Africa and South Africa. Nevertheless, it can be noted that the areas affected are much larger using Et-AED metric, with almost the entire land showing an increase in drought severity. Meteorological droughts, based on precipitation, show an increase in drought magnitude in areas of Central and South America, West Africa, South Australia and the Mediterranean region, although this increase is not as high as suggested by other drought indices (i.e., Et-AED, and soil moisture). This pattern is almost similar when considering precipitation-Et, although some areas of South America did not show an increase in drought severity, suggesting that—in specific regions—the increase in drought magnitude can be reduced if Et is included in the calculations. Drought magnitude trends based on runoff show smaller changes than considering exclusively precipitation, suggesting that CMIP6 models project a less increase in the magnitude of hydrological droughts than in the magnitude of meteorological (precipitation) droughts in a high emissions scenario. LAI did not show




Figure 3. Spatial distribution of the median trend in the magnitude of drought events between 1850 and 2100 (Factor: 100).

an increase in the magnitude of drought events in large areas of the world, except for parts of East Brazil. Thus, the spatial pattern was sparse on the global scale, with strong regional variability and a dominance of no changes or decrease in the magnitude of drought events in some regions (e.g., South America, Southeast Asia, Central Europe, and North America). Notably, the NPP-based assessment showed a strong reinforcement of drought magnitude in the high latitudes of the Northern Hemisphere. Rather, in some areas of Africa, South America and Southeast Asia, a decrease in the magnitude of the drought episodes, based on the NPP, was noted. Changes in the duration of drought events were almost similar to those of drought magnitude, particularly in terms of spatial patterns and the behavior of the different drought metrics (Figure S4 in Supporting Information S1).

Some drought metrics show high consistency in identifying positive trends in drought magnitude among the different models. Figure 4 shows the percentage of models showing positive and statistically significant trends in drought magnitude between 1850 and 2100. As depicted, almost all models defined the same the regions with strong increase in drought magnitude considering precipitation-AED and Et-AED. This agreement was much lower for soil moisture, even in large regions where the multimodel median values showed an increase in drought magnitude. A representative example is found in southern North America and South Africa, where multimodel medians showed a large increase in drought magnitude, while less than 40% of the models showed a positive and significant trend. In other regions whereas decline in drought magnitude was observed like northern South America or the Mediterranean, the percentage of models showing significant declining trends was roughly 50%, suggesting a strong uncertainty in model projections. Notably, although precipitation, precipitation-Et and runoff showed a drought increase in fewer regions than soil moisture, the consistency of this increase among models seems to be greater. More than 50% of the models suggest a positive and statistically significant increase in drought magnitude in northern South America and Central America, the Mediterranean and southern Africa for




Figure 4. Percentage of models showing positive and statistically significant trends in drought magnitude from 1850 to 2100.




Figure 5. Percentage of models showing negative and statistically significant trends in drought magnitude from 1850 to 2100.

precipitation. A similar pattern was evident for vast areas in North and South America, Central Africa, and Central and South Asia when considering P-Et. This suggests that Et projections suppress the trend toward higher drought magnitudes in Southern Africa in comparison to precipitation-based projections, with only few models showing a positive and statistically significant trend. Interestingly, for runoff almost 50% of the models suggested a significant increase in drought magnitude in large regions of the Northern Hemisphere (e.g., Alaska, Labrador, Scandinavia, West Russia), while they did not witness a relevant increase in drought magnitude based on precipitation and precipitation-Et metrics. In the same context, apart from the high latitudes of the Northern Hemisphere, there were no regions where more than 30% of models showed an increase in drought magnitude for the NPP. Interestingly, results demonstrate that drought magnitude based on LAI will not change anywhere worldwide, with almost no model suggests significant changes.

Like drought magnitude, similar patterns of drought duration changes were observed globally (Figure S5 in Supporting Information S1), with majority of the models suggesting no significant changes in ecological and agricultural droughts across majority of the world regions under scenarios of high greenhouse gas emissions.

The negative trends in drought magnitude (Figure 5) and duration (Figure S6 in Supporting Information S1) indicated few regions and metrics in which the models agree on a decrease in drought severity, mainly for precipitation in the high latitudes of the Northern Hemisphere. Even for LAI and NPP, the percentage of models that showed a decrease in drought magnitude is low. As depicted, although some areas, based on some metrics, showed a projected decrease in drought duration and magnitude with multimodel medians (e.g., Southeast Asia with LAI, Central Africa with the NPP, West Russia with soil moisture), there is still large inconsistency among the models. In the same context, while a steady increase in drought duration and magnitude was projected for some regions and variables, only few areas witnessed a decrease in drought duration and magnitude, irrespective of drought metric used. Thus, although there are important uncertainties between drought metrics and models related to the increase of drought duration and magnitude, there is a high consistency between models and metrics concerning drought decrease since drought magnitude and duration are not expected to decrease under a scenario of high greenhouse gasses emissions.

3.2. Spatio-Temporal Relationships Among Drought Metrics

In addition to knowing the consistency of trends between different drought metrics and models, it is also relevant to analyze the consistency of the temporal relationship in the drought severity based on these metrics (Figure S7 in Supporting Information S1). As illustrated, we found strong annual relationships between some pairs of drought indices in the historical period. For example, the correlation was higher than 0.8 between precipitation and precipitation-AED and between precipitation and precipitation-Et in most areas of the world. Also, a high correlation was observed between precipitation-AED and precipitation-Et, with few exceptions, mainly in arid and semiarid regions where correlations decreased. Other pairs of drought metrics showed lower relationships on global scale, with important spatial differences. For example, the relationship between precipitation and Et-AED was only high in water-limited regions, where Et is mostly determined by water availability. It is worth mentioning that the relationship between precipitation (and also between the other climatic metrics) and soil

moisture was low in most regions. Thus, the correlation with soil moisture was higher considering precipitation-AED and particularly Et-AED in regions like South America, Africa, and South Asia. LAI and NPP showed high correlations particularly in water-limited and cold regions. Nevertheless, these two ecological variables showed low correlations with the different meteorological drought metrics, suggesting that the interannual variability of agricultural and ecological droughts simulated by models is independent from those of climatic droughts in most regions of the world. This pattern was also observed considering soil moisture, with low correlations found between the interannual variability of soil moisture and the NPP and LAI in most regions, irrespective of biome types and bioclimatic conditions. The relationship between precipitation and runoff was high in most regions of the world, except for North America and most of Eurasia. In contrast, the relationship between interannual variability of runoff and soil moisture tended to be low globally, apart from the Mediterranean, northern South America, and Africa. Similarly, ecological metrics (i.e., NPP and LAI) showed low correlations with runoff worldwide.

Overall, these results suggest that, except for the high relationship between different climate metrics and their corresponding spatial differences that are mainly determined by the average water availability and temperature, the temporal relationship between different drought metrics was generally low in most regions of the world. This relationship was particularly low between climatic and vegetation metrics, as well as between soil moisture and other drought metrics.

The spatial pattern and the magnitude of the temporal relationships between the different variables, analyzed by means of partial correlations, did not show important changes considering future simulations (2015–2100), as compared with historical simulations (Figure S8 in Supporting Information S1), albeit with some important exceptions (Figure 6). For example, the relationship between the interannual variability of precipitation and other climatic drought metrics generally decreased, which is quite relevant in some areas of Central Asia considering precipitation-AED, but also in the Sahel and high latitudes of the Northern Hemisphere considering Et-AED. On the contrary, the relationship between precipitation and precipitation-Et remained stable for both the historical period and future. Also, we found a decrease in the relationship between precipitation-AED and precipitation-Et in some regions of Europe, South America, and Africa. The relationship between LAI and NPP was stable for the historical period and future simulations in most regions, albeit with a trend to reinforce in some regions. In addition, the relationship between precipitation and LAI tended to reinforce in the high latitudes of the Northern Hemisphere. This was also observed with the NPP, although a decline in the correlation between precipitation and NPP was observed in the Mediterranean, southern North America and northern South America. While the relationship between NPP and precipitation-AED was low during the historical period, this relationship was projected to decline further in the future, particularly in arid regions, the Amazon basin, and some wet areas of Africa. The decrease in the relationship with the NPP was even more severe when considering Et-AED, with an overall global decline. In addition, the relationship between NPP and soil moisture is likely to decline over large areas (e.g., the Mediterranean, northern South America, southern Africa, and Australia). Finally, the relationship of the runoff to other drought metrics tended to be stable between the historical period and the future high emission scenario, although a decreasing correlation with precipitation was observed in Scandinavia, and particularly with precipitation-AED and Et-AED in most Africa and the Amazon basin.

The temporal agreement in drought conditions among the different metrics is small in most regions during the historical period (Figure S9 in Supporting Information S1), suggesting that the annual drought conditions tend to differ noticeably between metrics. There was some agreement in the identified drought periods between precipitation and precipitation-AED, except in arid lands. A similar pattern was also noted between precipitation and precipitation-Et in wet regions and between precipitation-AED and Et-AED in arid lands. Nevertheless, the agreement in the occurrence of droughts between climatic, ecologic, and hydrologic metrics was small. Herein, it is worth to note that while our analysis is restricted to annual droughts to reduce the role of seasonality and the lags in the response of hydrological, agricultural and ecological drought conditions to meteorological droughts and irrespective of the physical consistency among models, drought periods mostly do not coincide in time among the different metrics. For the projected scenario, the temporal agreement between metrics shows some increase (Figure S10 in Supporting Information S1). This is particularly relevant in some regions, such as the Mediterranean region, southern Africa, the Amazon basin, and Central America when comparing drought episodes recorded with precipitation and precipitation-AED, precipitation-Et, Et-AED and soil moisture and also between precipitation-AED and precipitation-Et and between Et-AED and soil moisture, particularly in water-limited regions. The agreement in the temporal identification of drought conditions also increases when comparing




Figure 6. Differences in the median Kendall's τ partial correlations between the projected (2015–2100) and historical period (1850–2014) for the different models.

the climatic indices and the runoff in some areas, particularly in the Amazon and the humid regions of Africa, suggesting an agreement in annual droughts between some pairs of drought metrics, especially in water-limited or humid regions (Figure 7).

The temporal agreement between annual droughts was low during the historical period between the different metrics, and also with low spatial agreement, suggesting that the global spatial patterns of annual drought severity usually did not agree between drought metrics obtained from CMIP6 simulations (Figure 8). The spatial agreement of drought conditions tends to increase under future climate change, in particular for some metrics (e.g., precipitation-AED and precipitation-Et, precipitation-AED and Et-AED, precipitation-AED and soil moisture). Nevertheless, the spatial agreement between droughts on the annual scale between climatic indices, runoff, and ecological droughts was low in both the historical experiment and the projected scenario, indicating spatial differences in replicating annual droughts among the different drought metrics obtained from ESMs.




Figure 7. Differences in the average percentage of temporal agreement among the different metrics between the projected (2015–2100) and the historical period (1850–2014) for the different models.

4. Discussion

This study analyzed long-term evolution of different drought metrics on a global scale using CMIP6 models from 1850 to 2100. These metrics represent different climatic, hydrologic, and ecological variables. Results were presented for the historical experiment (1850–2014) and future projections (2015–2100) under a high-emission scenario (SSP5-8.5). While numerous studies assessed drought severity for future climate using CMIP6 models (e.g., Cook et al., 2020; Guo et al., 2022; Papalexou et al., 2021; Ukkola et al., 2020; Wang et al., 2021; Zhao & Dai, 2022), our assessment employed a larger number of drought metrics, including climate-based (precipitation, precipitation-AED, precipitation-Et, Et-AED), hydrological-based (soil moisture and runoff) and plant physiology-based metrics (LAI and NPP). An evaluation of this variety of different metrics is essential to assess different drought types (meteorological, agricultural/ecological and hydrological) and to determine their consistency in terms of projected drought severity.




Figure 8. Evolution of the spatial agreement of dry conditions between the different drought metrics.

4.1. Global Drought Projections Based on Different Metrics

Our results, as suggested by most models and drought metrics, suggest that in a high emissions scenario drought would increase in southern North America, Central America, the Amazon region, the Mediterranean, southern Africa, and southern Australia, which agrees with earlier studies (e.g., Cook et al., 2020; Seneviratne et al., 2021; Ukkola et al., 2020; Wang et al., 2021; Zhao & Dai, 2022). These projections must be considered carefully since the models are affected by substantial biases (Adeyeri et al., 2022, 2023), strong differences with the trends in observations during the historical period (Vicente-Serrano, Miralles, et al., 2022b) and limitations to consider key ecohydrological processes (e.g., the interactions between plant root systems, soil moisture and runoff generation with groundwater) (Miguez-Macho & Fan, 2021; Ndehedehe et al., 2023).

In accordance with previous studies (Cook et al., 2020; Scheff et al., 2021), our results showed important differences in drought projections as a function of drought metrics. For example, the use of AED-based drought metrics (e.g., the Standardized Precipitation Evapotranspiration Index (SPEI)) revealed that drought severity is

likely to be enhanced in future, as compared to those metrics based on precipitation, precipitation-Et, and runoff. This finding agrees with some investigations based on CMIP6 (e.g., Zeng et al., 2022), and CMIP5 outputs (e.g., Cook et al., 2014) and also by studies that employed other metrics like the Palmer Drought Severity Index (PDSI) (e.g., Scheff et al., 2021; Yang et al., 2021; Zhao & Dai, 2022). The different magnitude of drought projections as simulated based on hydrological (i.e., runoff) and climatic drought indices (which use AED in the calculations) is behind the suggested overestimation of drought severity based on climatic indices under high emissions climate change scenarios (Berg & McColl, 2021; Berg & Sheffield, 2018; Greve et al., 2019; Scheff, 2018).

While it can be argued that focusing on the metrics directly indicative of impacts in agricultural, ecological and hydrological systems (i.e., soil moisture, runoff, NPP, and LAI) instead of climatic proxies of drought severity can be a more practical approach (McColl et al., 2022), it is necessary to note that models show higher uncertainties in simulating complex hydrological and plant physiology processes than climate variables, irrespective of the existing uncertainties related to the simulation of different meteorological processes and also the possible coupling mechanisms between the land and the atmosphere. For example, the spatial and temporal variability in soil moisture involves several processes, some of them are unknown, while others are difficult to simulate (Lu et al., 2019; van den Hurk et al., 2011). This may explain poor agreement between soil moisture observations and model simulations (Ford & Quiring, 2019; Yuan & Quiring, 2017). Streamflow generation is also very complex and models usually fail to simulate hydrological droughts (Barella-Ortiz & Quintana-Seguí, 2018; Tallaksen & Stahl, 2014). Plant physiology is also a key factor controlling both hydrological, agricultural and ecological droughts, and models show strong limitations and uncertainties in simulating plant physiological processes and water interchanges with soil and atmosphere (Liu et al., 2020). These problems are even more important for future climate projections (Gentine et al., 2019) in which the models show strong uncertainties when simulating plant physiological processes (Padrón et al., 2022), and other processes may introduce other sources of uncertainty (e.g., the role of atmospheric CO₂ concentrations) (De Kauwe et al., 2021; Vicente-Serrano, Miralles, et al., 2022). For example, Li et al. (2023) have showed very difficult to separate the radiative and physiological effect of the CO₂ concentrations on hydrological and ecological simulations. These authors showed that the increase of the water use efficiency by plants associated to CO₂ is limited by the increase of the temperature and the vapor pressure deficit. This mechanisms could enhance evapotranspiration and plant water stress under scenarios characterized by a high warming (as the SSP5-8.5 used here). In fact, these mechanisms seem to be incompatible with the limited increase of drought severity based on runoff and soil moisture projections. For these reasons, although some studies argue that ecological and hydrological drought metrics obtained from model simulations may be more accurate than AED-based climatic indices, we should consider that these metrics are affected by stronger biophysical and hydrological uncertainties. On the contrary, the drought indices based on climate variables, although simpler conceptually, would be less constrained by these problems.

One of the novelties of our study is the use of diverse metrics, which is fundamental to address drought characteristics and impacts. In particular, we employed the Standardized Evapotranspiration Deficit Index (SEDI), based on the difference between Et and AED, which is informative on plant water stress (Alsafadi et al., 2022; Jiang et al., 2022; Kim & Rhee, 2016; Li et al., 2019, 2020; Vicente-Serrano et al., 2018; Zhang et al., 2019) with several biogeographic implications (Stephenson, 1990). Changes in the SEDI, both in spatial patterns and drought severity, were almost similar, or even stronger than those obtained by the SPEI, and are characterized by an increase in drought severity under future scenarios of high anthropogenic emissions. In addition, we used two eco-physiological metrics, LAI and NPP, which have been considered by few studies as metrics of drought severity in model simulations (e.g., Scheff et al., 2021). As opposed to the SEDI, our assessment based on the LAI and NPP did not suggest an increase in agricultural and ecological drought severity, except for the high latitudes of the Northern Hemisphere. This regional pattern may be explained by the role of snow and permafrost melt processes that could affect water availability (Chen et al., 2021).

4.2. Uncertainties in Ecological Drought Projections

The picture provided by the eight drought metrics showed some paradoxical projections that are difficult to explain by plant physiological processes. In particular, different studies focusing on plant physiology have highlighted that plant mortality could strongly increase in the future as a consequence of increased plant water stress and air temperature (e.g., Brodribb et al., 2020; McDowell & Allen, 2015; Williams et al., 2013; Xu et al., 2019). This assessment is consistent with observations of ecological and agricultural impacts of droughts, which have reinforced by the observed increase in AED (Allen et al., 2010; Asseng et al., 2015; Breshears

et al., 2005, 2013; Carnicer et al., 2011; Lobell et al., 2011; Sánchez-Salguero et al., 2017). In opposition to this empirical evidence and also to the strong increase of drought severity as suggested by some climatic indices, LAI-based drought projections suggest small changes in drought severity in the future high-emissions scenario.

This limited increase in drought severity based on ecological metrics is difficult to be supported according to the widely known response of plants to water availability (Vicente-Serrano, Quiring, et al., 2020) and atmospheric water demand (Breshears et al., 2013; Grossiord et al., 2020), particularly in water-limited regions where meteorological droughts (e.g., southern Africa, southern North America, and the Mediterranean), and AED are projected to increase (Scheff & Frierson, 2015; Vicente-Serrano, McVicar, et al., 2020). These conditions would lead to a remarkable increase in plant water stress incompatible with increases in LAI and NPP. Thus, the unique explanation of limited increase in ecological droughts in water-limited regions, where climate aridity is projected to increase, is related to the physiological effects of the atmospheric CO₂ concentrations (Gonsamo et al., 2021; Mankin et al., 2017; Scheff et al., 2022). Several studies have showed a reduction in the leaf stomatal conductance and plant resistance to water stress in response to enhanced atmospheric CO₂ concentrations (e.g., Ainsworth & Long, 2005; Ceulemans & Mousseau, 1994; Donohue et al., 2013; Green et al., 2020). However, the effects of increasing CO₂ concentrations on ecological and agricultural drought severity are very complex (Allen et al., 2015; De Kauwe et al., 2021), and there are still several uncertainties in the assessment of these effects based on ESMs (De Kauwe et al., 2021; Gentine et al., 2019), which tend to overestimate the effects of increasing CO₂ concentrations on plant physiology (Kolby Smith et al., 2015; Marchand et al., 2020; Zhao et al., 2020). Moreover, CO₂ effects would not ameliorate plant stress during periods of water deficit, given that leaf stomatal conductance would not be controlled by CO₂ concentrations, but mostly by soil moisture content (Menezes-Silva et al., 2019; Morgan et al., 2004; Xu et al., 2019). Therefore, the assessment of future agricultural and ecological droughts based on model simulations should be considered highly uncertain given the current evidence of the responses of plants to enhanced water stress and AED and the several sources of uncertainty in the modeling of the carbon cycle by the ESMs (Padrón et al., 2022). Thus, it is difficult to support that ecological droughts will not increase in areas in which models show a strong decrease in precipitation and a remarkable increase in AED.

4.3. Uncertainties in Hydrological Drought Projections

Our results indicate that future projections of droughts quantified with soil moisture tend to resemble the pattern of the projections of drought severity using precipitation-AED. This seems to disagree with some previous studies that had suggested less increase in soil moisture deficits than the decrease in meteorological indices including AED in future drought projections (Berg & Sheffield, 2018; Milly & Dunne, 2016). This disagreement can be basically explained by the different statistical methods used to assess the future projections of these metrics. The assessment is strongly affected by the autocorrelation of some of the drought metrics (e.g., the PDSI), as well as by focusing on changes in the average values versus the tails of the complete set of the distribution values (Vicente-Serrano, Domínguez-Castro, et al., 2020). Thus, the last IPCC report has showed a strong increase in drought severity worldwide based on extreme events of the total column soil moisture, particularly during the boreal summer season (Seneviratne et al., 2021). This increase in the duration and magnitude of soil moisture deficits would be coherent with an increase in agricultural and ecological drought severity, even more considering the strong increase in AED, as projected by the CMIP models (Scheff & Frierson, 2015; Vicente-Serrano, McVicar, et al., 2020d), which would cause enhanced plant stress. Also, uncertainties in the projected Et are noticeably affect drought projections based on precipitation-Et, which is usually considered a metric of water availability. Thus, the projections of precipitation showed a stronger increase in drought duration and magnitude than projections based on precipitation-Et and runoff, which seems to introduce some incoherence regarding to what would be expected with Et in a warming world. It would be expected that hydrological droughts will not increase at similar rates of agricultural and ecological droughts, in response to increased AED. This is basically because the response of streamflow to enhanced AED is expected to be lower than to precipitation, as observed with streamflow data (Ficklin et al., 2018; Vicente-Serrano et al., 2019; Yang et al., 2018). This issue has been well-established based on the ESMs, as runoff simulations mostly respond to precipitation at short time scales (Scheff et al., 2022). However, even responding more to precipitation than to AED, it is difficult to support a smaller increase in drought severity by runoff than by precipitation under scenarios of a high increase in AED. Probably, this behavior would be explained by the suppression of Et as a consequence of the decreased leaf stomatal conductance given the enhanced atmospheric CO₂ concentrations, which would reduce the severity of hydrological droughts (Milly & Dunne, 2016; Roderick et al., 2015; Yang et al., 2019). However, a main

constrain of this assessment is that the influence of this mechanism on future Et is highly uncertain in ESMs (Vicente-Serrano, García-Herrera, et al., 2022). Moreover, Et is also observed to increase during dry periods (Zhao et al., 2022) and evaporation in surface water bodies is expected to increase in future scenarios (Wang et al., 2018). For these reasons, we find uncertain to support that hydrological droughts quantified using precipitation-Et and runoff would increase less than meteorological droughts, based on precipitation.

4.4. Relationship Among Drought Metrics

In addition to the comparative assessment of drought trends based on different drought metrics, another aspect of novelty in our study is that it assesses the spatial and temporal relationship between different drought metrics under the historical experiment and future SSP5-8.5 scenario. Specifically, we found that the temporal relationship between the precipitation-based climatic metrics (i.e., precipitation, precipitation-AED, and P-Et) is high worldwide, with some spatial exceptions (e.g., in water-limited regions for P-Et). This behavior is expected given that precipitation is a main controller of the interannual variability of climate drought indices (Tomas-Burguera et al., 2020; Vicente-Serrano et al., 2015). For example, in the case of precipitation-AED, precipitation explains more than 90% of the variability of this metric, while AED is only relevant during periods of precipitation deficit, particularly in water-limited regions (Tomas-Burguera et al., 2020). This main role of precipitation is also observed in other drought indices such as the PDSI (van der Schrier et al., 2013; Vicente-Serrano et al., 2015). On the other hand, under the SSP5-8.5 scenario, the correlation between precipitation and AED-based drought indices is expected to decrease, suggesting a greater role of AED. Nevertheless, this temporal relationship remains high in most world regions.

The close relationship found between climate drought indices in historical and future simulations contrasts with the low correlations found between climatic and ecological drought indices, given the low percentage of years when drought conditions coincide following meteorological and ecological metrics. The interannual variability of LAI and NPP showed high agreement in both the historical period and in the future scenario. This is in agreement with observations recorded in the last decades using vegetation activity from satellites (as a surrogate of the leaf area) and tree-ring growth (as a surrogate of NPP) (Vicente-Serrano et al., 2016, 2020c). Nevertheless, unexpectedly, we noted a poor relationship between the temporal evolution of both LAI and NPP and the climatic drought indices, albeit with the use of a wide set of metrics used here that highly represent plant water stress conditions (e.g., Et-AED). Moreover, this low relationship is also found between the ecological variables and soil moisture, which is one of the main factors controlling vegetation activity and carbon uptake worldwide (Green et al., 2019). This low relationship between climatic indices (and soil moisture) and ecological metrics could be explained by the uncoupling between water availability and plant water requirements as a consequence of the physiological effects of atmospheric CO₂ concentrations (as discussed above). Nevertheless, low interannual correlations were also found in the historical experiment. We consider that the low relationship between ecological drought metrics and climatic and soil moisture metrics introduces another important source of uncertainty in the assessment of the drought severity under future climate scenarios. It is expected that the agreement between NPP, LAI, and the different climatic metrics and soil moisture should be high, given the climate forcings used in the historical experiment. Thus, based on different vegetation metrics, numerous studies found strong temporal correlations between climate drought indices and soil moisture and different ecological measurements in the past decades, including satellite metrics (e.g., Bachmair et al., 2018; Vicente-Serrano et al., 2013), and tree ring growth (e.g., Orwig & Abrams, 1997; Vicente-Serrano et al., 2014). This unexpectedly low correlation between climatic drought metrics, soil moisture deficits and agricultural and ecological metrics during the historical experiment suggests that the temporal decoupling between these metrics is not related to the possible physiological effects of the enhanced CO₂ concentrations. Rather, it can probably be due to the existing limitations of the models in reproducing the real physiological response of vegetation to water deficits. Moreover, in addition to the low temporal agreement, there is a general spatial disconnection between the occurrence of climatic and ecological droughts in different regions worldwide, which introduce an additional uncertainty.

The temporal agreement between climatic drought metrics, soil moisture, precipitation-Et, and runoff is also low, both in the historical experiment and the SSP5-8.5 scenario. With the exception of the tropical and subtropical regions in the case of runoff, the remaining world showed low correlations with climatic metrics. Thus, the temporal correlations were low between the interannual variability of soil moisture and runoff in most regions of the world. This suggests that, considering climatic and hydrological drought metrics, the consistency of ESMs simulations on long temporal scales (i.e., annual) may be also affected by uncertainties. Thus, as opposed to

CMIP6 outputs, the interannual variability of observed streamflow is highly consistent with climate variables in most basins of the world (Dai, 2021).

5. Conclusions

This study provided new evidence on the interannual relationships and long-term trends between drought types based on different drought metrics obtained from ESM simulations. In the existing debate related to the assessment of drought severity in future climate projections based on different metrics from CMIP models, this study stresses some uncertainties of this assessment and the limited spatial and temporal consistency among different types of drought metrics that usually show agreement with observations, particularly with the use of hydrological (runoff and soil moisture) and ecological (LAI and NPP) drought metrics that are directly generated from model outputs. The main conclusion of this study is that the coherence of the trends and the interannual relationships between drought metrics suggest uncertainties that can largely impact any robust assessment of drought projections under scenarios of high emissions of greenhouse gases. Although some previous studies have suggested that the use of climatic drought indices could overestimate drought severity under future scenarios, this study indicates that projections based on hydrological (i.e., soil moisture and runoff) and ecological drought metrics (i.e., NPP and LAI) can be affected by important inconsistencies, particularly for the projected interannual relationship between drought metrics. We believe that still there are several sources of uncertainty, particularly linked to the plant processes and the physiological influences of the enhanced CO₂ atmospheric concentrations, which have important implications for the assessment of both ecological and hydrological droughts in future scenarios. Recent evidence highlights increased drought effects on crop systems and natural environments in response to drought events characterized by warmer conditions (Breshears et al., 2013; Fontes et al., 2018; Williams et al., 2013), but also hydrological implications given enhanced evaporation from crops, natural vegetation, and water bodies (Althoff et al., 2020; Friedrich et al., 2018; Vicente-Serrano et al., 2017). Although the response of plant physiology and hydrological processes could change in the future, with more adaptive strategies to much warmer conditions leading to a reduction in the severity of hydrological, agricultural, and ecological droughts compared to climatic droughts conditions, uncertainties persist in the assessment of processes related to elevated CO₂ concentrations, primarily due to the intricate challenge of distinguishing between radiative and physiological effects.

Drought severity projections are an extremely relevant topic with several environmental and socioeconomic implications, which deserves robust studies. Nevertheless, we must be aware that assessments based on model projections may be affected by considerable uncertainties. Indeed, improving the knowledge and modeling of the complex processes involved could reduce these uncertainties, but we are probably still far from finding this solution. A focus on simple, but robust models, as suggested by McColl et al. (2022), could be a better approach to improve the assessment of future drought severity. However, this assessment may actually be simpler, as in future periods of precipitation deficits (anthropogenic or naturally induced), the projected increased warming will undoubtedly cause more stress on hydrological and environmental systems as observed in near-present climate.

Acknowledgments

This work was supported by projects PCI2019-103631, PID2019-108589RA-I00, PID2021-122314OB-I00, SPIP2022-02857, TED2021-129152B-C41 and TED2021-129152B-C43, financed by the Ministerio de Ciencia, Innovación y Universidades, Spain (MCIN/10.13039/501100011033) and “ERDF A way of making Europe”; LINKB20080 of the programme i-LINK 2021 of CSIC and CROSSDRO financed by AXIS (Assessment of Cross(X) - sectoral climate Impacts and pathways for Sustainable transformation), JPI-Climate co-funded call of the European Commission. This study was also supported by the “Unidad Asociada CSIC–Universidade de Vigo: Grupo de Física de la Atmósfera y del Océano.” The work of Luis Gimeno-Sotelo was financially supported by CSIC Programa JAE and a “Ministerio de Ciencia, Innovación y Universidades” PhD grant (reference: PRE2022-101497). The EPhysLab group was also funded by Xunta de Galicia and ERDF, under project ED431C 2021/44 “Programa de Consolidación e Estructuración de Unidades de Investigación Competitivas.”

Data Availability Statement

The data from the CMIP6 models is available at the World Climate Research Programme (WCRP, <https://esgf-node.llnl.gov/search/cmip6/>). From that page, it is possible to download the necessary data for this research. For each CMIP6 model, which is selected in Source ID, it consists of selecting Experiment ID equal to “historical” and “ssp585”, Frequency equal to “mon” and Variable ID equal to “pr” (Precipitation), “evspsbl” (Evaporation Including Sublimation and Transpiration), “mrros” (Surface Runoff), “mrso” (Total Soil Moisture Content), “lai” (Leaf Area Index) and “npp” (Net Primary Production on Land as Carbon Mass Flux) and those necessary to compute the atmospheric evaporative demand, that is, “hurs” (Near-Surface Relative Humidity), “rsds” (Surface Downwelling Shortwave Radiation), “tasmax” (Daily Maximum Near-Surface Air Temperature), “tasmin” (Daily Minimum Near-Surface Air Temperature) and “tran” (Transpiration).

References

- Adeyeri, O. E., Zhou, W., Laux, P., Ndehedehe, C. E., Wang, X., Usman, M., & Akinsanola, A. A. (2023). Multivariate drought monitoring, propagation, and projection using bias-corrected general circulation models. *Earth's Future*, 11(4), e2022EF003303. <https://doi.org/10.1029/2022ef003303>

- Adeyeri, O. E., Zhou, W., Wang, X., Zhang, R., Laux, P., Ishola, K. A., & Usman, M. (2022). The trend and spatial spread of multisectoral climate extremes in CMIP6 models. *Scientific Reports*, 12(1), 21000. <https://doi.org/10.1038/s41598-022-25265-4>
- Ainsworth, E. A., & Long, S. P. (2005). What have we learned from 15 years of free-air CO₂ enrichment (FACE)? A meta-analytic review of the responses of photosynthesis, canopy properties and plant production to rising CO₂. *New Phytologist*, 165(2), 351–372. <https://doi.org/10.1111/j.1469-8137.2004.01224.x>
- Allen, C. D. C. D., Breshears, D. D. D., & McDowell, N. G. N. G. (2015). On underestimation of global vulnerability to tree mortality and forest die-off from hotter drought in the Anthropocene. *Ecosphere*, 6(8), art129–55. <https://doi.org/10.1890/ES15-00203.1>
- Allen, C. D. C. D., Macalady, A. K. A. K., Chenchouni, H., Bachelet, D., McDowell, N., Vennetier, M., et al. (2010). A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests. *Forest Ecology and Management*, 259(4), 660–684. <https://doi.org/10.1016/j.foreco.2009.09.001>
- Almazroui, M., Saeed, S., Saeed, F., Islam, M. N., & Ismail, M. (2020). Projections of precipitation and temperature over the South Asian countries in CMIP6. *Earth Systems and Environment*, 4(2), 297–320. <https://doi.org/10.1007/s41748-020-00157-7>
- Alsafadi, K., Al-Ansari, N., Mokhtar, A., Mohammed, S., Elbeltagi, A., Sh Sammen, S., & Bi, S. (2022). An evapotranspiration deficit-based drought index to detect variability of terrestrial carbon productivity in the Middle East. *Environmental Research Letters*, 17(1), 014051. <https://doi.org/10.1088/1748-9326/ac4765>
- Althoff, D., Rodrigues, L. N., & da Silva, D. D. (2020). Impacts of climate change on the evaporation and availability of water in small reservoirs in the Brazilian savannah. *Climatic Change*, 159(2), 215–232. <https://doi.org/10.1007/s10584-020-02656-y>
- Asseng, S., Ewert, F., Martre, P., Rötter, R. P., Lobell, D. B., Cammarano, D., et al. (2015). Rising temperatures reduce global wheat production. *Nature Climate Change*, 5(2), 143–147. <https://doi.org/10.1038/nclimate2470>
- Bachmair, S., Kohn, I., & Stahl, K. (2015). Exploring the link between drought indicators and impacts. *Natural Hazards and Earth System Sciences*, 15(6), 1381–1397. <https://doi.org/10.5194/nhess-15-1381-2015>
- Bachmair, S., Svensson, C., Hannaford, J., Barker, L. J., & Stahl, K. (2016). A quantitative analysis to objectively appraise drought indicators and model drought impacts. *Hydrology and Earth System Sciences*, 20(7), 2589–2609. <https://doi.org/10.5194/hess-20-2589-2016>
- Bachmair, S., Tanguy, M., Hannaford, J., & Stahl, K. (2018). How well do meteorological indicators represent agricultural and forest drought across Europe? *Environmental Research Letters*, 13(3), 034042. <https://doi.org/10.1088/1748-9326/aaafda>
- Barella-Ortiz, A., & Quintana-Seguí, P. (2018). Evaluation of drought representation and propagation in Regional Climate Model simulations over Spain. *Hydrology and Earth System Sciences Discussions*, 1–32. <https://doi.org/10.5194/hess-2018-603>
- Berg, A., & McColl, K. A. (2021). No projected global drylands expansion under greenhouse warming. *Nature Climate Change*, 11(4), 331–337. <https://doi.org/10.1038/s41558-021-01007-8>
- Berg, A., & Sheffield, J. (2018). Climate change and drought: The soil moisture perspective. *Current Climate Change Reports*, 4(2), 180–191. <https://doi.org/10.1007/s40641-018-0095-0>
- Breshears, D. D., Adams, H. D., Eamus, D., McDowell, N. G., Law, D. J., Will, R. E., et al. (2013). The critical amplifying role of increasing atmospheric moisture demand on tree mortality and associated regional die-off. *Frontiers in Plant Science*, 4(AUG). <https://doi.org/10.3389/fpls.2013.00266>
- Breshears, D. D., Cobb, N. S., Rich, P. M., Price, K. P., Allen, C. D., Balice, R. G., et al. (2005). Regional vegetation die-off in response to global-change-type drought. *Proceedings of the National Academy of Sciences of the United States of America*, 102(42), 15144–15148. <https://doi.org/10.1073/pnas.0505734102>
- Brodribb, T. J., Powers, J., Cochard, H., & Choat, B. (2020). Hanging by a thread? Forests and drought. *Science*, 368(6488), 261–266. American Association for the Advancement of Science. <https://doi.org/10.1126/science.aat7631>
- Carnicer, J., Coll, M., Ninyerola, M., Pons, X., Sánchez, G., & Peñuelas, J. (2011). Widespread crown condition decline, food web disruption, and amplified tree mortality with increased climate change-type drought. *Proceedings of the National Academy of Sciences of the United States of America*, 108(4), 1474–1478. <https://doi.org/10.1073/pnas.1010070108>
- Ceulemans, R., & Mousseau, M. (1994). Tansley review No. 71. Effects of elevated atmospheric CO₂ on woody plants. *New Phytologist*, 127(3), 425–446. [Wiley, New Phytologist Trust]. <https://doi.org/10.1111/j.1469-8137.1994.tb03961.x>
- Chen, Y., Romps, D. M., Seeley, J. T., Veraverbeke, S., Riley, W. J., Mekonnen, Z. A., & Randerson, J. T. (2021). Future increases in Arctic lightning and fire risk for permafrost carbon. *Nature Climate Change*, 11(5), 404–410. <https://doi.org/10.1038/s41558-021-01011-y>
- Cook, B. I., Mankin, J. S., Marvel, K., Williams, A. P., Smerdon, J. E., & Anchukaitis, K. J. (2020). Twenty-first century drought projections in the CMIP6 forcing scenarios. *Earth's Future*, 8(6), e2019EF001461. John Wiley & Sons, Ltd. <https://doi.org/10.1029/2019EF001461>
- Cook, B. I., Smerdon, J. E., Seager, R., & Coats, S. (2014). Global warming and 21st century drying. *Climate Dynamics*, 43(9), 2607–2627. <https://doi.org/10.1007/s00382-014-2075-y>
- Dai, A. (2012). Increasing drought under global warming in observations and models. *Nature Climate Change*, 3(1), 52–58. Nature Publishing Group. <https://doi.org/10.1038/nclimate1633>
- Dai, A. (2021). Hydroclimatic trends during 1950–2018 over global land. *Climate Dynamics*, 56(11), 4027–4049. <https://doi.org/10.1007/s00382-021-05684-1>
- De Kauwe, M. G., Medlyn, B. E., & Tissue, D. T. (2021). To what extent can rising [CO₂] ameliorate plant drought stress? *New Phytologist*, 231(6), 2118–2124. John Wiley & Sons, Ltd. <https://doi.org/10.1111/nph.17540>
- Donohue, R. J., Roderick, M. L., McVicar, T. R., & Farquhar, G. D. (2013). Impact of CO₂ fertilization on maximum foliage cover across the globe's warm, arid environments. *Geophysical Research Letters*, 40(12), 3031–3035. <https://doi.org/10.1002/grl.50563>
- Douville, H., Raghavan, K., & Renwick, J. (2021). Water cycle changes. In *Climate change 2021: The physical climate, basis. Contribution of working group I to the sixth assessment report of the intergovernmental panel on climate change*.
- Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., & Taylor, K. E. (2016). Overview of the Coupled Model Inter-comparison Project Phase 6 (CMIP6) experimental design and organization. *Geoscientific Model Development*, 9(5), 1937–1958. <https://doi.org/10.5194/gmd-9-1937-2016>
- Ficklin, D. L., Abatzoglou, J. T., Robeson, S. M., Null, S. E., & Knouft, J. H. (2018). Natural and managed watersheds show similar responses to recent climate change. *Proceedings of the National Academy of Sciences of the United States of America*, 115(34), 201801026–8557. National Academy of Sciences. <https://doi.org/10.1073/PNAS.1801026115>
- Fleig, A. K., Tallaksen, L. M., Hisdal, H., & Demuth, S. (2006). A global evaluation of streamflow drought characteristics. *Hydrology and Earth System Sciences*, 10(4), 535–552. <https://doi.org/10.5194/hess-10-535-2006>
- Fontes, C. G., Dawson, T. E., Jardine, K., McDowell, N., Gimenez, B. O., Anderegg, L., et al. (2018). Dry and hot: The hydraulic consequences of a climate change-type drought for Amazonian trees. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 373(1760), 20180209. <https://doi.org/10.1098/rstb.2018.0209>

- Ford, T. W., & Quiring, S. M. (2019). *Comparison of contemporary in situ, model, and satellite remote sensing soil moisture with a focus on drought monitoring*. Water Resources Research.
- Friedrich, K., Grossman, R. L., Huntington, J., Blanken, P. D., Lenters, J., Holman, K. D., et al. (2018). Reservoir evaporation in the Western United States. *Bulletin of the American Meteorological Society*, 99(1), 167–187. <https://doi.org/10.1175/BAMS-D-15-00224.1>
- Gentine, P., Green, J., Guerin, M., Humphrey, V., Seneviratne, S., Zhang, Y., & Zhou, S. (2019). Coupling between the terrestrial carbon and water cycles – A review. *Environmental Research Letters*, 14(8), 083003. <https://doi.org/10.1088/1748-9326/ab22d6>
- Gonsamo, A., Ciais, P., Miralles, D. G., Sitch, S., Dorigo, W., Lombardozzi, D., et al. (2021). Greening drylands despite warming consistent with carbon dioxide fertilization effect. *Global Change Biology*. John Wiley & Sons, Ltd, 27(14), 3336–3349. <https://doi.org/10.1111/gcb.15658>
- Green, J. K., Berry, J., Ciais, P., Zhang, Y., & Gentine, P. (2020). Amazon rainforest photosynthesis increases in response to atmospheric dryness. *Science Advances*, 6(47), eabb7232. <https://doi.org/10.1126/sciadv.abb7232>
- Green, J. K., Seneviratne, S. I., Berg, A. M., Findell, K. L., Hagemann, S., Lawrence, D. M., & Gentine, P. (2019). Large influence of soil moisture on long-term terrestrial carbon uptake. *Nature*, 565(7740), 476–479. <https://doi.org/10.1038/s41586-018-0848-x>
- Greve, P., Roderick, M. L., Ukkola, A. M., & Wada, Y. (2019). The aridity Index under global warming. *Environmental Research Letters*, 14(12), 124006. <https://doi.org/10.1088/1748-9326/ab5046>
- Grossiord, C., Buckley, T. N., Cernusak, L. A., Novick, K. A., Poulter, B., Siegwolf, R. T. W., et al. (2020). Plant responses to rising vapor pressure deficit. *New Phytologist*, 226(6), 1550–1566. John Wiley & Sons, Ltd. <https://doi.org/10.1111/nph.16485>
- Guo, H., He, S., Li, M., Bao, A., Chen, T., Zheng, G., & De Maeyer, P. (2022). Future changes of drought characteristics in Coupled Model Intercomparison Project phase 6 Shared Socioeconomic Pathway scenarios over Central Asia. *International Journal of Climatology*, 42(7), 3888–3908. <https://doi.org/10.1002/joc.7450>
- Gutiérrez, J. M., Jones, R. G., Narisma, G. T., Alves, L. M., Amjad, M., Gorodetskaya, I. V., et al. (2021). Atlas. In V. Masson-Delmotte, P. Zhai, A. Pirani, S. L. Connors, C. Péan, S. Berger, et al. (Eds.), *Climate change 2021: The physical science basis. Contribution of working group I to the sixth assessment report of the intergovernmental panel on climate change* (pp. 1927–2058). Cambridge University Press. <https://doi.org/10.1017/9781009157896.021>
- Hlavinka, P., Trnka, M., Semerádová, D., Dubrovský, M., Žalud, Z., & Možný, M. (2009). Effect of drought on yield variability of key crops in Czech Republic. *Agricultural and Forest Meteorology*, 149(3), 431–442. <https://doi.org/10.1016/j.agrformet.2008.09.004>
- Jiang, W., Niu, Z., Wang, L., Yao, R., Gui, X., Xiang, F., & Ji, Y. (2022). Impacts of drought and climatic factors on vegetation dynamics in the Yellow River Basin and Yangtze River Basin, China. *Remote Sensing*, 14(4), 930. <https://doi.org/10.3390/rs14040930>
- Kendall, M. G. (1938). A new measure of rank correlation. *Biometrika*, 30(1/2), 81–93. <https://doi.org/10.2307/2332226>
- Kendall, M. G. (1948). Rank correlation methods.
- Kim, D., & Rhee, J. (2016). A drought index based on actual evapotranspiration from the Bouchet hypothesis. *Geophysical Research Letters*, 43(19), 10277–10285. <https://doi.org/10.1002/2016GL070302>
- Kim, S. (2015). ppcor: An R package for a fast calculation to semi-partial correlation coefficients. *Communications for Statistical Applications and Methods*, 22(6), 665–674. <https://doi.org/10.5351/csam.2015.22.6.665>
- Kolby Smith, W., Reed, S. C., Cleveland, C. C., Ballantyne, A. P., Anderegg, W. R. L., Wieder, W. R., et al. (2015). Large divergence of satellite and Earth system model estimates of global terrestrial CO₂ fertilization. *Nature Climate Change*, 6(3), 306–310. Nature Publishing Group. <https://doi.org/10.1038/nclimate2879>
- Kolmogorov, A. N. (1933). Sulla determinazione empirica di una legge di distribuzione. *Giornale dell'Istituto Italiano Degli Attuari*, 4, 83–91.
- Li, F., Xiao, J., Chen, J., Ballantyne, A., Jin, K., Li, B., et al. (2023). Global water use efficiency saturation due to increased vapor pressure deficit. *Science*, 381(6658), 672–677. <https://doi.org/10.1126/science.adf5041>
- Li, G., Sun, S., Han, J., Yan, J., Liu, W., Wei, Y., et al. (2019). Impacts of Chinese Grain for Green program and climate change on vegetation in the Loess Plateau during 1982–2015. *Science of the Total Environment*, 660, 177–187. <https://doi.org/10.1016/j.scitotenv.2019.01.028>
- Li, H., Li, Z., Chen, Y., Xiang, Y., Liu, Y., Kayumba, P. M., & Li, X. (2021). Drylands face potential threat of robust drought in the CMIP6 SSPs scenarios. *Environmental Research Letters*, 16(11), 114004. <https://doi.org/10.1088/1748-9326/ac2bce>
- Li, J., Wang, Z., & Lai, C. (2020). Severe drought events inducing large decrease of net primary productivity in mainland China during 1982–2015. *Science of the Total Environment*, 703, 703. <https://doi.org/10.1016/j.scitotenv.2019.135541>
- Liu, Y., Kumar, M., Katul, G. G., Feng, X., & Konings, A. G. (2020). Plant hydraulics accentuates the effect of atmospheric moisture stress on transpiration. *Nature Climate Change*, 10(7), 691–695. <https://doi.org/10.1038/s41558-020-0781-5>
- Lloyd-Hughes, B. (2014). The impracticality of a universal drought definition. *Theoretical and Applied Climatology*, 117(3–4), 607–611. <https://doi.org/10.1007/s00704-013-1025-7>
- Lobell, D. B. (2014). Climate change adaptation in crop production: Beware of illusions. *Global Food Security*, 3(2), 72–76. <https://doi.org/10.1016/j.gfs.2014.05.002>
- Lobell, D. B., Schlenker, W., & Costa-Roberts, J. (2011). Climate trends and global crop production since 1980. *Science*, 333(6042), 616–620. <https://doi.org/10.1126/science.1204531>
- Lu, J., Carbone, G. J., & Grego, J. M. (2019). Uncertainty and hotspots in 21st century projections of agricultural drought from CMIP5 models. *Scientific Reports*, 9(1), 4922. <https://doi.org/10.1038/s41598-019-41196-z>
- Mankin, J. S., Smerdon, J. E., Cook, B. I., Williams, A. P., & Seager, R. (2017). The curious case of projected twenty-first-century drying but greening in the American west. *Journal of Climate*, 30(21), 8689–8710. American Meteorological Society. <https://doi.org/10.1175/JCLI-D-17-0213.1>
- Mann, H. B. (1945). Nonparametric tests against trend. *Econometrica: Journal of the Econometric Society*, 13(3), 245–259. <https://doi.org/10.2307/1907187>
- Marchand, W., Girardin, M. P., Hartmann, H., Depardieu, C., Isabel, N., Gauthier, S., et al. (2020). Strong overestimation of water-use efficiency responses to rising CO₂ in tree-ring studies. *Global Change Biology*. John Wiley & Sons, Ltd, 26(8), 4538–4558. <https://doi.org/10.1111/gcb.15166>
- Martin, E. R. (2018). Future projections of global pluvial and drought event characteristics. *Geophysical Research Letters*, 45(21). <https://doi.org/10.1029/2018GL079807>
- McColl, K. A., Roderick, M. L., Berg, A., & Scheff, J. (2022). The terrestrial water cycle in a warming world. *Nature Climate Change*, 12(7), 604–606. <https://doi.org/10.1038/s41558-022-01412-7>
- McDowell, N. G., & Allen, C. D. (2015). Darcy's law predicts widespread forest mortality under climate warming. *Nature Climate Change*, 5(7), 669–672. <https://doi.org/10.1038/nclimate2641>
- Menezes-Silva, P. E., Loram-Lourenço, L., Alves, R. D. F. B., Sousa, L. F., Almeida, S. E. d. S., & Farnese, F. S. (2019). Different ways to die in a changing world: Consequences of climate change for tree species performance and survival through an ecophysiological perspective. *Ecology and Evolution*, 9(20), 11979–11999. <https://doi.org/10.1002/ece3.5663>

- Meng, Y., Hao, Z., Feng, S., Zhang, X., & Hao, F. (2022). Increase in compound dry-warm and wet-warm events under global warming in CMIP6 models. *Global and Planetary Change*, 210, 103773. <https://doi.org/10.1016/j.gloplacha.2022.103773>
- Miguez-Macho, G., & Fan, Y. (2021). Spatiotemporal origin of soil water taken up by vegetation. *Nature*, 598(7882), 624–628. <https://doi.org/10.1038/s41586-021-03958-6>
- Milly, P. C. D., & Dunne, K. A. (2016). Potential evapotranspiration and continental drying. *Nature Climate Change*, 6(10), 946–949. <https://doi.org/10.1038/nclimate3046>
- Morgan, J. A., Pataki, D. E., Körner, C., Clark, H., Del Grosso, S. J., & Grünzweig, J. M. (2004). Water relations in grassland and desert ecosystems exposed to elevated atmospheric CO₂. *Oecologia*, 140(1), 11–25. <https://doi.org/10.1007/s00442-004-1550-2>
- Naumann, G., Alfieri, L., Wyser, K., Mentaschi, L., Betts, R. A., Carrao, H., et al. (2018). Global changes in drought conditions under different levels of warming. *Geophysical Research Letters*, 45(7), 3285–3296. <https://doi.org/10.1002/2017GL076521>
- Naumann, G., Cammalleri, C., Mentaschi, L., & Feyen, L. (2021). Increased economic drought impacts in Europe with anthropogenic warming. *Nature Climate Change*, 11(6), 485–491. <https://doi.org/10.1038/s41558-021-01044-3>
- Ndehedehe, C. E., Ferreira, V. G., Adeyeri, O. E., Correa, F. M., Usman, M., Oussou, F. E., et al. (2023). Global assessment of drought characteristics in the Anthropocene. *Resources, Environment and Sustainability*, 12, 100105. <https://doi.org/10.1016/j.resenv.2022.100105>
- O'Connor, P., Murphy, C., Matthews, T., & Wilby, R. (2022). Relating drought indices to impacts reported in newspaper articles. *International Journal of Climatology*, 43(4), 1796–1816. <https://doi.org/10.1002/joc.7946>
- Orwig, D. A., & Abrams, M. D. (1997). Variation in radial growth responses to drought among species, site, and canopy strata. *Trees: Structure and Function*, 11(8), 474–484. <https://doi.org/10.1007/s004680050110>
- Padrón, R. S., Gudmundsson, L., Decharme, B., Ducharme, A., Lawrence, D. M., Mao, J., et al. (2020). Observed changes in dry-season water availability attributed to human-induced climate change. *Nature Geoscience*, 13(7), 477–481. <https://doi.org/10.1038/s41561-020-0594-1>
- Padrón, R. S., Gudmundsson, L., Liu, L., Humphrey, V., & Seneviratne, S. I. (2022). Drivers of intermodel uncertainty in land carbon sink projections. *Biogeosciences*, 19(23), 5435–5448. <https://doi.org/10.5194/bg-19-5435-2022>
- Papalexiou, S. M., Rajulapati, C. R., Andreadis, K. M., Foufoula-Georgiou, E., Clark, M. P., & Trenberth, K. E. (2021). Probabilistic evaluation of drought in CMIP6 simulations. *Earth's Future*. John Wiley & Sons, Ltd, 9(10), e2021EF002150. <https://doi.org/10.1029/2021EF002150>
- Pereira, L. S., Allen, R. G., Smith, M., & Raes, D. (2015). Crop evapotranspiration estimation with FAO56: Past and future. *Agricultural Water Management*, 147, 4–20. <https://doi.org/10.1016/j.agwat.2014.07.031>
- Quiring, S. M., & Papakryiakou, T. N. (2003). An evaluation of agricultural drought indices for the Canadian prairies. *Agricultural and Forest Meteorology*, 118(1), 49–62. [https://doi.org/10.1016/S0168-1923\(03\)00072-8](https://doi.org/10.1016/S0168-1923(03)00072-8)
- Rettie, F. M., Gayler, S., Weber, T. K., Tesfaye, K., & Streck, T. (2023). High-resolution CMIP6 climate projections for Ethiopia using the gridded statistical downscaling method. *Scientific Data*, 10(1), 442. <https://doi.org/10.1038/s41597-023-02337-2>
- Ridder, N. N., Ukkola, A. M., Pitman, A. J., & Perkins-Kirkpatrick, S. E. (2022). Increased occurrence of high impact compound events under climate change. *npj Climate and Atmospheric Science*, 5(1), 3. <https://doi.org/10.1038/s41612-021-00224-4>
- Roderick, M. L., Greve, P., & Farquhar, G. D. (2015a). On the assessment of aridity with changes in atmospheric CO₂. *Water Resources Research*, 51(7), 5450–5463. <https://doi.org/10.1002/2015WR017031>
- Sánchez-Salguero, R., Camarero, J. J., Carrer, M., Gutiérrez, E., Alla, A. Q., Andreu-Hayles, L., et al. (2017). Climate extremes and predicted warming threaten Mediterranean Holocene firs forests refugia. *Proceedings of the National Academy of Sciences of the United States of America*, 114(47), E10142–E10150. <https://doi.org/10.1073/pnas.1708109114>
- Scheff, J. (2018). Drought indices, drought impacts, CO₂, and warming: A historical and geologic perspective. *Current Climate Change Reports*, 4(2), 202–209. <https://doi.org/10.1007/s40641-018-0094-1>
- Scheff, J., Coats, S., & Laguë, M. M. (2022). Why do the global warming responses of land-surface models and climatic dryness metrics disagree? *Earth's Future*, 10(8). <https://doi.org/10.1029/2022EF002814>
- Scheff, J., & Frierson, D. M. W. (2015). Terrestrial aridity and its response to greenhouse warming across CMIP5 climate models. *Journal of Climate*, 28(14), 5583–5600. <https://doi.org/10.1175/JCLI-D-14-00480.1>
- Scheff, J., Mankin, J. S., Coats, S., & Liu, H. (2021). CO₂-plant effects do not account for the gap between dryness indices and projected dryness impacts in CMIP6 or CMIP5. *Environmental Research Letters*, 16(3), 034018. IOP Publishing. <https://doi.org/10.1088/1748-9326/abd8fd>
- Seneviratne, S., Zhang, X., Adnan, M., Badi, W., Dereczynski, C., Di Luca, A., et al. (2021). Weather and climate extreme events in a changing climate. In V. MassonDelmotte & E. Al (Eds.), *Climate change 2021: The physical science basis. Contribution of working group I to the sixth assessment report of the intergovernmental panel on climate change*. Cambridge University Press.
- Smirnov, N. (1948). Table for estimating the goodness of fit of empirical distributions. *The Annals of Mathematical Statistics*, 19(2), 279–281. <https://doi.org/10.1214/aoms/1177730256>
- Spinoni, J., Barbosa, P., Buccignani, E., Cassano, J., Cavazos, T., Christensen, J. H., & Dosio, A. (2020). Future global meteorological drought hot spots: A study based on CORDEX data. *Journal of Climate*, 33(9), 3635–3661. American Meteorological Society. <https://doi.org/10.1175/JCLI-D-19-0084.1>
- Stagge, J. H., Kohn, I., Tallaksen, L. M., & Stahl, K. (2015). Modeling drought impact occurrence based on meteorological drought indices in Europe. *Journal of Hydrology*, 530, 37–50. <https://doi.org/10.1016/j.jhydrol.2015.09.039>
- Stagge, J. H., Tallaksen, L. M., Gudmundsson, L., Van Loon, A. F., & Stahl, K. (2015). Candidate distributions for climatological drought indices (SPI and SPEI). *International Journal of Climatology*, 35(13), 4027–4040. John Wiley and Sons Ltd. <https://doi.org/10.1002/joc.4267>
- Stephenson, N. L. (1990). Climatic control of vegetation distribution: The role of the water balance. *The American Naturalist*, 135(5), 649–670. <https://doi.org/10.1086/285067>
- Stocker, T. F., Qin, D., Plattner, G.-K., Alexander, L. V., Allen, S. K., Bindoff, N. L., & Xie, S.-P. (2013). Technical summary. In *Climate change 2013: The physical science basis*.
- Tallaksen, L. M., Madsen, H., & Clausen, B. (1997). On the definition and modelling of streamflow drought duration and deficit volume. *Hydrological Sciences Journal*, 42(1), 15–33. Taylor & Francis Group. <https://doi.org/10.1080/02626669709492003>
- Tallaksen, L. M., & Stahl, K. (2014). Spatial and temporal patterns of large-scale droughts in Europe: Model dispersion and performance. *Geophysical Research Letters*, 41(2), 429–434. <https://doi.org/10.1002/2013GL058573>
- Thorarindottir, T. L., Sillmann, J., Haugen, M., Gissibl, N., & Sandstad, M. (2020). Evaluation of CMIP5 and CMIP6 simulations of historical surface air temperature extremes using proper evaluation methods. *Environmental Research Letters*, 15(12), 124041. <https://doi.org/10.1088/1748-9326/abc778>
- Tomas-Burguera, M., Vicente-Serrano, S. M., Peña-Angulo, D., Domínguez-Castro, F., Noguera, I., & El Kenawy, A. (2020a). Global characterization of the varying responses of the standardized precipitation evapotranspiration index to atmospheric evaporative demand. *Journal of Geophysical Research: Atmospheres*, 125(17). Blackwell Publishing Ltd. <https://doi.org/10.1029/2020JD033017>

- Ukkola, A. M., De Kauwe, M. G., Roderick, M. L., Abramowitz, G., & Pitman, A. J. (2020). Robust future changes in meteorological drought in CMIP6 projections despite uncertainty in precipitation. *Geophysical Research Letters*, 47(11), e2020GL087820. John Wiley & Sons, Ltd. <https://doi.org/10.1029/2020GL087820>
- van den Hurk, B., Best, M., Dirmeyer, P., Pitman, A., Polcher, J., & Santanello, J. (2011). Acceleration of land surface model development over a decade of glass. *Bulletin of the American Meteorological Society*, 92(12), 1593–1600. American Meteorological Society. <https://doi.org/10.1175/BAMS-D-11-00007.1>
- van der Schrier, G., Barichivich, J., Briffa, K. R., & Jones, P. D. (2013). A scPDSI-based global data set of dry and wet spells for 1901–2009. *Journal of Geophysical Research: Atmospheres*, 118(10), 4025–4048. <https://doi.org/10.1002/jgrd.50355>
- Van Loon, A. F., Gleeson, T., Clark, J., Van Dijk, A. I. J. M., Stahl, K., Hannaford, J., et al. (2016). Drought in the anthropocene. *Nature Geoscience*, 9(2), 89–91. <https://doi.org/10.1038/ngeo2646>
- Vicente-Serrano, S. M., & Beguería, S. (2016). Comment on “Candidate distributions for climatological drought indices (SPI and SPEI)” by James H. Stagge et al. *International Journal of Climatology*, 36(4), 2120–2131. <https://doi.org/10.1002/joc.4474>
- Vicente-Serrano, S. M., Beguería, S., & López-Moreno, J. I. (2010). A multiscale drought index sensitive to global warming: The standardized precipitation evapotranspiration index. *Journal of Climate*, 23(7), 1696–1718. <https://doi.org/10.1175/2009JCLI2909.1>
- Vicente-Serrano, S. M., Beguería, S., Lorenzo-Lacruz, J., Camarero, J. J., López-Moreno, J. I., Azorin-Molina, C., et al. (2012). Performance of drought indices for ecological, agricultural, and hydrological applications. *Earth Interactions*, 16(10), 1–27. <https://doi.org/10.1175/2012EI000434.1>
- Vicente-Serrano, S. M., Camarero, J. J., & Azorin-Molina, C. (2014). Diverse responses of forest growth to drought time-scales in the Northern Hemisphere. *Global Ecology and Biogeography*, 23(9), 1019–1030. <https://doi.org/10.1111/geb.12183>
- Vicente-Serrano, S. M., Camarero, J. J., Olano, J. M., Martín-Hernández, N., Peña-Gallardo, M., Tomás-Burguera, M., et al. (2016). Diverse relationships between forest growth and the Normalized Difference Vegetation Index at a global scale. *Remote Sensing of Environment*, 187, 14–29. <https://doi.org/10.1016/j.rse.2016.10.001>
- Vicente-Serrano, S. M., der Schrier, G., Beguería, S., Azorin-Molina, C., & Lopez-Moreno, J.-I. (2015). Contribution of precipitation and reference evapotranspiration to drought indices under different climates. *Journal of Hydrology*, 526, 42–54. <https://doi.org/10.1016/j.jhydrol.2014.11.025>
- Vicente-Serrano, S. M., Domínguez-Castro, F., McVicar, T. R., Tomas-Burguera, M., Peña-Gallardo, M., Noguera, I., et al. (2020). Global characterization of hydrological and meteorological droughts under future climate change: The importance of timescales, vegetation-CO₂ feedbacks and changes to distribution functions. *International Journal of Climatology*, 40(5), 2557–2567. <https://doi.org/10.1002/joc.6350>
- Vicente-Serrano, S. M., García-Herrera, R., Peña-Angulo, D., Tomas-Burguera, M., Domínguez-Castro, F., Noguera, I., et al. (2022). Do CMIP models capture long-term observed annual precipitation trends? *Climate Dynamics*, 58(9–10), 2825–2842. <https://doi.org/10.1007/s00382-021-06034-x>
- Vicente-Serrano, S. M., Gouveia, C., Camarero, J. J., Beguería, S., Trigo, R., López-Moreno, J. I., & Sanchez-Lorenzo, A. (2013). Response of vegetation to drought time-scales across global land biomes. *Proceedings of the National Academy of Sciences of the United States of America*, 110(1), 52–57. National Academy of Sciences. <https://doi.org/10.1073/pnas.1207068110>
- Vicente-Serrano, S. M., Martín-Hernández, N., Camarero, J. J. J., Gazol, A., Sánchez-Salguero, R., Peña-Gallardo, M., & Diego Galván, J. (2020). Linking tree-ring growth and satellite-derived gross primary growth in multiple forest biomes. Temporal-scale matters. *Ecological Indicators*, 108, 105753. <https://doi.org/10.1016/j.ecolind.2019.105753>
- Vicente-Serrano, S. M., McVicar, T. R., Miralles, D. G., Yang, Y., & Tomas-Burguera, M. (2020). Unraveling the influence of atmospheric evaporative demand on drought and its response to climate change. *Wiley Interdisciplinary Reviews: Climate Change*, 11(2), e632. <https://doi.org/10.1002/wcc.632>
- Vicente-Serrano, S. M., Miralles, D. G., Domínguez-Castro, F., Azorin-Molina, C., El Kenawy, A., McVicar, T. R., et al. (2018). Global assessment of the standardized evapotranspiration deficit index (SEDI) for drought analysis and monitoring. *Journal of Climate*, 31(14), 5371–5393. <https://doi.org/10.1175/jcli-d-17-0775.1>
- Vicente-Serrano, S. M., Miralles, D. G., McDowell, N., Brodribb, T., Domínguez-Castro, F., & Leung, R. (2022). The uncertain role of rising atmospheric CO₂ on global plant transpiration. *Earth-Science Reviews*, 230, 104055. <https://doi.org/10.1016/j.earscirev.2022.104055>
- Vicente-Serrano, S. M., Peña-Gallardo, M., Hannaford, J., Murphy, C., Lorenzo-Lacruz, J., Domínguez-Castro, F., et al. (2019). Climate, irrigation, and land-cover change explain streamflow trends in countries bordering the Northeast Atlantic. *Geophysical Research Letters*, 46(19), 10821–10833. <https://doi.org/10.1029/2019gl084084>
- Vicente-Serrano, S. M., Quiring, S. M., Peña-Gallardo, M., Yuan, S., & Domínguez-Castro, F. (2020). A review of environmental droughts: Increased risk under global warming? *Earth-Science Reviews*, 201, 102953. <https://doi.org/10.1016/j.earscirev.2019.102953>
- Vicente-Serrano, S. M., Zabalza-Martínez, J., Borrás, G., López-Moreno, J. I., Pla, E., Pascual, D., et al. (2017). Effect of reservoirs on streamflow and river regimes in a heavily regulated river basin of Northeast Spain. *Catena*, 149, 727–741. <https://doi.org/10.1016/j.catena.2016.03.042>
- Wang, T., Tu, X., Singh, V. P., Chen, X., & Lin, K. (2021). Global data assessment and analysis of drought characteristics based on CMIP6. *Journal of Hydrology*, 596, 126091. <https://doi.org/10.1016/j.jhydrol.2021.126091>
- Wang, W., Lee, X., Xiao, W., Liu, S., Schultz, N., Wang, Y., et al. (2018). Global lake evaporation accelerated by changes in surface energy allocation in a warmer climate. *Nature Geoscience*, 11(6), 410–414. <https://doi.org/10.1038/s41561-018-0114-8>
- Wilhite, D. A., & Buchanan-Smith, M. (2005). Drought as hazard: Understanding the natural and social context. In *Drought and water crises: Science, technology, and management issues*.
- Williams, A. P., Allen, C. D., Macalady, A. K., Griffin, D., Woodhouse, C. A., Meko, D. M., & McDowell, N. G. (2013). Temperature as a potent driver of regional forest drought stress and tree mortality. *Nature Climate Change*, 3(3), 292–297. <https://doi.org/10.1038/nclimate1693>
- Wu, H., Lei, H., Lu, W., & Liu, Z. (2022). Future changes in precipitation over the upper Yangtze River basin based on bias correction spatial downscaling of models from CMIP6. *Environmental Research Communications*, 4(4), 045002. <https://doi.org/10.1088/2515-7620/ac620e>
- Xu, C., McDowell, N. G., Fisher, R. A., Wei, L., Sevanto, S., Christoffersen, B. O., et al. (2019). Increasing impacts of extreme droughts on vegetation productivity under climate change. *Nature Climate Change*, 9(12), 948–953. <https://doi.org/10.1038/s41558-019-0630-6>
- Yang, Y., Roderick, M. L., Yang, D., Wang, Z., Ruan, F., McVicar, T. R., et al. (2021). Streamflow stationarity in a changing world. *Environmental Research Letters*, 16(6), 064096. <https://doi.org/10.1088/1748-9326/ac08c1>
- Yang, Y., Roderick, M. L., Zhang, S., McVicar, T. R., & Donohue, R. J. (2019). Hydrologic implications of vegetation response to elevated CO₂ in climate projections. *Nature Climate Change*, 9(1), 44–48. <https://doi.org/10.1038/s41558-018-0361-0>
- Yang, Y., Zhang, S., McVicar, T. R., Beck, H. E., Zhang, Y., & Liu, B. (2018). Disconnection between trends of atmospheric drying and continental runoff. *Water Resources Research*, 54(7), 4700–4713. <https://doi.org/10.1029/2018WR022593>
- Yuan, S., & Quiring, S. M. (2017). Evaluation of soil moisture in CMIP5 simulations over the contiguous United States using in situ and satellite observations. *Hydrology and Earth System Sciences*, 21(4), 2203–2218. <https://doi.org/10.5194/hess-21-2203-2017>

- Zeng, J., Li, J., Lu, X., Wei, Z., Shangguan, W., Zhang, S., et al. (2022). Assessment of global meteorological, hydrological and agricultural drought under future warming based on CMIP6. *Atmospheric and Oceanic Science Letters*, 15(1), 100143. <https://doi.org/10.1016/j.aosl.2021.100143>
- Zhang, X., Li, M., Ma, Z., Yang, Q., Lv, M., & Clark, R. (2019). Assessment of an evapotranspiration deficit drought index in relation to impacts on ecosystems. *Advances in Atmospheric Sciences*, 36(11), 1273–1287. <https://doi.org/10.1007/s00376-019-9061-6>
- Zhao, M., Liu, Y., & Konings, A. G. (2022). Evapotranspiration frequently increases during droughts. *Nature Climate Change*, 12(11), 1024–1030. <https://doi.org/10.1038/s41558-022-01505-3>
- Zhao, Q., Zhu, Z., Zeng, H., Zhao, W., & Myneni, R. B. (2020). Future greening of the Earth may not be as large as previously predicted. *Agricultural and Forest Meteorology*, 292–293, 108111. <https://doi.org/10.1016/j.agrformet.2020.108111>
- Zhao, T., & Dai, A. (2021). CMIP6 model-projected hydroclimatic and drought changes and their causes in the 21st century. *Journal of Climate*, 1–58. American Meteorological Society. <https://doi.org/10.1175/JCLI-D-21-0442.1>
- Zhao, T., & Dai, A. (2022). CMIP6 model-projected hydroclimatic and drought changes and their causes in the twenty-first century. *Journal of Climate*, 35(3), 897–921. American Meteorological Society. <https://doi.org/10.1175/JCLI-D-21-0442.1>
- Zhu, Y., & Yang, S. (2021). Interdecadal and interannual evolution characteristics of the global surface precipitation anomaly shown by CMIP5 and CMIP6 models. *International Journal of Climatology*, 41(S1), E1100–E1118. <https://doi.org/10.1002/joc.6756>