

Influence of conventional resistance training compared to core exercises on road cycling power output

Sebastian Sitko¹, Isaac López-Laval Sr.¹, Rafel Cirer-Sastre²

1. Physiatry and nursing, University of Zaragoza, Zaragoza, ESP 2. Institut Nacional d'Educació Física de Catalunya, Universitat de Lleida, Lleida, ESP

Corresponding author: Isaac López-Laval Sr., isaac@unizar.es

Abstract

Conventional strength training and core exercises are commonly prescribed to improve cycling performance. Although previous studies have explored the utility of strength training in various cycling populations, this intervention has never been compared to core exercises. 36 trained road cyclists were divided in 3 groups of 12 participants that performed either no strength training, conventional strength training or core exercises, in all cases together with their regular cycling training during a 12-week period. Peak power outputs across different durations (5 seconds, 60 seconds, 5 minutes, and 20 minutes) were recorded before and after the intervention. The results of the present study showed higher increases in relative power output with conventional strength training when compared to core training and no strength training for all measured durations: 5-sec $\Delta = 1.25$ W/kg vs 0.47 W/kg and -0.17 W/kg; 60-sec ($\Delta = 0.51$ W/kg vs 0.13 W/kg and 0.02 W/kg; 5-min $\Delta = 0.22$ W/kg vs 0.06 W/kg and 0.05 W/kg and 20-min $\Delta = 0.22$ W/kg vs 0.07 W/kg and 0.06 W/kg. According to the data obtained in this study, conventional strength training is superior to core exercises and no strength training in trained road cyclists, and accordingly, it is recommended that this population incorporates strength training during their regular weekly workouts.

Categories: Sports Medicine

Keywords: strength, core training, endurance, bike, power output

Introduction

Road cycling is an endurance sport characterized by large training volumes and prolonged periods of moderate force production [1,2]. However, during decisive competition phases, short high intensity efforts are necessary in order to perform [3]. This complex and variable requirements in force production may be addressed on the bike but, as previous evidence pointed out, the addition of conventional strength training may result in even better results. Strength training may delay activation of less efficient type II fibers, improve neuromuscular efficiency, convert fast-twitch type IIX fibers into more fatigue-resistant type IIA fibers and improve muscle-tendinous stiffness [4]. On the other hand, core training may have the potential to increase joint range and muscle extensibility; improve joint stability, enhance muscle performance, and optimize movement function [5].

The incorporation of strength training during the preparatory period of trained road cyclists has received increasing attention since the beginning of the century. Several authors have reported improvements in various markers of performance after the addition of conventional strength training increases in type IIA fiber proportions and decreases in type IIX; improvements in pedaling efficacy and cycling economy after fatigue; increases in peak power output, time trial performance and power output at fixed blood lactate concentrations are some of the highlighted changes [6-8]. However, a previous study has also reported no effect of strength training when added to cycling training [9]. Although the consensus is towards a positive effect of these types of interventions, some degree of discrepancy still exists in the scientific and coaching field regarding the overall effect of strength training on cycling performance.

As road cycling is characterized by large training volumes performed under fixed postures in which the stabilizers and compensatory muscles of the trunk are being activated, the selective training of these muscle groups has received increased attention in the last years [2,10]. The practical coaching field has implemented training sessions with several exercises targeting the core muscles (transverse abdominis, multifidus, internal and external obliques, erector spinae, diaphragm, pelvic floor muscles, and the rectus abdominis [6]. Training of these muscles is seen as pivotal for efficient biomechanical function to maximize force generation and minimize joint loads in other endurance sports as swimming, although the scientific evidence behind these interventions in cycling is lacking as of today [11,12].

Previous research that investigated the effect of strength training interventions on cycling performance evaluated laboratory parameters such as lactate profiles, VO₂ kinetics, muscle fiber composition and torque profiles [13]. Concretely, strength training interventions resulted in an earlier occurrence of peak torque

values, higher power outputs at fixed blood lactates, transitions towards slower twitch muscle fiber types and improvements in gross efficiency represented as the power to VO₂ relationship [14,15] Although these values are useful for the prediction of cycling performance, during the last years the attention has focused towards the power profile, which allows to accurately predict performance after the analysis of power outputs registered with a mobile power meter [3]. This method allows an accurate, flexible, and inexpensive longitudinal tracking of performance without resorting to the laboratory. Previous research has shown that 5-second, 60-second, 5-minute and 20-minute intervals can be used to track neuromuscular and glycolytic changes, maximal oxygen consumption and maximal metabolic stable state, respectively [16-18].

Given the relative uncertainty regarding the true effect of conventional strength training on cycling performance, the lack of previous studies regarding the utility of core training and the easy implementation of power profiling as a valid tool to monitor changes in performance experienced after strength training, the objective of the current study was to compare the power profile of trained road cyclists after 12 weeks of a) conventional strength training; b) core training and c) no strength training during the preparatory phase of the annual training plan.

Materials And Methods

The current study was structured as an exploratory intervention aimed at assessing training effects. A test protocol, outlined below, was conducted both at baseline (pre-intervention) and after a 12-week intervention period (post-intervention). The study duration, spanning from November to January, encompassed the off-season and the initial phase of preparation leading up to the competitive season.

Participants

Thirty-six cyclists volunteered, with mean (standard deviation) [range] characteristics as follows: age = 28.8 (4.2) (21, 37) years, height = 179.3 (5.1) (169, 190) cm, body mass = 69.1 (4.5) (60, 80) kg. Random allocation divided participants into three groups of n = 12: cycling, cycling & core, and cycling & strength. All participants trained at the same facility, underwent assessments using identical equipment, and were evaluated by the same researcher. Inclusion criteria included possessing a cycling license (World Tour, Elite/U23, Masters, or recreational), absence of surgical procedures or injuries in the preceding 6 months, and no use of performance-enhancing supplements or drugs during the same period. Following informed consent and completion of a health-screening questionnaire, participants were enrolled in the study, adhering to ethical guidelines outlined in the 2013 Declaration of Helsinki and approved by the Research Ethics Committee of the autonomous region of Aragon, Spain (PI23/131) [19]. Grouped participant characteristics are presented in Table 1.

Variable	Cycling (n=12)	Cycling & Core (n=12)	Cycling & Strength (n=12)
Age (years)	30 (4) (22, 36)	29 (5) (21, 37)	28 (4) (22, 36)
Body height (cm)	180 (6) (170, 190)	179 (5) (169, 187)	179 (5) (170, 185)
Body mass (kg)	69 (6) (60, 80)	69 (3) (62, 73)	70 (4) (60, 74)

TABLE 1: Summary of participant characteristics.

Note: Data are expressed as mean (standard deviation) (range)

Procedures

Subjects underwent anthropometric evaluation and cycling tests before and after the 12-week intervention period, conducted within the morning hours (between 10:00 AM and 12:00 PM) to mitigate diurnal hormonal fluctuations. Data collection occurred under consistent environmental conditions (17-18°C, 45-55% relative humidity). Cycling tests were performed on personal bikes mounted on a Tacx Neo 2T Smart bike trainer (Tacx International, Rijksstraatweg, the Netherlands), with power output measured using Favero Assioma pedals [16]. Maximal oxygen consumption was estimated using a formula based on relative power output obtained during a 5-minute interval [20].

In order to obtain the power profile of each participant, subjects performed an adaptation of the testing protocol suggested by Allen & Coggan, as follows: (a) 20 minutes at a self-selected easy intensity, (b) 3 1-minute fast pedaling accelerations (100-105 rpm) with a 1-minute recovery between efforts, (c) 5 minutes at a self-selected easy intensity, (d) 1 all out 5 second sprint, (e) 5 minutes at a self-selected easy intensity, (f) 1-minute all-out effort, (g) 5-minute all-out effort, (h) 10 minutes at a self-selected easy intensity and 5 minutes of resting [2]. The main part of the test consisted of a

20-minute maximal effort, where subjects were asked to produce the highest mean PO possible for this duration and adopt their personal pacing strategies [2,21-23]. Participants were familiarized with the protocol as it commonly included in their normal testing routine. They could also view their progress on a computer monitor and were provided with information regarding time to completion and gear choice. All other information was blinded, no verbal encouragement was provided, and water was allowed ad libitum. FTP was determined as 95% of the mean power output of the 20-minute effort. Maximal power outputs for each selected duration were registered in relative values (w/kg) considering the weight obtained during the anthropometric evaluation.

Anthropometric evaluation

Body mass and fat mass were assessed using the electrical impedance method (BC-602; Tanita Co., Tokyo, Japan) in the morning, while height was measured using a SECA 214 stadiometer (Seca; Hamburg, Germany), graduated up to 1 mm.

Endurance training

All participants performed the same training sessions during the 12 weeks of the intervention period. Training was prescribed based on six power zones relative to the functional threshold power (FTP), which was calculated by subtracting 5% to the maximal 20-minute power output obtained in the baseline testing [2]. The power zones were set up as follows: <55% Zone 1; 56-75% Zone 2; 76-90% Zone 3; 91-105% Zone 4; 106-120% Zone 5 and >121% Zone 6 [2]. Participants performed 4 weekly sessions riding in Zone 2 (2h on Tuesday, 2h on Thursday, 4h on Saturday and 4h on Sunday). Compliance was verified by analyzing files obtained from the bike computer of the participant. 54 out of 56 participants (94%) achieved a compliance of >90% when Training Stress Score of the full intervention period was assessed.

Strength training

The strength training exercises were based on previous research and in the following order: (half squat, leg press with one leg at a time, one-legged hip flexion, and ankle plantar flexion) and were performed twice weekly (Monday and Wednesday) [24]. 3-minute rests were allowed between sets. All cyclists were supervised by an investigator at all workouts during the first 2 weeks and thereafter at least once every second week throughout the intervention period. During the 12 weeks of the intervention period, cyclists trained with 6 repetitions maximum (RM) sets until failure. The cyclists were encouraged to increase their RM loads after 4 and 8 weeks of the intervention period and they were allowed assistance on the last repetition. The number of sets in each exercise was always three. Over the entire training period, one session was not performed due to illness.

Core training

The core exercises (glute bridge, abdominal plank and prone back extension) were performed as recommended in a previous study designed with cyclists [25]. The glute bridge and prone back extension consisted of a 2-second concentric phase, 2-second isometric phase and 4-second eccentric phase. Both exercises incorporated 10 repetitions for each set. The abdominal plank was maintained for 30 seconds. 8 sets of each exercise were performed with a 60-second rest in-between. These exercises were performed twice weekly (Monday and Wednesday). All cyclists were supervised by an investigator at all workouts during the first 2 weeks and thereafter at least once every second week throughout the intervention period. Over the entire training period, two sessions were not performed due to illness.

Statistical analyses

Data are described as mean (standard deviation) (range). Baseline differences between the three groups were assessed by univariate analyses of variance (ANOVA). To compare the effectiveness of each intervention, a one-way between-groups analysis of covariance (ANCOVA) was conducted. The base analysis model was built with group allocation as the independent variable and the absolute change (Δ = post-pre) as the dependent variable, with the corresponding baseline scores as covariates. Prior to the analysis, the assumptions of linearity, homogeneity of regression slopes, normality of residuals, homogeneity of variances and absence of outlier values were inspected and, accordingly, the following decisions were made: 5-min RPO violated slope homogeneity and was heteroscedastic and, consequently, its modeling included an interaction term with group-by-baseline data and the White-Huber heteroscedasticity correction; and 60-sec RPO had baseline differences and, consequently, its modeling included an interaction term with group-by-baseline data. Post-hoc pairwise comparisons were based on model estimated marginal means and reported as estimated means and/or mean changes (Δ) and their 95% confidence interval (CI). Effect sizes for ANCOVA terms were reported as partial eta squared (η^2), and for pairwise comparisons as Cohen's d , in both cases with their respective 95% CI. Analyses were performed using R version 4.2.2 (R Core Team) and statistical significance was assumed when $p < 0.05$.

Results

Anthropometric data

At baseline, all groups had similar Body Mass ($F(2, 33) = 0.06, p = 0.94$), Fat Mass ($F(2, 33) = 3.07, p = 0.06$), Body Mass Index ($F(2, 33) = 1.21, p = 0.31$), and VO2max ($F(2, 33) = 1.11, p = 0.34$). Changes after training were also comparable among groups in Body Mass ($F(2, 32) = 0.12, p = 0.89$), Fat Mass ($F(2, 30) = 0.94, p = 0.402$), and Body Mass Index ($F(2, 32) = 0.026, p = 0.97$). By contrast, VO2max improvements were different among groups ($F(2, 32) = 6.84, p = 0.003; h2P = 0.30, 95\% CI (0.08, 1.00)$), being higher in the Cycling & Strength group ($\Delta = 2.14 \text{ ml/min/kg}, \Delta 95\% CI (1.4, 2.47) \text{ ml/min/kg}$) compared with the Cycling-only ($\Delta = 0.35 \text{ ml/min/kg}, \Delta 95\% CI (-0.27, 1.21) \text{ ml/min/kg}, t(32) = 3.22, p = 0.008, d = 1.14, d 95\% CI (0.38, 1.88)$) and the Cycling & Core group ($\Delta = 0.36 \text{ ml/min/kg}, \Delta 95\% CI (-0.2, 1.25) \text{ ml/min/kg}, t(32) = 3.2, p = 0.008, d = 1.13, d 95\% CI (0.38, 1.87)$).

Main relative power output differences

Individual relative power output (RPO) is summarized in Table 2 and individual trends can be inspected in Figure 1 in Figure 1) There were no group differences in baseline 5-sec RPO ($F(2, 33) = 1.39, p = 0.263$), 5-min RPO ($F(2, 33) = 1.11, p = 0.34$), and 20-min RPO ($F(2, 33) = 0.27, p = 0.763$), but a statistically significant and large main effect of group in 60-sec RPO ($F(2, 33) = 4.16, p = 0.024; h2P = 0.20, 95\% CI (0.02, 1.00)$). Adjusted by baseline data, RPO improvements after the intervention were different among groups in all tests: 5-sec RPO ($F(2, 32) = 14.09, p < .001; h2P = 0.47, 95\% CI (0.24, 1.00)$), 60-sec RPO ($F(2, 30) = 11.96, p < .001; h2P = 0.44, 95\% CI (0.20, 1.00)$), 5-min RPO ($F(2, 32) = 5.77, p = 0.008, h2P = 0.28, 95\% CI (0.06, 1.00)$), and 20-min RPO ($F(2, 32) = 11.72, p < .001; h2P = 0.42, 95\% CI (0.19, 1.00)$).

Variable	Cycling		Cycling & Core		Cycling & Strength		Adj. Group dif.	Adj. Time dif.
	Pre	Post	Pre	Post	Pre	Post		
5-sec RPO (W/kg)	14.97 (0.22)	14.82 (0.63)	14.93 (0.41)	15.44 (0.57)	15.12 (0.17)	16.31 (0.73)	F = 1.39, p = 0.263	F = 14.09, p < .001
60-sec RPO (W/kg)	8.98 (0.14)	9.04 (0.17)	9.31 (0.44)	9.38 (0.41)	9.07 (0.19)	9.61 (0.32)	F = 4.16, p = 0.024	F = 11.96, p < .001
5-min RPO (W/kg)	6 (0.12)	6.04 (0.14)	6.03 (0.09)	6.09 (0.18)	6.05 (0.06)	6.3 (0.19)	F = 1.11, p = 0.34	F = 5.77, p = 0.008
20-min RPO (W/kg)	5 (0.08)	5.08 (0.14)	4.98 (0.06)	5.03 (0.12)	4.99 (0.06)	5.21 (0.14)	F = 0.27, p = 0.763	F = 11.72, p < .001

TABLE 2: Participants performance in the Relative Power Output tests (RPO) by group and time.

Note: Data are expressed as mean (standard deviation)




FIGURE 1: Individual changes in Relative Power Output by test.

Post-hoc group contrasts by variable

Absolute mean changes are provided in Figure 2. Statistical contrasts revealed that adjusted by baseline, the mean 5-sec RPO improvements were higher in Cycling & Strength group ($\Delta = 1.25$ W/kg, Δ 95% CI (0.86, 1.64) W/kg) compared to the Cycling-only ($\Delta = -0.17$ W/kg, Δ 95% CI (-0.55, 0.21) W/kg, $t(32) = 5.3$, $p < 0.001$, $d = 1.88$, d 95% CI (1.04, 2.7)) and Cycling & Core ($\Delta = 0.47$ W/kg, Δ 95% CI (0.09, 0.85) W/kg, $t(32) = 2.86$, $p = 0.02$, $d = 1.01$, d 95% CI (0.27, 1.74)). Additionally, 5-min RPO was higher too in the Cycling & Core compared to Cycling-only ($t(32) = 2.46$, $p = 0.049$, $d = 0.87$, d 95% CI (1.14, 1.59)).

Improvements in 60-sec RPO were higher too in the Cycling & Strength group ($\Delta = 0.51$ W/kg, Δ 95% CI (0.34, 0.67) W/kg) compared to the Cycling-only ($\Delta = 0.02$ W/kg, Δ 95% CI (-0.21, 0.24) W/kg, $t(30) = 3.58$, $p = 0.003$, $d = 1.31$, d 95% CI (0.51, 2.09)) and Cycling & Core ($\Delta = 0.13$ W/kg, Δ 95% CI (-0.04, 0.3) W/kg, $t(30) = 3.22$, $p = 0.008$, $d = 1.18$, d 95% CI (0.39, 1.94)).

RPO in 5-min also improved more in the Cycling & Strength group ($\Delta = 0.22$ W/kg, Δ 95% CI (0.14, 0.32) W/kg) compared to the Cycling-only ($\Delta = 0.05$ W/kg, Δ 95% CI (-0.04, 0.14) W/kg, $t(30) = 2.9$, $p = 0.018$, $d = 1.06$, d 95% CI (0.29, 1.82)) and Cycling & Core ($\Delta = 0.06$ W/kg, Δ 95% CI (-0.02, 0.14) W/kg, $t(30) = 2.8$, $p = 0.023$, $d = 1.02$, d 95% CI (0.26, 1.78)).

Finally, the mean 20-min RPO improvement was greater in the Cycling & Strength group ($\Delta = 0.22$ W/kg, Δ 95% CI (0.17, 0.28) W/kg) compared to the Cycling-only ($\Delta = 0.07$ W/kg, Δ 95% CI (0.02, 0.13) W/kg, $t(32) = 3.98$, $p = 0.001$, $d = 1.41$, d 95% CI (0.62, 2.17)) and Cycling & Core ($\Delta = 0.06$ W/kg, Δ 95% CI (0.00, 0.11) W/kg, $t(32) = 4.37$, $p < 0.001$, $d = 1.54$, d 95% CI (0.75, 2.32)).




FIGURE 2: . Absolute mean changes in Relative Power Output by group and test.

Discussion

The objective of the current study was to compare the effects of conventional strength training, core training and no strength training in trained road cyclists. The main findings were as follows: a) Conventional strength training was superior to core training and no training for the improvement of 5-second, 60-second, 5-minute and 20-minute power outputs; b) Core training was superior to no training for the improvement of 5-second power outputs only and c) There were no differences in body composition changes across the three groups although the VO₂max increase was larger in the conventional strength training group.

To the best of the author's knowledge, this is the first study to compare conventional strength versus core training for the improvement of road cycling performance. The results support previous evidence that suggested the general utility of conventional strength exercises when contextualized into a concurrent strength and endurance training program [24,14,15]. This was not the case for core exercises, a lack of effect that suggests that these types of interventions do not benefit road cyclists. Several theoretical benefits have been suggested for core stability and strength training: increases in joint range and muscle extensibility; improvements in joint stability, enhanced muscle performance and optimized movement function are among the most reported [6]. Some of these possible benefits could be of interest for road cyclists given that most of them translate into better exercise economy [1,26]. Given that an improvement in exercise economy may not necessarily translate into an increase in mean maximal power outputs, the current study could have overlooked this hypothetical benefit, which should be explored in the future [7]. Despite this possibility, most cyclists have limited time to train and need to limit cycling time in order to incorporate some kind of strength training into their routine. Accordingly, the results of the current study suggest most cyclists would make a better use of their time by performing conventional strength exercises rather than core training.

Core training resulted in an increase in 5-second RPO when compared to no strength training. This finding should be further contextualized as the no strength training group actually decreased its sprinting power after the 12-week intervention period. This could be related to the fact that no sprints were performed during the study. Sprinting power isn't normally the main objective during the preparatory period of the annual training cycle. Given the time and effort required to perform the core training sessions, the cost/benefit ratio does not seem very attractive in this case and time of the season [10,18]. Theoretically, an improvement in core strength and stability could result in an optimized movement pattern during sprinting, which would result in an increase in power output during this race moment [10]. However, as of today, it is impossible to know whether an improvement in core strength and stability is the cause of this effect: there is lack of a gold standard method for measuring core stability and strength during sporting movements. Further, few studies have observed any performance enhancement in sporting activities despite observing improvements in core stability and core strength following a core training program [6]. Finally, there are no official guidelines nor scientific evidence regarding the best intensity, volume and distribution of core training sessions for the improvement of sporting performance. Accordingly, the intervention used in the current study, although based on previous research with cyclists, may have produced an insufficient stimulus to produce any measurable change in performance [27]. Given all the above, the results of the

current study do not support core training as a time-efficient strategy to improve road cycling performance.

No significant differences in the anthropometric characteristics were observed after 12 weeks of intervention. This finding is interesting given that previous research has reported increases in muscle mass, changes in fiber type and alterations of body mass after conventional strength training performed with heavy weights [1,17,24]. As the main interest of the present study was to assess the relative power output after the intervention period, changes in body mass and fat mass but not muscle mass were tracked. Therefore, the evolution of the specific components of body composition cannot be discussed in this case. Given the lack of difference in the evolution of body mass and the clear increase in VO₂max in the conventional strength training group, this change can be attributed to the increase in absolute power output observed in this group [25]. Improvements in either fractional or maximal oxygen utilization have been reported in previous research and are probably related to postponed activation of less efficient type II muscle fibers, conversion of type IIX fibers into more fatigue-resistant IIa fibers, and increased muscle mass and rate of force development [17,27]. Given these results, conventional strength training is an interesting addition to cycling training in order to improve VO₂max.

The current study presents several limitations that should be addressed in future research. First, for obvious reasons, participants couldn't be blinded and accordingly were exposed to the placebo effect. Concretely, past research shows that participants' belief in the efficacy of an intervention may influence findings in sport science research [28]. The conventional strength training group could have been exposed to this effect and, consequently, could have reported better results [29]. Second, the sample was conformed by participants of different cycling levels. The sample size was insufficient to compare across performance groups and try to extrapolate whether the findings obtained in the current study could be applied to all cyclist irrespective of their level. Finally, several important physiological parameters such as lactate or heart rate were not monitored in this study. This is important as improvements in relative blood lactate values or cardiac drift can be obtained even in the absence of increases in power output [30].

Conclusions

Road cyclists are often limited by their time availability in order to choose the best training plan for maximal performance optimization. The incorporation of strength training sessions into the weekly training routine may reduce the time spent on the bicycle and, accordingly, this decision should not be taken lightly. The present study suggests that 12 weeks of conventional strength training added to cycling during the preparatory phase of the season results in increased power outputs over the entire power curve and improves VO₂max. These results were superior to those obtained from core training, which only increased power output during sprints. Given the notable performance gains and great adherence observed in the current study, it is recommended that cyclists include bi-weekly strength training sessions into their pre-season. The exercises should target the main muscles activated during the pedal stroke (half squat, leg press with one leg at a time, one-legged hip flexion, and ankle plantar flexion) with intensities of 6RM and three sets per exercise.

Additional Information

Disclosures

Human subjects: Consent was obtained or waived by all participants in this study. Research Ethics Committee of the autonomous region of Aragon issued approval PI23/131. Dirección General de Aragon (DGA). **Animal subjects:** All authors have confirmed that this study did not involve animal subjects or tissue. **Conflicts of interest:** In compliance with the ICMJE uniform disclosure form, all authors declare the following: **Payment/services info:** All authors have declared that no financial support was received from any organization for the submitted work. **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

References

1. Aagaard P, Andersen JL, Bennekou M, et al.: Effects of resistance training on endurance capacity and muscle fiber composition in young top-level cyclists. *Scand J Med Sci Sports*. 2011, 21:298-307. [10.1111/j.1600-0858.2010.01283.x](https://doi.org/10.1111/j.1600-0858.2010.01283.x)
2. Allen, H.; Coggan, A: Training and racing with a power meter VeloPress . Boulder, CO; 2019.
3. Bell PG, Furber MJ: VAN Someren KA, Antón-Solanas A . Swart J. The physiological profile of a multiple tour de france winning cyclist. *Med Sci Sports Exerc*. 2017, 49:115-123. [10.1249/MSS.0000000000001068](https://doi.org/10.1249/MSS.0000000000001068)
4. Rønnestad BR, Mujika I: Optimizing strength training for running and cycling endurance performance: A review. *Scand J Med Sci Sports*. 2014, 24:605-612. [10.1111/sms.12104](https://doi.org/10.1111/sms.12104)
5. Hansen EA, Rønnestad BR, Vegge G, Raastad T: Cyclists' improvement of pedaling efficacy and performance after heavy strength training. *Int J Sports Physiol Perform*. 2012, 7:513-521. [10.1123/ijsp.7.4.513](https://doi.org/10.1123/ijsp.7.4.513)
6. Hibbs AE, Thompson KG, French D, Wrigley A, Spears I: Optimizing performance by improving core stability and core strength. *Sports Med*. 2008, 38:995-1008. [10.2165/00007256-200838120-00004](https://doi.org/10.2165/00007256-200838120-00004)
7. Leo P, Spragg J, Podlogar T, Lawley JS, Mujika I: Power profiling and the power-duration relationship in cycling: a narrative review. *Eur J Appl Physiol*. 2022, 122:301-316. [10.1007/s00421-021-04833-y](https://doi.org/10.1007/s00421-021-04833-y)

8. Levin GT, Mcguigan MR, Laursen PB: Effect of concurrent resistance and endurance training on physiologic and performance parameters of well-trained endurance cyclists. *J Strength Cond Res.* 2009, 23:2280-2286. [10.1519/JSC.0b013e3181b990c2](https://doi.org/10.1519/JSC.0b013e3181b990c2)
9. Lucia A, Hoyos J, Chicharro JL: Physiology of professional road cycling. *Sports Med.* 2001, 31:325-337. [10.2165/00007256-200131050-00004](https://doi.org/10.2165/00007256-200131050-00004)
10. Menaspà P, Quod M, Martin DT, Peiffer JJ, Abbiss CR: Physical demands of sprinting in professional road cycling. *Int J Sports Med.* 2015, 36:1058-1062. [10.1055/s-0035-1554697](https://doi.org/10.1055/s-0035-1554697)
11. Luo S, Soh KG, Soh KL, Sun H, Nasiruddin NJM, Du C, Zhai X: Effect of core training on skill performance among athletes: a systematic review. *Front Physiol.* 2022, 6:915259. [10.3389/fphys.2022.915259](https://doi.org/10.3389/fphys.2022.915259)
12. Kibler WB, Press J, Sciascia A: The role of core stability in athletic function. *Sports Med.* 2006, 36:189-198. [10.2165/00007256-200636030-00001](https://doi.org/10.2165/00007256-200636030-00001)
13. Castronovo AM, Conforto S, Schmid M, Bibbo D, D'Alessio T: How to assess performance in cycling: the multivariate nature of influencing factors and related indicators. *Front Physiol.* 2013, 21:116. [10.3389/fphys.2013.00116](https://doi.org/10.3389/fphys.2013.00116)
14. Rønnestad BR, Hansen J, Hollan I, Ellefsen S: Strength training improves performance and pedaling characteristics in elite cyclists. *Scand J Med Sci Sports.* 2015, 25:89-98. [10.1111/sms.12257](https://doi.org/10.1111/sms.12257)
15. Vikmoen O, Rønnestad BR: A Comparison of the effect of strength training on cycling performance between men and women. *J Funct Morphol Kinesiol.* 2021, 17:29. [10.3390/jfmk6010029](https://doi.org/10.3390/jfmk6010029)
16. Montalvo-Pérez A, Alejo LB, Valenzuela PL, et al.: Validity of the Favero Assioma duo Power Pedal System for measuring power output and cadence. *Sensors (Basel).* 2021, 24:2277. [10.3390/s21072277](https://doi.org/10.3390/s21072277)
17. Mujika I, Rønnestad BR, Martin DT: Effects of increased muscle strength and muscle mass on endurance-cycling performance. *Int J Sports Physiol Perform.* 2016, 11:283-9. [10.1123/IJSPP.2015-0405](https://doi.org/10.1123/IJSPP.2015-0405)
18. Navalta JW, Hrcir SP: Core stabilization exercises enhance lactate clearance following high-intensity exercise. *J Strength Cond Res.* 2007, 21:1305-1309. [10.1519/R-21546.1](https://doi.org/10.1519/R-21546.1)
19. Pallares JG, Hernández-Belmonte A, Valenzuela PL, Muriel X, Mateo-March M, Barranco-Gil D, Lucia A: Field-derived maximal power output in cycling: an accurate indicator of maximal performance capacity? . *Int J Sports Physiol Perform.* 2022, 27:1558-1564. [10.1123/ijsp.2022-0208](https://doi.org/10.1123/ijsp.2022-0208)
20. Rønnestad BR, Hansen EA, Raastad T: Effect of heavy strength training on thigh muscle cross-sectional area, performance determinants, and performance in well-trained cyclists. *Eur J Appl Physiol.* 2010, 108:965-975. [10.1007/s00421-009-1307-z](https://doi.org/10.1007/s00421-009-1307-z)
21. Sitko S, Cirer-Sastre R, Corbi F, López-Laval I: Functional threshold power as an alternative to lactate thresholds in road cycling. *J Strength Cond Res.* 2022, 1:3179-3183. [10.1519/JSC.0000000000004070](https://doi.org/10.1519/JSC.0000000000004070)
22. Sitko S, Cirer-Sastre R, Corbi F, López-Laval I: Five-minute power-based test to predict maximal oxygen consumption in road cycling. *Int J Sports Physiol Perform.* 2022, 1:9-15. [10.1123/ijsp.2020-0923](https://doi.org/10.1123/ijsp.2020-0923)
23. Sitko S, Cirer-Sastre R, Corbi F, López-Laval I: Power assessment in road cycling: a narrative review. *Sustainability.* 2020, 12:5216. [10.3390/su12125216](https://doi.org/10.3390/su12125216)
24. Rønnestad BR, Mujika I: Optimizing strength training for running and cycling endurance performance: A review. *Scand J Med Sci Sports.* 2014, 24:603-612. [10.1111/sms.1210](https://doi.org/10.1111/sms.1210)
25. Sunde A, Støren O, Bjerkaas M, Larsen MH, Hoff J, Helgerud J: Maximal strength training improves cycling economy in competitive cyclists. *J Strength Cond Res.* 2010, 24:2157-2165. [10.1519/JSC.0b013e3181aeb16a](https://doi.org/10.1519/JSC.0b013e3181aeb16a)
26. Vikmoen O, Ellefsen S, Trøen Ø, Hollan I, Hanestadhaugen M, Raastad T, Rønnestad BR: Strength training improves cycling performance, fractional utilization of VO₂max and cycling economy in female cyclists. *Scand J Med Sci Sports.* 2016, 26:384-396. [10.1111/sms.12468](https://doi.org/10.1111/sms.12468)
27. Warburton DER, Jamnik V, Bredin SSD, Shephard RJ, Gledhill N: The 2019 Physical Activity Readiness Questionnaire for Everyone (PAR-Q+) and electronic Physical Activity Readiness Medical Examination (ePARmed-X+): 2019 PAR-Q+. *Heal Fit J Can.* 2018, 11:80-83. [10.14288/hfjc.v11i4.27](https://doi.org/10.14288/hfjc.v11i4.27)
28. Beedie CJ, Coleman DA, Foad AJ: Positive and negative placebo effects resulting from the deceptive administration of an ergogenic aid. *Int J Sport Nutr Exerc Metab.* 2007, 17:259-269. [10.1123/ijsnem.17.3.259](https://doi.org/10.1123/ijsnem.17.3.259)
29. Eng J: Sample size estimation: how many individuals should be studied? . *Radiology.* 2003, 227:309-313. [10.1148/radiol.2272012051](https://doi.org/10.1148/radiol.2272012051)
30. Formenti D, Trecroci A, Cavaggioni L, Caumo A, Alberti G: Heart rate response to a marathon cross-country skiing race: A case study. *Sport Scie Heal.* 2015, 11:125-128. [10.1007/s11532-014-0187-8](https://doi.org/10.1007/s11532-014-0187-8)