


Are perceived barriers to physical activity related to depression, anxiety and stress among adolescents? The EHDLA study

Edina Maria de Camargo,¹ Sitong Chen,² Rubén López-Bueno,³ Arthur Eumann Mesas,⁴ Bruno Bizzozero-Peroni,^{4,5} Nerea Martín-Calvo,^{6,7,8} Estela Jiménez-López,^{4,9} José Francisco López-Gil ¹⁰

To cite: de Camargo EM, Chen S, López-Bueno R, et al. Are perceived barriers to physical activity related to depression, anxiety and stress among adolescents? The EHDLA study. *BMJ Open Sport & Exercise Medicine* 2024;**10**:e002069. doi:10.1136/bmjsem-2024-002069

► Additional supplemental material is published online only. To view, please visit the journal online (<https://doi.org/10.1136/bmjsem-2024-002069>).

Accepted 1 September 2024



© Author(s) (or their employer(s)) 2024. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr José Francisco López-Gil; josefranciscolepezgil@gmail.com

ABSTRACT

Objective The aim of this study was twofold: first, to examine the association between perceived barriers to physical activity (PA) practice and depression, anxiety and stress in a sample of Spanish adolescents; and second, to determine which barriers are specifically associated with depression, anxiety and stress.

Methods This cross-sectional study was conducted with 765 adolescents aged 12–17 (55.6% girls) in the *Valle of Ricote*, Murcia, Spain. Depression, anxiety and stress symptoms were assessed using the Depression, Anxiety and Stress Scale (DASS-21), with validated cut points employed to determine the presence of each of these mental conditions. The perception of barriers to PA was assessed using a validated questionnaire for the Spanish adolescent population.

Results The barrier 'Because I feel that my physical appearance is worse than that of others' was related to a higher likelihood of having depression (OR=2.41; 95% CI 1.35 to 4.28; p=0.003), anxiety (OR=2.65; 95% CI 1.51 to 4.71; p=0.001) and stress (OR=2.82; 95% CI 1.59 to 5.07; p<0.001). Similarly, the barrier 'Because nobody encourages me to engage in physical activity' was related to a higher likelihood of having depression (OR=1.92; 95% CI 1.08 to 3.43; p=0.026), anxiety (OR=1.97; 95% CI 1.11 to 3.50; p=0.021) and stress (OR=1.99; 95% CI 1.12 to 3.59; p=0.021).

Conclusion Perceived barriers to PA related to physical appearance and social support seem to be associated with a greater likelihood of depression, anxiety and stress among Spanish adolescents.

INTRODUCTION

According to the WHO,¹ 970 million people worldwide had a mental disorder in 2019.² Depression and anxiety are the most prevalent mental disorders and main causes of disability, being considered among the top 10 causes of disability-adjusted life-years in youths aged 10–24 years.³ Thus, the presence of mental health symptoms in adolescence, even when subclinical, predicts psychopathology in the future and can contribute to increased

WHAT IS ALREADY KNOWN ON THIS TOPIC

- ⇒ The relationship between perceived barriers to engaging in physical activity (PA) and the presence of depression, anxiety and stress has not been investigated in the adolescent population.
- ⇒ Gaining insight into the factors related to PA engagement and mental health might aid the design of more effective interventions.

WHAT THIS STUDY ADDS

- ⇒ Perceiving barriers related to physical appearance and perceiving a lack of social support seem to be associated with symptoms of depression, anxiety and stress in adolescents.
- ⇒ Promoting PA by reducing some specific barriers could be an important step in reducing symptoms of depression, anxiety and stress in adolescents.

morbidity, mortality and disability.^{4–6} Nevertheless, half of mental disorders develop before the age of 18.⁶ This leads to a strong call for interventions aimed at promoting mental health in children and adolescents.^{7,8}

Strategies focused on the improvement of some modifiable lifestyle factors, such as the practice of physical activity (PA),⁹ have been proposed as potential contributors to the prevention or decrease of depression, anxiety and stress symptoms.¹⁰ The importance of PA has been highlighted worldwide, as it provides multiple benefits related to health in children and adolescents, including their mental health.¹¹ The lifestyle acquired during childhood and adolescence can determine their future health and possibilities.¹² Investing in the health and well-being of adolescents not only yields current benefits but also has long-term advantages for the upcoming decades and the next generation.¹² Numerous studies indicate that active individuals during childhood and adolescence are likely to maintain an active lifestyle in adulthood, and active

adults, in turn, tend to encourage PA in their children.¹³ However, approximately 81% of adolescents aged 11–17 years worldwide accumulate insufficient PA, with significant differences in prevalence according to sex, region and country.¹⁴

Different factors are thought to influence engagement in PA during adolescence. These factors encompass the existence of perceived barriers, an element that, according to an individual's perception, poses challenges in adopting a physically active lifestyle, either individually¹⁵ or in combination.^{16–17} In this line, systematic reviews have indicated an inverse association between the perception and number of certain barriers, such as the lack of social support or the self-perceived physical appearance, and levels of practised PA.^{18–20} Additionally, low PA levels are considered a serious threat to the health and well-being of the population^{3 21–23} and have been related to symptoms of depression, anxiety and stress in adolescents.^{7–9}

The development and persistence of mental health issues are considered multifactorial, involving complex interactions between genetic and environmental factors.²⁴ Furthermore, different variables, including comorbidity and social or lifestyle factors, have been examined as potential protective or risk factors for mental health problems.²⁵ Scientific evidence indicates that participation in PA can support the current and future mental health of young people.^{26–27} For example, findings from a systematic review and meta-analysis by Rodriguez-Ayllon *et al*²⁶ suggest that PA interventions can enhance adolescents' mental health. Furthermore, observational studies indicate that promoting PA and reducing sedentary behaviour may protect adolescents' mental health. Addressing the reduction of perceived barriers to practising PA is an important consideration alongside promoting other healthy behaviours, such as improving the quality of sleep, all of which can promote the reduction of depressive, anxiety and stress symptoms.^{7–9 11} Therefore, gaining insight into the factors related to PA engagement and mental health might aid the design of more effective interventions. Nevertheless, the relationship between perceived barriers to engaging in PA and the presence of depression, anxiety and stress has not been investigated in the adolescent population.

Despite extensive research on the benefits of PA on mental health,²⁶ there is limited knowledge about how specific perceived barriers are associated with mental health outcomes in this population. Therefore, there is a need to understand the relationship between perceived barriers to PA and mental health issues in adolescents. This study is novel as it is the first to investigate these associations in a sample of adolescents, providing insights that could contribute to the development of strategies aimed at promoting PA by mitigating barriers, thereby lowering the levels of depressive, anxiety and stress symptoms within this population. It is hypothesised that a higher number of perceived barriers to PA, such as concerns about physical appearance and lack of social support,

are positively associated with higher symptoms of depression, anxiety and stress in adolescents, independent of their PA levels. Therefore, the aim of the present study was twofold: first, to examine the association between perceived barriers to PA practice and depression, anxiety and stress in a sample of Spanish adolescents, and second, to determine which barriers are specifically associated with depression, anxiety and stress

MATERIAL AND METHODS

Population and study design

This cross-sectional study includes data from the Eating Healthy and Daily Life Activities (EHDLA) study, where adolescents from three secondary schools from the *Valle de Ricote* (Region of Murcia, Spain) were assessed. The methodology of the EHDLA study has been previously published.²⁸ Regarding participation, the parents or legal guardians of the adolescents were provided with a written informed consent form to be signed previously. The adolescents were also asked about their willingness to participate in the study. The entire data collection process was carried out during physical education classes. The inclusion criteria for the study were as follows: aged between 12 and 17 years old and registered and/or lived in the *Valle de Ricote*. Regarding exclusion criteria, adolescents are exempt from physical education at school and pathology that contraindicates PA or demands special attention and pharmacological treatment.

Procedures

Perceived barriers to PA practice (independent variables)

A scale of perceived barriers to PA in adolescents, previously validated among the Spanish population, was applied²⁹ to assess perceived barriers to PA in adolescents. The scale is composed of twelve items that are included in four distinct dimensions: barriers due to incompatibility with other tasks (two items), self-concept barriers (four items), motivational barriers (four items) and social barriers (two items). Each item is scored with a 5-point Likert-type response scale, from 1 (strongly disagree) to 5 (strongly agree).²⁹ For further analyses, we categorised the items as 'agree' (4 or 5 points on a Likert scale) or 'not agree' (1, 2 and 3 points on a Likert scale). In addition, we determined the number of perceived barriers to PA (ie, items with 'agree' responses).

Depression, anxiety and stress (dependent variables)

Depressive, anxiety and stress symptoms were evaluated by the *Depression, Anxiety and Stress Scale* (DASS-21),³⁰ which includes 21 items scored on a 4-point Likert-type scale. The score of each item ranges from 0 (did not apply to me at all) to 3 (applied to me very much or most of the time) and is divided uniformly into three subscales: depression, anxiety and stress. The depression subscale assesses symptoms like inertia, anhedonia, dysphoria, lack of interest/involvement, self-depreciation, devaluation of life and discouragement. The anxiety subscale evaluates the excitation of the autonomous nervous

system, musculoskeletal effects, situational anxiety and subjective anxiety experiences. Finally, the stress subscale assesses difficulty in relaxing, such as nervous excitation, easy perturbation/agitation, irritability/exaggerated reaction and impatience. The Spanish version, which has adequate reliability, was used in this study.³¹ The cut-off points established for the presence of depression, anxiety and stress were ≥ 6 points, ≥ 6 points and ≥ 5 points, respectively, due to the adequate sensitivity and specificity that these cut-off points have shown in adolescents.³² It is important to mention that higher scores represent more severe symptoms on the DASS-21.

Covariates

Sex and age were self-reported. Socioeconomic status (SES) was assessed using the Family Affluence Scale-III,³³ which has a final score that ranges from 0 to 13 points. The body weight of the adolescents was measured using an electronic scale (with an accuracy of 0.1 kg) (Tanita BC-545, Tokyo, Japan), while a portable height rod determined height with an accuracy of 0.1 cm (Leicester Tanita HR 001, Tokyo, Japan). The body mass index (BMI) was calculated by dividing body weight (in kg) and height (in square metres). PA and sedentary behaviour were assessed using the Spanish version of the Youth Activity Profile.³⁴ The Youth Activity Profile is a 15-item self-administered 7-day (previous week) recall questionnaire appropriate for young people aged 8 and 17 years.³⁵ The items use a 5-point Likert scale and are separated into three sections: activity at school, activity out of school and sedentary habits.³⁶ Sleep duration was evaluated by asking respondents for weekdays and weekends separately: 'What time does your child usually go to bed?' and 'What time does your child usually get up?'. The average daily sleep duration was calculated for each adolescent, considering the average nighttime sleep duration on weekdays and weekends.³⁶

Statistical analysis

For categorical variables, the descriptive data are presented as the number (n) and the percentage (%) of observations. For continuous variables, descriptive data were presented as median and IQR values since visual methods (ie, density and quantile-quantile plots) and the Shapiro-Wilk test showed a non-normal distribution of the variables. To assess the potential moderating effect of sex on the relationship between barriers to PA and depression, anxiety or stress, an interaction term was included in the generalised logistic regression model. The results indicated a non-significant interaction between barriers to PA and sex (depression: $p=0.397$; anxiety: $p=0.757$; stress: $p=0.826$), suggesting that the association of barriers with depression, anxiety or stress do not differ between sexes. Therefore, both sexes were examined jointly. To examine the association between the number of perceived barriers for PA and the number of depression, anxiety, or stress symptoms among adolescents without any parametric assumptions on the nature

of the relationship, generalised additive models (GAMs) were applied. The GAM is a flexible model that can capture non-linear connections in data without requiring a predefined mathematical structure. GAMs were chosen for their flexibility in capturing complex non-linear relationships through smooth functions, which adaptively fit the data without requiring prespecified knot placement. Restricted maximum likelihood for smoothness selection was applied,³⁷ with a shrinkage approach employed as a function of thin plate regression spline smoothers.³⁸ To quantify the degree of non-linearity of the curve, we used the effective df (edf) of the GAM. To determine the OR and its 95% CI of the association of barriers to PA status (ie, low, medium and high) or each specific barrier to PA with depression, anxiety or stress status, we carried out binary logistic regression analyses. Furthermore, we applied a correction for multiple comparisons using the false discovery rate p value method developed by Benjamini and Hochberg.³⁹ In addition, we computed prediction margins and average marginal effects of the relationship between barriers to PA (ie, for each additional barrier, based on the status of barriers) and levels of depression, anxiety and stress (ie, for each additional symptom, based on the status of depression, anxiety and stress). Models were adjusted for sex, age, SES, BMI, overall sleep duration, PA and sedentary behaviour. All statistical analyses were conducted with the statistical software R (V.4.3.2) (R Core Team, Vienna, Austria) and RStudio (2023.09.1+494) (Posit, Boston, MA, USA). Statistical significance was set up with a p value <0.05 .

RESULTS

Table 1 shows the main characteristics of the study participants. A total of 31.2% of adolescents were classified as having depression, 34.9% as having anxiety and 40.3% as having stress. Online supplemental table S1 shows the perceived barriers to the practice of PA in adolescents. The barriers most declared by adolescents were 'Because I have a lot of homework' (42.6%) and 'Because I am afraid of making a fool of myself' (19.1%). The least frequently declared barriers were 'Because I consider that I already do enough physical activity in my physical education classes' (7.8%) and 'Because nobody encourages me to engage in physical activity' (9.9%).

Figure 1 presents smoothed functions from GAMs for the number of barriers for PA and as a function of depression, anxiety or stress score. Examining the figure, edf and approximate significance of the smooth terms, we observed a nonlinear relationship between the number of barriers to PA and depressive symptoms ($F=5.43$; $edf=1.89$; $p<0.001$), anxiety symptoms ($F=5.08$; $edf=2.28$; $p<0.001$) and stress symptoms ($F=4.42$; $edf=2.29$; $p<0.001$). The predictive margins for each additional symptom of depression, anxiety or stress are shown in online supplemental figure S1. For each additional barrier to PA, the probability of having one more symptom of depression, anxiety and stress was 61.8% (95% CI 39.9% to 83.7%,

Table 1 Descriptive data of the study participants (n=765)

Variables		Perceived barriers to physical activity		
		Low n=548; 71.6%	Medium n=171; 22.4%	High n=46; 6.0%
Age (years)	Mean (SD)	13.9 (1.5)	14.3 (1.6)	14.0 (1.5)
Sex	Boys, n (%)	273 (49.8)	50 (29.2)	17 (37.0)
	Girls, n (%)	275 (50.2)	121 (70.8)	29 (63.0)
Socioeconomic status (FAS-III score)	Mean (SD)	8.2 (2.1)	8.0 (2.0)	7.7 (2.4)
Physical activity (YAP-S score)	Mean (SD)	2.7 (0.7)	2.5 (0.6)	2.6 (0.8)
Sedentary behaviours (YAP-S score)	Mean (SD)	2.6 (0.6)	2.6 (0.6)	2.9 (0.7)
Overall sleep duration (minutes)	Mean (SD)	497.3 (53.4)	481.3 (58.8)	476.5 (64.0)
Body mass index (kg/m ²)	Mean (SD)	22.5 (4.7)	23.5 (5.1)	24.6 (5.3)
Excess weight status	No, n (%)	417 (76.1)	121 (70.8)	29 (63.0)
	Yes, n (%)	131 (23.9)	50 (29.2)	17 (37.0)
DASS-21				
Depression symptoms (score)	Mean (SD)	3.7 (4.5)	6.7 (6.4)	7.4 (6.2)
Depression (status)*	No, n (%)	410 (74.8)	94 (55.0)	22 (47.8)
	Yes, n (%)	138 (25.2)	77 (45.0)	24 (52.2)
Anxiety symptoms (score)	Mean (SD)	3.4 (4.0)	6.2 (5.6)	6.5 (5.4)
Anxiety (status)†	No, n (%)	394 (71.9)	83 (48.5)	21 (45.7)
	Yes, n (%)	154 (28.1)	88 (51.5)	25 (54.3)
Stress symptoms (score)	Mean (SD)	4.5 (4.4)	7.4 (5.4)	7.1 (5.2)
Stress (status)‡	No, n (%)	371 (67.7)	69 (40.4)	17 (37.0)
	Yes, n (%)	177 (32.3)	102 (59.6)	29 (63.0)

Data are expressed as the mean (SD) or count (percentages).
 *Cut-off point for depression symptoms ≥ 6 points.
 †Cut-off point for anxiety symptoms ≥ 6 points.
 ‡Cut-off point for stress symptoms ≥ 5 points.
 DASS-21, Depression, Anxiety and Stress Scale-21; FAS-III, Family Affluence Scale-III; YAP-S, Spanish Youth Active Profile.

$p < 0.001$), 52.7% (95% CI 31.0% to 74.5%, $p < 0.001$) and 56.2% (95% CI 33.8% to 78.6%, $p < 0.001$), respectively.

Figure 2 displays the adjusted ORs of having depression, anxiety and stress in adolescents according to their perceived barriers to PA status. Compared with those with low perceived barriers to PA, adolescents with medium or high barriers to PA had higher probabilities of having depression (medium perceived barriers to PA: OR=1.97; 95% CI 1.35 to 2.88; $p < 0.001$; high perceived barriers to PA: OR=2.62; 95% CI 1.36 to 5.05; $p = 0.004$), anxiety (medium perceived barriers to PA: OR=2.29; 95% CI 1.58 to 3.32; $p < 0.001$; high perceived barriers to PA: OR=2.55; 95% CI 1.34 to 4.89; $p = 0.004$) and stress (medium perceived barriers to PA: OR=2.62; 95% CI 1.81 to 3.82; $p < 0.001$; high perceived barriers to PA: OR=3.31; 95% CI 1.72 to 6.52; $p < 0.001$). Additionally, online supplemental figure S2 displays the predictive margins for having depression, anxiety or stress for each additional perceived barrier to PA. Thus, the probability of having depression, anxiety and stress was 3.2% (95% CI 2.0% to 4.5%, $p < 0.001$), 3.7% (95% CI 2.4% to 5.1%, $p < 0.001$) and 4.1% (95% CI 2.7% to 5.4%, $p < 0.001$), respectively.

Table 2 shows the results of the logistic regression analyses examining the relationship between perceived barriers to PA and symptoms of depression, anxiety and stress in adolescents before and after adjusting for age, sex, SES, BMI, PA, sedentary behaviour and overall sleep duration. Barrier 8, 'Because I feel that my physical appearance is worse than that of others', was related to a higher likelihood of having depression (OR=2.41; 95% CI 1.35 to 4.28; $p = 0.003$), anxiety (OR=2.65; 95% CI 1.51 to 4.71; $p = 0.001$) and stress symptoms (OR=2.82; 95% CI 1.59 to 5.07; $p < 0.001$). Similarly, barrier 9, 'Because nobody encourages me to engage in physical activity', was related to a higher likelihood of having depression (OR=1.92; 95% CI 1.08 to 3.43; $p = 0.026$), anxiety (OR=1.97; 95% CI 1.11 to 3.50; $p = 0.021$) and stress symptoms (OR=1.99; 95% CI 1.12 to 3.59; $p = 0.021$).

DISCUSSION

Main findings and comparison with previous studies

Overall, the present study is the first to examine the association between the perception of barriers to PA practice and depression, anxiety and stress symptoms. Compared






Figure 1 Curvilinear associations between the number of barriers to physical activity (PA) and the number of depression, anxiety or stress symptoms among adolescents. Adjusted for age, sex, socioeconomic status, body mass index, PA, sedentary behaviour and overall sleep duration.

Depression status: OR (95% CI, *p*-value)

Low perceived barriers to PA	Reference
Medium perceived barriers to PA	1.97 (1.35–2.88, <i>p</i> < 0.001)
High perceived barriers to PA	2.62 (1.36–5.05, <i>p</i> = 0.004)

Anxiety status: OR (95% CI, *p*-value)

Low perceived barriers to PA	Reference
Medium perceived barriers to PA	2.29 (1.58–3.32, <i>p</i> < 0.001)
High perceived barriers to PA	2.55 (1.34–4.89, <i>p</i> = 0.004)

Stress status: OR (95% CI, *p*-value)

Low perceived barriers to PA	Reference
Medium perceived barriers to PA	2.62 (1.81–3.82, <i>p</i> < 0.001)
High perceived barriers to PA	3.31 (1.72–6.52, <i>p</i> < 0.001)




Figure 2 Adjusted ORs and 95% CIs of having depression, anxiety and stress symptoms according to barriers to physical activity (PA) status among adolescents. Adjusted for age, sex, socioeconomic status, body mass index, PA, sedentary behaviour and overall sleep duration.

with adolescents with low perceived barriers to PA (ie, from 0 to 1 barrier), those with two more barriers showed a greater number of depression, anxiety and stress symptoms, as well as a higher likelihood of having depression, anxiety and stress. Specifically, the barriers related to physical appearance and lack of encouragement were associated with higher odds of depression, anxiety and stress after adjusting for several covariates (ie, sociodemographic, socioeconomic, anthropometric and lifestyle variables). Previous studies have identified associations between PA and physical appearance,^{40–42} as well as lack of encouragement,^{43 44} which may support the present study's findings. For instance, regarding barriers related to physical appearance, the systematic review by Gualdi-Russo *et al*⁴⁰ showed that PA involvement is related to lower concerns about body image perception and enhances body satisfaction (ie, as PA increases, body image dissatisfaction decreases). In relation to barriers linked to lack of encouragement, the systematic review by Mendonça *et al*⁴³ indicated that participants who received more overall social support and support from parents, friends and family exhibited higher levels of PA. Furthermore, a study with Brazilian adolescents found that social support from parents and friends was associated with higher moderate to vigorous PA levels. Several factors could explain our findings, such as the relationship between self-perception of physical appearance and mental health symptoms like depression, anxiety and stress; the impact

of social support on these mental health issues; and the influence of PA on depression, anxiety and stress.

First, adolescents who do not conform to societal beauty standards or have different body shapes and sizes (ie, obesity) may experience stigma and discrimination, which can be detrimental to their mental well-being.^{42 45} In the same line, adolescents who are dissatisfied with their physical appearance may experience negative body image self-perception, low self-esteem and lower body confidence.⁴⁰ On the other hand, comparing themselves to others who may appear to be more physically fit or athletic may further contribute to negative emotions. These feelings and the fear of judgement or scrutiny from others can lead to reluctance to engage in PA,^{42 45} especially in public settings.⁴⁰ Similarly, negative comments, bullying or feeling judged due to their appearance can significantly impact self-esteem, self-worth and overall psychological health.⁴¹ This, in turn, can create a barrier to engaging in PA, as it may be associated with potential exposure to such negative experiences.¹⁹ A systematic review reported a significantly elevated prevalence of depression and anxiety symptoms in children/adolescents with overweight/obesity in contrast to those without.⁴² Given the association between inadequate PA and an energy imbalance, it is reasonable to suggest that weight gain and its potential adverse effects on mental health might serve as a mediator in the association between barriers to PA and heightened symptoms of

Table 2 ORs and 95% CIs of having depression, anxiety and stress symptoms according to specific barriers to physical activity among adolescents (N=765)

Barriers	Depression	
	OR (95% CI, p value) (univariable)	OR (95% CI, p value) (multivariable)
Because I have a lot of homework	1.22 (0.90 to 1.67, p=0.199)	0.83 (0.58 to 1.20, p=0.330)
Because training days don't work well for me	0.95 (0.63 to 1.40, p=0.784)	0.83 (0.51 to 1.31, p=0.430)
Because my friends don't engage in physical activity	1.37 (0.84 to 2.20, p=0.203)	0.81 (0.44 to 1.44, p=0.475)
Because I don't have enough physical fitness	2.65 (1.74 to 4.05, p<0.001)	1.12 (0.64 to 1.94, p=0.700)
Because I'm not interested in physical activity	1.65 (1.01 to 2.67, p=0.041)	1.37 (0.71 to 2.60, p=0.343)
Because I feel embarrassed about my body when I engage in physical activity	4.22 (2.88 to 6.22, p<0.001)	1.67 (0.93 to 2.97, p=0.085)
Because I don't enjoy physical activity	1.22 (0.77 to 1.91, p=0.389)	0.67 (0.35 to 1.27, p=0.228)
Because I feel that my physical appearance is worse than that of others	4.37 (3.00 to 6.42, p<0.001)	2.41 (1.35 to 4.28, p=0.003)
Because nobody encourages me to engage in physical activity	2.90 (1.80 to 4.72, p<0.001)	1.92 (1.08 to 3.43, p=0.026)
Because there are no physical activities that I enjoy	1.87 (1.22 to 2.86, p=0.004)	1.26 (0.72 to 2.19, p=0.417)
Because I am afraid of making a fool of myself	3.26 (2.25 to 4.74, p<0.001)	1.04 (0.60 to 1.78, p=0.892)
Because I consider that I already do enough physical activity in my physical education classes	1.20 (0.68 to 2.07, p=0.513)	0.95 (0.47 to 1.87, p=0.895)
	Anxiety	
	OR (95% CI, p value) (univariable)	OR (95% CI, p value) (multivariable)
Because I have a lot of homework	1.24 (0.92 to 1.68, p=0.158)	0.83 (0.58 to 1.18, p=0.296)
Because training days don't work well for me	1.23 (0.84 to 1.78, p=0.289)	1.14 (0.73 to 1.77, p=0.554)
Because my friends don't engage in physical activity	1.20 (0.74 to 1.93, p=0.446)	0.65 (0.36 to 1.16, p=0.150)
Because I don't have enough physical fitness	2.99 (1.97 to 4.58, p<0.001)	1.45 (0.84 to 2.51, p=0.179)
Because I'm not interested in physical activity	1.74 (1.08 to 2.79, p=0.023)	1.33 (0.71 to 2.50, p=0.370)
Because I feel embarrassed about my body when I engage in physical activity.	3.77 (2.58 to 5.56, p<0.001)	1.40 (0.78 to 2.49, p=0.251)
Because I don't enjoy physical activity	1.38 (0.88 to 2.13, p=0.154)	0.80 (0.43 to 1.47, p=0.475)
Because I feel that my physical appearance is worse than that of others	4.20 (2.88 to 6.18, p<0.001)	2.65 (1.51 to 4.71, p=0.001)
Because nobody encourages me to engage in physical activity	2.87 (1.78 to 4.69, p<0.001)	1.97 (1.11 to 3.50, p=0.021)
Because there are no physical activities that I enjoy	2.12 (1.39 to 3.23, p<0.001)	1.38 (0.80 to 2.37, p=0.244)
Because I am afraid of making a fool of myself	2.80 (1.94 to 4.06, p<0.001)	0.83 (0.47 to 1.41, p=0.490)
Because I consider that I already do enough physical activity in my physical education classes	1.27 (0.73 to 2.16, p=0.389)	0.87 (0.44 to 1.67, p=0.676)
	Stress	
	OR (95% CI, p value) (univariable)	OR (95% CI, p value) (multivariable)
Because I have a lot of homework	1.48 (1.11 to 1.99, p=0.008)	1.05 (0.75 to 1.48, p=0.769)
Because training days don't work well for me	1.15 (0.79 to 1.66, p=0.468)	0.92 (0.60 to 1.42, p=0.722)
Because my friends don't engage in physical activity	1.56 (0.98 to 2.48, p=0.062)	0.97 (0.55 to 1.70, p=0.918)
Because I don't have enough physical fitness	2.87 (1.88 to 4.43, p<0.001)	1.36 (0.78 to 2.37, p=0.284)
Because I'm not interested in physical activity	1.80 (1.12 to 2.89, p=0.015)	1.40 (0.75 to 2.62, p=0.294)
Because I feel embarrassed about my body when I engage in physical activity	3.38 (2.31 to 5.00, p<0.001)	1.05 (0.58 to 1.87, p=0.878)

Continued

Table 2 Continued

	Stress	
	OR (95% CI, p value) (univariable)	OR (95% CI, p value) (multivariable)
Because I don't enjoy physical activity	1.42 (0.92 to 2.20, p=0.109)	0.89 (0.48 to 1.62, p=0.707)
Because I feel that my physical appearance is worse than that of others	4.21 (2.86 to 6.27, p<0.001)	2.82 (1.59 to 5.07, p<0.001)
Because nobody encourages me to engage in physical activity	2.83 (1.74 to 4.67, p<0.001)	1.99 (1.12 to 3.59, p=0.021)
Because there are no physical activities that I enjoy	2.22 (1.46 to 3.40, p<0.001)	1.55 (0.91 to 2.67, p=0.109)
Because I am afraid of making a fool of myself	2.96 (2.04 to 4.31, p<0.001)	0.95 (0.55 to 1.62, p=0.861)
Because I consider that I already do enough physical activity in my physical education classes	0.92 (0.53 to 1.56, p=0.751)	0.55 (0.28 to 1.07, p=0.085)

Adjusted for age, sex, socioeconomic status, body mass index, physical activity, sedentary behaviour and overall sleep duration. Bold indicates a p value < 0.05.

depression, anxiety and stress. Future prospective studies are necessary to explore this potential pathway.

On the other hand, social support has been cross-sectionally and longitudinally associated with the practice of PA in adolescents.⁴³ Otherwise, having social support that encourages and participates in PA can positively influence mental health outcomes.^{43 46} However, when individuals perceive a lack of support or encouragement from their friends, family or community, it can undermine their motivation and confidence to engage in physical activities.^{43 46} Furthermore, the absence of a supportive environment can increase feelings of loneliness and isolation, which are risk factors for depression, anxiety and stress.^{10 27 47 48} Therefore, increasing parents' and friends' encouragement to higher practice of PA among adolescents can be essential.⁴⁴ For instance, de Camargo *et al*⁴⁴ showed that Brazilian adolescents who received social support from parents or friends had greater odds of engaging in PA and meeting the PA WHO recommendations.⁴⁹ Additionally, the literature has shown that doing PA together,⁴⁴ watching other adolescents doing it,⁴³ talking about PA with other adolescents, praising⁴⁰ and providing support for active commuting⁵⁰ can contribute to adolescents being physically active.

Additionally, regular PA is an effective coping mechanism for managing and reducing symptoms of stress, anxiety and depression.^{10 27 47 48} When individuals face barriers to PA related to physical appearance or lack of social support, they may lose out on the benefits of exercise as a coping strategy.^{10 27 47 48} And, without these beneficial effects, they may struggle to find alternative healthy ways to manage their mental health, leading to a higher likelihood of experiencing symptoms of depression, anxiety and stress.^{10 27 47 48} An overview of systematic reviews, which included the results of 97 reviews (1039 trials and 128 119 participants), synthesised the evidence on the effects of PA interventions on symptoms of depression, anxiety and psychological distress and showed that PA can be effective in reducing mild-to-moderate

symptoms of depression, anxiety and psychological distress compared with usual care.⁴⁷ In the present study, some barriers related to PA (self-perceived physical appearance and lack of social support) were associated with an increased likelihood of depression, anxiety and stress. These barriers may lead to a lower level of PA in adolescents, which could lead to greater symptoms of depression, anxiety and stress. Understanding perceived barriers to PA practice may contribute to developing strategies to promote PA for this population,¹⁹ and improve their mental health.⁴⁷

This study was not without limitations. First, the cross-sectional study does not establish cause-and-effect associations. Although the causal relationship between healthy behaviours (eg, PA, adequate sleep and low perceived barriers to PA) and mental health problems remains unclear, evidence suggests a significant association between these variables. Furthermore, the self-report questionnaires used in this study may lead to social desirability and recall bias. On the other hand, this study has some strengths that must be recognised. For instance, this is the first study to examine the relationship between perceived barriers to PA and depression, anxiety and stress symptoms in a sample of Spanish adolescents. Moreover, the analyses presented were adjusted for numerous covariates that could influence the results obtained, providing greater robustness to the findings. However, residual confusion is still possible. Despite these limitations, the implications of this study for adolescent health and exercise programmes are significant, suggesting that interventions should focus on addressing perceived barriers to PA. Potential intervention designs could include programmes that enhance social support from family and peers, which is related to PA levels and mental health outcomes. Additionally, efforts to improve adolescents' body image and self-esteem could reduce the impact of physical appearance-related barriers. Future steps could involve developing comprehensive school-based programmes incorporating these elements,

promoting an environment that supports regular PA and concurrently addresses mental health.

CONCLUSIONS

The results showed that perceiving barriers related to physical appearance and perceiving a lack of social support seem to be associated with symptoms of depression, anxiety and stress in adolescents. One of the most relevant practical applications of this study is that promoting PA by reducing some specific barriers could be an important step in reducing symptoms of depression, anxiety and stress in adolescents.

Author affiliations

¹Department of Physical Education, Universidade Federal do Paraná, Curitiba, Brazil

²Institute for Health and Sport, Victoria University, Melbourne, Victoria, Australia

³Department of Physical Medicine and Nursing, University of Zaragoza, Zaragoza, Spain

⁴Health and Social Research Center, Universidad de Castilla-La Mancha, Cuenca, Spain

⁵Higher Institute of Physical Education, Universidad de la República, Rivera, Uruguay

⁶IdiSNA, Instituto de Investigación Sanitaria de Navarra, Pamplona, Spain

⁷Department of Preventive Medicine and Public Health, Facultad de Medicina, Universidad de Navarra, Pamplona, Spain

⁸Pathophysiology of Obesity and Nutrition, Centro de Investigación Biomédica en Red, Instituto de Salud Carlos III, Madrid, Spain

⁹Centro de Investigación Biomédica en Red de Salud Mental, Instituto de Salud Carlos III, Madrid, Spain

¹⁰One Health Research Group, Universidad de Las Américas, Quito, Ecuador

X Edina Maria de Camargo @camargoedina_, Sitong Chen @chenst1212, Rubén López-Bueno @rlopezbu, Arthur Eumann Mesas @MesasArthur, Estela Jiménez-López @ELopezJimenez and José Francisco López-Gil @JFLopezGil

Acknowledgements The authors wish to extend their thanks to *Ayuntamiento de Archena*, along with all the teenagers, parents or legal guardians, physical education instructors, schools and staff members who participated.

Contributors Conceptualisation: JFL-G and EMdC; methodology: JL-G; formal analysis: JFL-G; data curation: JFL-G; writing—original draft preparation: EMdC and JFL-G; writing—review and editing: SC, RB, AEM, BB-P, NM-C and EJ-L. All authors have read and agreed to the published version of the manuscript.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient and public involvement Patients and/or the public were involved in the design, or conduct, or reporting, or dissemination plans of this research. Refer to the Methods section for further details.

Patient consent for publication Consent obtained from parent(s)/guardian(s).

Ethics approval This study involves human participants and this research project obtained ethics approval from the Bioethics Committee of the University of Murcia (ID 2218/2018) and the Ethics Committee of the Albacete University Hospital Complex and the Albacete Integrated Care Management (ID 2021-85). It was carried out following the Declaration of Helsinki, respecting the human rights of the participants enrolled. Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request. Not applicable.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability

of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

José Francisco López-Gil <http://orcid.org/0000-0002-7412-7624>

REFERENCES

- World Health Organization. Mental disorders. 2022. Available: <https://www.who.int/news-room/fact-sheets/detail/mental-disorders>
- Institute for Health Metrics and Evaluation. Global health data exchange (GHDx). 2019. Available: <https://vizhub.healthdata.org/gbd-results>
- Vos T, Lim SS, Abbafati C, et al. Global burden of 369 diseases and injuries in 204 countries and territories, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet* 2020;396:1204–22.
- GBD 2019 Mental Disorders Collaborators. Global, regional, and national burden of 12 mental disorders in 204 countries and territories, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet Psychiatry* 2022;9:137–50.
- Liu NH, Daumit GL, Dua T, et al. Excess mortality in persons with severe mental disorders: a multilevel intervention framework and priorities for clinical practice, policy and research agendas. *World Psychiatry* 2017;16:30–40.
- Solmi M, Radua J, Olivola M, et al. Age at onset of mental disorders worldwide: large-scale meta-analysis of 192 epidemiological studies. *Mol Psychiatry* 2022;27:281–95.
- Colizzi M, Lasalvia A, Ruggeri M. Prevention and early intervention in youth mental health: is it time for a multidisciplinary and trans-diagnostic model for care? *Int J Ment Health Syst* 2020;14:23.
- Das JK, Salam RA, Lassi ZS, et al. Interventions for Adolescent Mental Health: An Overview of Systematic Reviews. *J Adolesc Health* 2016;59:S49–60.
- Lund C, Brooke-Sumner C, Baingana F, et al. Social determinants of mental disorders and the Sustainable Development Goals: a systematic review of reviews. *Lancet Psychiatry* 2018;5:357–69.
- Schuch FB, Vancampfort D. Physical activity, exercise, and mental disorders: it is time to move on. *Trends Psychiatry Psychother* 2021;43:177–84.
- Chaput J-P, Willumsen J, Bull F, et al. 2020 WHO guidelines on physical activity and sedentary behaviour for children and adolescents aged 5–17 years: summary of the evidence. *Int J Behav Nutr Phys Act* 2020;17:141.
- Patton GC, Sawyer SM, Santelli JS, et al. Our future: a Lancet commission on adolescent health and wellbeing. *Lancet* 2016;387:2423–78.
- Christofaro DGD, Andersen LB, Andrade SM de, et al. Adolescents' physical activity is associated with previous and current physical activity practice by their parents. *J Pediatr (Rio J)* 2018;94:48–55.
- García-Hermoso A, López-Gil JF, Ramírez-Vélez R, et al. Adherence to aerobic and muscle-strengthening activities guidelines: a systematic review and meta-analysis of 3.3 million participants across 32 countries. *Br J Sports Med* 2023;57:225–9.
- Sechrist KR, Walker SN, Pender NJ. Development and psychometric evaluation of the exercise benefits/barriers scale. *Res Nurs Health* 1987;10:357–65.
- Jodkowska M, Oblacińska A, Nałęcz H, et al. Perceived barriers for physical activity in overweight and obese adolescents and their association with health motivation. *Dev Period Med* 2017;21:248–58.
- Park JH, Moon JH, Kim HJ, et al. Sedentary Lifestyle: Overview of Updated Evidence of Potential Health Risks. *Korean J Fam Med* 2020;41:365–73.
- Christofoletti M, Streit IA, Garcia LMT, et al. Barreiras e facilitadores para a prática de atividade física em diferentes domínios no Brasil: uma revisão sistemática. *Ciênc saúde coletiva* 2022;27:3487–502.
- Rech CR, Camargo EM de, De Araujo PAB, et al. PERCEIVED BARRIERS TO LEISURE-TIME PHYSICAL ACTIVITY IN THE BRAZILIAN POPULATION. *Rev Bras Med Esporte* 2018;24:303–9.
- Sallis JF, Prochaska JJ, Taylor WC. A review of correlates of physical activity of children and adolescents. *Med Sci Sports Exerc* 2000;2000:963–75.

- 21 Ekelund U, Tarp J, Steene-Johannessen J, *et al.* Dose-response associations between accelerometry measured physical activity and sedentary time and all cause mortality: systematic review and harmonised meta-analysis. *BMJ* 2019;366:14570.
- 22 Lee IM, Shiroma EJ, Lobelo F, *et al.* Effect of physical inactivity on major non-communicable diseases worldwide: an analysis of burden of disease and life expectancy. *The Lancet* 2012;380:219–29.
- 23 World Health Organization. Global action plan on physical activity 2018–2030: more active people for a healthier world. 2018.
- 24 Tsuang MT, Bar JL, Stone WS, *et al.* Gene-environment interactions in mental disorders. *World Psychiatry* 2004;3:73–83.
- 25 Arango C, Dragioti E, Solmi M, *et al.* Risk and protective factors for mental disorders beyond genetics: an evidence-based atlas. *World Psychiatry* 2021;20:417–36.
- 26 Rodríguez-Ayllon M, Cadenas-Sánchez C, Estévez-López F, *et al.* Role of Physical Activity and Sedentary Behavior in the Mental Health of Preschoolers, Children and Adolescents: A Systematic Review and Meta-Analysis. *Sports Med* 2019;49:1383–410.
- 27 Recchia F, Bernal JDK, Fong DY, *et al.* Physical Activity Interventions to Alleviate Depressive Symptoms in Children and Adolescents: A Systematic Review and Meta-analysis. *JAMA Pediatr* 2023;177:132–40.
- 28 López-Gil JF. The Eating Healthy and Daily Life Activities (EHDLA) Study. *Children (Basel)* 2022;9:370.
- 29 Cabanas-Sánchez V, Tejero-González CM, Veiga OL. Construcción y validación de una escala breve de percepción de barreras para la práctica deportiva en adolescentes. *Rev Esp Salud Publica* 2012;86:435–43.
- 30 Lovibond PF, Lovibond SH. The structure of negative emotional states: comparison of the Depression Anxiety Stress Scales (DASS) with the Beck Depression and Anxiety Inventories. *Behav Res Ther* 1995;33:335–43.
- 31 Bados A, Solanas A, Andres R. Psychometric properties of the Spanish version of Depression, Anxiety and Stress Scales (DASS). *Psicothema* 2005;17:679e83.
- 32 Román F, Santibáñez P, Vinet EV. Uso de las Escalas de Depresión Ansiedad Estrés (DASS-21) como Instrumento de Tamizaje en Jóvenes con Problemas Clínicos. Esta investigación fue financiada por el Proyecto DIDUFRO DI11-0003 de la Universidad de La Frontera, Temuco – Chile. Además, contó con los aportes del Dr. David Mellor de Deakin University, Australia, cuya estadia en la Universidad de La Frontera fue financiada por el Programa CONICYT 80110035 de Atracción de Capital Humano Avanzado del Extranjero Modalidad Estadias Cortas (MEC). *Acta Invest Psicol* 2016;6:2325–36.
- 33 Currie C, Molcho M, Boyce W, *et al.* Researching health inequalities in adolescents: the development of the Health Behaviour in School-Aged Children (HBSC) family affluence scale. *Soc Sci Med* 2008;66:1429–36.
- 34 Segura-Díaz JM, Barranco-Ruiz Y, Saucedo-Araujo RG, *et al.* Feasibility and reliability of the Spanish version of the Youth Activity Profile questionnaire (YAP-Spain) in children and adolescents. *J Sports Sci* 2021;39:801–7.
- 35 Saint-Maurice PF, Welk GJ, Beyler NK, *et al.* Calibration of self-report tools for physical activity research: the Physical Activity Questionnaire (PAQ). *BMC Public Health* 2014;14:461.
- 36 Saint-Maurice PF, Welk GJ. Validity and Calibration of the Youth Activity Profile. *PLoS One* 2015;10:e0143949.
- 37 Wood SN. Fast Stable Restricted Maximum Likelihood and Marginal Likelihood Estimation of Semiparametric Generalized Linear Models. *J R Stat Soc Ser B* 2011;73:3–36.
- 38 Marra G, Wood SN. Practical variable selection for generalized additive models. *Comput Stat Data Anal* 2011;55:2372–87.
- 39 Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *J R Stat Soc Ser B* 1995;57:289–300.
- 40 Gualdi-Russo E, Rinaldo N, Zaccagni L. Physical Activity and Body Image Perception in Adolescents: A Systematic Review. *Int J Environ Res Public Health* 2022;19:13190.
- 41 Fowler LA, Kracht CL, Denstel KD, *et al.* Bullying experiences, body esteem, body dissatisfaction, and the moderating role of weight status among adolescents. *J Adolesc* 2021;91:59–70.
- 42 Wang S, Sun Q, Zhai L, *et al.* The Prevalence of Depression and Anxiety Symptoms among Overweight/Obese and Non-Overweight/Non-Obese Children/Adolescents in China: A Systematic Review and Meta-Analysis. *IJERPH* 2019;16:340.
- 43 Mendonça G, Cheng LA, Mélo EN, *et al.* Physical activity and social support in adolescents: a systematic review. *Health Educ Res* 2014;29:822–39.
- 44 de Camargo EM, da Costa CG, Piola TS, *et al.* Is Greater Social Support from Parents and Friends Related to Higher Physical Activity Levels among Adolescents? *Children (Basel)* 2023;10:701.
- 45 Liu Z, Sun L, Zhang Y, *et al.* The prevalence of underweight and obesity in Chinese children and adolescents with major depressive disorder and relationship with suicidal ideation and attempted suicide. *Front Psychiatry* 2023;14:1130437.
- 46 Laird Y, Fawkner S, Kelly P, *et al.* The role of social support on physical activity behaviour in adolescent girls: a systematic review and meta-analysis. *Int J Behav Nutr Phys Act* 2016;13:79.
- 47 Singh B, Olds T, Curtis R, *et al.* Effectiveness of physical activity interventions for improving depression, anxiety and distress: an overview of systematic reviews. *Br J Sports Med* 2023;57:1203–9.
- 48 Heissel A, Heinen D, Brokmeier LL, *et al.* Exercise as medicine for depressive symptoms? A systematic review and meta-analysis with meta-regression. *Br J Sports Med* 2023;57:1049–57.
- 49 Bull FC, Al-Ansari SS, Biddle S, *et al.* World Health Organization 2020 guidelines on physical activity and sedentary behaviour. *Br J Sports Med* 2020;54:1451–62.
- 50 Silva KS, Pizarro AN, Garcia LMT, *et al.* Which social support and psychological factors are associated to active commuting to school? *Prev Med* 2014;63:20–3.