

Genetic parameters and direct, maternal and heterosis effects on litter size in a diallel cross among three commercial varieties of Iberian pig

J. L. Noguera¹, N. Ibáñez-Escriche², J. Casellas³, J. P. Rosas⁴ and L. Varona^{5†} 

¹Genética i Milllora Animal, IRTA, Avinguda Alcalde Rovira Roure, 191, 25198 Lleida, Spain; ²Departament de Ciència Animal, Universitat Politècnica de València, 46071 València, Spain; ³Departament de Ciència Animal i dels Aliments, Universitat Autònoma de Barcelona, 08193 Bellaterra (Barcelona), Spain; ⁴Programa de Mejora Genética "CASTÚA", INGA FOOD S.A. (Nutreco Group), C./ Av. de la Rúa, 2, 06200 Almendralejo, Badajoz, Spain; ⁵Unidad de Genética Cuantitativa y Mejora Animal, Universidad de Zaragoza, Instituto Agroalimentario de Aragón (IA2), 50013 Zaragoza, Spain

(Received 21 September 2018; Accepted 15 April 2019; First published online 4 June 2019)

The Iberian pig is one of the pig breeds that has the highest meat quality. Traditionally, producers have bred one of the available varieties, exclusively, and have not used crosses between them, which has contrasted sharply with other populations of commercial pigs for which crossbreeding has been a standard procedure. The objective of this study was to perform an experiment under full diallel design among three contemporary commercial varieties of Iberian pig and estimate the additive genetic variation and the crossbreeding effects (direct, maternal and heterosis) for prolificacy. The data set comprised 18 193 records for total number born and number born alive from 3800 sows of three varieties of the Iberian breed (Retinto, Torbiscal and Entrepelado) and their reciprocal crosses (Retinto × Torbiscal, Torbiscal × Retinto, Retinto × Entrepelado, Entrepelado × Retinto, Torbiscal × Entrepelado and Entrepelado × Torbiscal), and a pedigree of 4609 individuals. The analysis was based on a multiple population repeatability model, and we developed a model comparison test that indicated the presence of direct line, maternal and heterosis effects. The results indicated the superiorities of the direct line effect of the Retinto and the maternal effect of the Entrepelado populations. All of the potential crosses produced significant heterosis, and additive genetic variation was higher in the Entrepelado than it was in the other two populations. The recommended cross for the highest yield in prolificacy is a Retinto father and an Entrepelado mother to generate a hybrid commercial sow.

Keywords: prolificacy, diallel cross, heritability, crossbreeding, Iberian pig

Implications

Reproductive efficiency is a limiting factor in the production of Iberian pigs. This study showed that litter size in the Iberian pig can be increased through two non-exclusive strategies. The use of within-line selection is supported by the presence of sufficient additive genetic variation, and crossbreeding is supported by estimates of the crossbreeding effects derived from a full diallel experiment among three varieties (Entrepelado, Retinto and Torbiscal). In addition, the optimal hybrid sow can be produced by crossing a Retinto father and an Entrepelado mother.

Introduction

The Iberian pig breed is recognized as one of the porcine populations that has the highest meat quality (Serra *et al.*, 1998; Gandemer, 2009), which is an important reason for its conservation because it has slower growth and lower feed efficiency (Barea *et al.*, 2011) and prolificacy (Silió *et al.*, 2001) than do other commercial pig populations. Furthermore, a profound transformation that has involved the replacement of many traditional producers by intensive management farms has affected Iberian pig production. Normative that regulates the Iberian pig products forces that the sow is Iberian, whereas the boar could be either Iberian, Duroc (DU) or hybrid between these two breeds (Boletín Oficial del Estado, 2014). In fact, a large proportion of Iberian pig farmers uses DU sires to obtain better growth rate and efficiency (Serrano *et al.*, 2008). Therefore, under intensive

[†] Present address: Luis Varona, Unidad de Genética Cuantitativa y Mejora Animal, Universidad de Zaragoza, Instituto Agroalimentario de Aragón (IA2), 50013 Zaragoza, Spain. E-mail: lvarona@unizar.es

management, improvement in the reproductive efficiency of Iberian sows is essential for its economic viability.

The number of weaned piglets per sow in a year (or numerical productivity) is one of the most important factors in the profitability of pig farms, and litter size is its most important component (Quinton *et al.*, 2006; Dekkers *et al.*, 2011). Two non-exclusive strategies can lead to the genetic improvement of litter size: (1) within-line selection and (2) appropriate crossbreeding between lines to exploit heterosis.

Traditionally, genetic improvement in litter size in Iberian pig populations has been limited (Fernández *et al.*, 2008), and Iberian pig farmers have not used crossbred sows; rather, they have bred one of the varieties of the Iberian breed, exclusively (Martínez *et al.*, 2000), which differed markedly from the practices used with other populations of commercial pigs in which selection efforts for prolificacy in maternal lines have been common and crossbreeding has been a standard procedure (Dekkers *et al.*, 2011). Several studies have identified genetic variability for prolificacy within (Rodríguez *et al.*, 1994, Fernández *et al.*, 2008) and between (García-Casco *et al.*, 2012) varieties of Iberian pig. Thus, within-line selection and the implementation of a maternal crossbreeding scheme are plausible strategies for genetic improvement; however, to identify the most appropriate strategy, reliable estimates of the heritability of prolificacy in each population and estimates of the crossbreeding parameters as defined by the Dickerson's model are required (Dickerson, 1969).

The objective of this study was to estimate the additive genetic variation (or heritability) and the crossbreeding effects (direct, maternal and heterosis) for prolificacy using data from an experiment based on a full diallel design (Hayman, 1954) among three contemporary commercial varieties of Iberian pig [Entrepelado (E), Retinto (R) and Torbiscal (T)] under intensive management conditions. The overarching objective of the study was to identify suitable genetic improvement strategies within a pyramidal breeding program among the Iberian pig populations.

Materials and methods

Animals and experimental design

The data set comprised 18 193 records for total number born (TNB) and number born alive (NBA) from 3800 sows that had been obtained from a full diallelic experiment among three varieties of the Iberian pig breed (Retinto x Retinto (RR), Torbiscal x Torbiscal (TT) and Entrepelado x Entrepelado (EE)) and their reciprocal crosses (Retinto x Torbiscal (RT), Torbiscal x Retinto (TR), Retinto x Entrepelado (RE), Entrepelado x Retinto (ER), Torbiscal x Entrepelado (TE) and Entrepelado x Torbiscal (ET)). The three varieties are recognized in Spain's official Iberian herd-book [Spanish Association of Iberian Purebred Pig Breeders (AECERIBER)]. A detailed description of their characteristics is provided by Ibáñez-Escriche *et al.* (2016). Purebred sows were located in two nucleus herds in intensive management systems

(Herds 1 and 2), and the service boars for these sows were kept at an artificial insemination center. A third herd (HERD 3) was a commercial production herd in which all sows, purebred or crossbred, were mated with boars from a DU population following the standard production system of Iberian pigs under intensive management. The distribution of the data between herds and breed of service sire and the average and SD of NBA and TNB for all crosses are presented in Table 1. In addition, the pedigree was extended back up to 3 generations, and it consisted of 4609 individuals. The number of founders in each per population was 47 Entrepelado (13 sires and 34 dams), 80 Retinto (18 sires and 62 dams) and 107 Torbiscal (38 sires and 69 dams).

Statistical analyses

TNB and NBA were analyzed with a multiple population repeatability model following García-Cortés and Toro (2006), which can divide the additive genetic values into separate fractions depending on their genetic origin. The general model of the analysis was as follows:

$$\mathbf{y} = \mathbf{X}\mathbf{b} + \sum_{i=E,R,T}^3 \mathbf{Z}_i\mathbf{a}_i + \mathbf{W}\mathbf{p} + \mathbf{e}$$

where $\mathbf{y}$ is the vector of phenotypic records (TNB and NBA), $\mathbf{b}$ is the vector of systematic effects: order of parity (six levels: first, second, third, fourth, fifth and sixth and beyond), herd-year-season (85 levels), genetic line of service boar [four levels: E, R, T and DU] and the genetic line of the sow (L_E , L_R , L_T for E, T and R, respectively), genetic line of the mother of the sow (M_E , M_R and M_T for E, T and R, respectively) and heterosis (H_{ER} , H_{ET} and H_{RT} between E and R, E and T and R and T, respectively) effects following Dickerson's model (Dickerson, 1969). Furthermore, $\mathbf{a}_i$ is the vector of random additive genetic effects of pure and crossbred individuals of the i th line –E, R or T– (with 1934, 2748 and 1765 levels, respectively), $\mathbf{p}$ is the vector of the permanent effect of the sow (3800 levels) and $\mathbf{e}$ is the residuals vector; $\mathbf{X}$ and $\mathbf{W}$ are known incidence matrices that link fixed and permanent random effects with the vector $\mathbf{y}$. Further, $\mathbf{Z}_E$, $\mathbf{Z}_R$ and $\mathbf{Z}_T$ are the matrices that links data with the random additive effects of each line with dimensions $18\,193 \times 1934$, $18\,193 \times 2748$ and $18\,193 \times 1765$ levels, respectively. The model allows for a different additive genetic variance component for each genetic origin, whereas the permanent environmental ($\mathbf{p}$) and the residual effects ($\mathbf{e}$) were assumed homogeneous between lines.

The model assumed flat prior distributions for systematic effects ($\mathbf{b}$) and multivariate Gaussian distributions for the additive ($\mathbf{a}_i$), permanent environmental ($\mathbf{p}$) and residual effects ($\mathbf{e}$). The permanent environmental effects and the residuals were assumed identically and independently distributed; therefore, their prior distribution was as follows:

$$\mathbf{p} \sim N(0, I\sigma_p^2) \quad \mathbf{e} \sim N(0, I\sigma_e^2)$$

Table 1. Number of recorded parities by Iberian pig breed of sow, service sire and herd, and the mean and SD of the NBA and the TNB

Dam (number of sows)	Sire							Total	NBA Mean (SD)	TNB Mean (SD)
	HERD 1			HERD 2			HERD 3			
	E	R	T	E	R	T	DU			
EE (707)	517	752	5	255	811	70	433	2843	7.73 (2.15)	7.95 (2.16)
ER (527)	—	—	—	—	—	—	2336	2336	8.31 (2.25)	8.53 (2.27)
ET (177)	—	—	—	—	—	—	942	942	7.81 (2.25)	8.02 (2.25)
RE (196)	—	—	—	—	—	—	806	806	8.55 (2.43)	8.84 (2.45)
RR (874)	1450	491	96	655	277	633	870	4472	8.05 (2.19)	8.39 (2.20)
RT (488)	—	—	—	—	—	—	2450	2450	8.31 (2.43)	8.60 (2.43)
TE (36)	—	—	—	—	—	—	193	193	8.34 (2.47)	8.69 (2.49)
TR (343)	—	—	—	—	—	—	1993	1993	8.29 (2.28)	8.53 (2.32)
TT (452)	197	808	58	109	507	247	232	2158	7.21 (2.12)	7.59 (2.13)
Total (3 800)	2 164	2 051	159	1 019	1 595	950	10 255	18 193	8.01 (2.27)	8.29 (2.28)

E = Entrepelado; R = Retinto; T = Torbiscal; DU = Duroc; EE = Entrepelado × Entrepelado; ER = Entrepelado × Retinto; ET = Entrepelado × Torbiscal; RE = Retinto × Entrepelado; RR = Retinto × Retinto; RT = Retinto × Torbiscal; TE = Torbiscal × Entrepelado; TR = Torbiscal × Retinto; TT = Torbiscal × Torbiscal; NBA = number born alive; TNB = total number born.

The assumed prior distribution for the additive effects was as follows:

$$\mathbf{a}_E \sim N(0, A_E \sigma_{a_E}^2) \quad \mathbf{a}_R \sim N(0, A_R \sigma_{a_R}^2) \quad \mathbf{a}_T \sim N(0, A_T \sigma_{a_T}^2)$$

where A_E , A_R and A_T are the partial relationship matrices generated by the founders of E, R and T, respectively. The calculation of those matrices was performed following the algorithm proposed by García-Cortés and Toro (2006), and they are illustrated with a tiny example in the Appendix. Prior distributions for the variance components ($\sigma_{(a_E)}^2$, $\sigma_{(a_R)}^2$, $\sigma_{(a_T)}^2$, σ_p^2 and σ_e^2) were uniform.

The above model leads to the following mixed model equations:

$$\begin{bmatrix} X'X & X'Z_E & X'Z_R & X'Z_T & X'W \\ Z_E'X & Z_E'Z_E + A_E^- \frac{\sigma_e^2}{\sigma_{a_E}^2} & Z_E'Z_R & Z_E'Z_T & Z_E'W \\ Z_R'X & Z_R'Z_E & Z_R'Z_R + A_R^- \frac{\sigma_e^2}{\sigma_{a_R}^2} & Z_R'Z_T & Z_R'W \\ Z_T'X & Z_T'Z_E & Z_T'Z_R & Z_T'Z_T + A_T^- \frac{\sigma_e^2}{\sigma_{a_T}^2} & Z_T'W \\ W'X & W'Z_E & W'Z_R & W'Z_T & W'W + I \frac{\sigma_e^2}{\sigma_p^2} \end{bmatrix} \begin{bmatrix} \hat{\mathbf{b}} \\ \hat{\mathbf{a}}_E \\ \hat{\mathbf{a}}_R \\ \hat{\mathbf{a}}_T \\ \hat{\mathbf{p}} \end{bmatrix} = \begin{bmatrix} X'y \\ Z_E'y \\ Z_R'y \\ Z_T'y \\ W'y \end{bmatrix}$$

where A_E^- , A_R^- and A_T^- are the inverses of non-zero part of the population specific partial numerator relationship matrices (see the Appendix).

From that general full model (LMH), the following models were defined by fixing to zero the direct (MH), maternal (LH), heterosis (LM), direct and maternal (H), direct and heterosis (M) and maternal and heterosis effects (L). The analysis of each model was performed using a Gibbs sampling algorithm

(Gelfand and Smith, 1990) with a single long chain of 1 000 000 iterations, following a 'burn in' period of 250 000 iterations.

Model comparisons

The models were compared using the deviance information criteria (DIC) (Spiegelhalter *et al.*, 2002) and the logarithm of the conditional predictive ordinate (LogCPO) (Gelfand, 1996).

Deviance information criteria: The DIC compares the global quality of two or more hierarchical models accounting for model complexity. For a particular model M, the DIC is defined as follows:

$$DIC_M = 2\bar{D}_M - D(\bar{\theta}_M),$$

where $\bar{D}_M$ is the posterior expectation of the deviance $D(\bar{\theta}_M)$ and $D(\bar{\theta}_M) = -2\log(p(\mathbf{y}|\bar{\theta}_M))$ is the deviance evaluated at the posterior mean estimate of the parameter vector ($\bar{\theta}_M$). The computation of DIC comprises two terms, $\bar{D}_M$ is a measure of model fit and $\bar{D}_M - D(\bar{\theta}_M)$ is related to the effective number of parameters. Models that have the smallest DIC have the best global fit after accounting for model complexity.

Log marginal probability: If we consider the data vector $\mathbf{y} = (y_i, \mathbf{y}_{-i})$, where y_i is the i th datum and $\mathbf{y}_{-i}$ is the vector of data with the i th datum deleted, the conditional predictive distribution has a probability density equal to the following:

$$p(y_i|\mathbf{y}_{-i}) = \int p(y_i|\mathbf{y}_{-i}, \theta) p(\theta|\mathbf{y}_{-i}) d\theta,$$

where θ is the vector of unknown parameters and random effects in the model. Therefore, $p(y_i|\mathbf{y}_{-i})$ can be interpreted as the probability of each datum given the remaining data and is the conditional predictive ordinate (CPO) for the

Table 2. Differences with the best model for DIC and the LogCPO for NBA and TNB among three varieties of Iberian pig

Model	NBA		TNB	
	DIC	LogCPO	DIC	LogCPO
LMH	—	—	—	—
MH	6.3	−0.9	5.9	−0.4
LH	6.3	−5.0	5.7	−2.9
LM	23.8	−19.8	22.4	−20.2
L	26.3	−22.3	25.8	−23.0
M	26.6	−20.9	26.1	−23.3
H	7.5	−6.9	8.0	−4.8

LMH = full model with direct line; maternal and heterosis effects; MH = reduced model with maternal and heterosis effects; LH = reduced model with direct line and heterosis effects; LM = reduced model with direct line and maternal effects; L = reduced model with direct line effects; M = reduced model with maternal effects; H = reduced model with Heterosis effects; DIC = deviance information criteria; LogCPO = logarithm of the conditional predictive ordinate; NBA = number born alive; TNB = total number born.

i th datum. The pseudo log-marginal probability of the data is as follows:

$$\sum_i \ln p(y_i | \mathbf{y}_{-i}).$$

A Monte Carlo approximation of the CPO (Gelfand, 1996)

is $\sum_i \ln \hat{p}(y_i | \mathbf{y}_{-i})$, where $\hat{p}(y_i | \mathbf{y}_{-i}) = N \left[\sum_{j=1}^N \frac{1}{p(y_i | \theta^j)} \right]^{-1}$,

and N is the number of Markov chain Monte Carlo (MCMC) draws, and θ^j is the j th draw from the posterior distribution of the corresponding parameter. The higher the value of the LogCPO, the better the fit of the model to the data.

Results

For both NBA and TNB, the model with best fit with DIC and LogCPO was the complete LMH model, followed by a group of models (LH, MH and H) whose differences from the best model ranged from 5.9 to 8.0 units for DIC and from 0.4 to 6.9 for LogCPO (Table 2). The LM, L and M models had a worse adjustment and their differences from the LMH model ranged from 22.4 to 26.6 for DIC and from 19.8 to 23.3 for LogCPO.

The posterior mean estimates ($\pm$ posterior SDs) of the additive genetic variance ranged from 0.371 ± 0.106 (T) to 0.665 ± 0.123 (E) and from 0.418 ± 0.115 (T) to 0.717 ± 0.126 (E) for NBA and TNB, respectively, and the posterior mean estimates ($\pm$ posterior SDs) of the permanent environmental and residual variances were 0.361 ± 0.051 and 4.020 ± 0.048 for NBA and 0.371 ± 0.053 and 4.029 ± 0.048 for TNB (Table 3). The posterior mean ($\pm$ posterior SDs) estimates for the population-specific heritabilities were 0.078 ± 0.021 (T), 0.084 ± 0.017 (R) and 0.131 ± 0.022 (E) for NBA and 0.086 ± 0.022 (T), 0.090 ± 0.017 (R) and 0.140 ± 0.022 (E) for TNB. The results from the MH, LH,

LM, L, M and H models were similar (Supplementary Tables S1 and S2 for NBA and TNB, respectively).

The posterior mean ($\pm$ posterior SD) estimates of the differences between L_E and L_R , L_E and L_T and L_R and L_T were -0.684 ± 0.317 , 0.114 ± 0.324 and 0.798 ± 0.210 for NBA and -0.876 ± 0.327 , -0.024 ± 0.334 and 0.852 ± 0.217 piglets for TNB (Table 4). In addition, the posterior mean ($\pm$ posterior SD) estimates of the differences of M_E with M_R and M_T were 0.443 ± 0.123 and 0.450 ± 0.144 for NBA and 0.554 ± 0.117 and 0.533 ± 0.117 piglets for TNB. The differences between M_R and M_T were only 0.007 ± 0.104 and -0.021 ± 0.107 . Those differences between the maternal effects were supported by the comparisons between reciprocals. The posterior mean ($\pm$ posterior SD) estimates of the differences between ER and RE and between ET and TE were -0.443 ± 0.123 and -0.554 ± 0.125 and -0.450 ± 0.144 and -0.534 ± 0.147 for NBA and TNB, respectively. The differences between TR and RT were negligible (-0.007 ± 0.104 and 0.021 ± 0.107). Furthermore, the posterior mean ($\pm$ posterior SD) estimates of the heterosis effects ranged from 0.600 ± 0.129 (H_{ET}) to 0.690 ± 0.092 (H_{RT}) for NBA and from 0.622 ± 0.131 (H_{ET}) to 0.666 ± 0.093 (H_{RT}) for TNB (Table 4). The results from the other models followed the same pattern (Supplementary Tables S3 and S4 for NBA and TNB, respectively).

The posterior mean and SDs of differences between each cross with respect to the RE ranged from -0.443 (with ER) to -1.566 piglets (with TT) piglets for NBA and from -0.464 (with TE) to -1.607 (with TT) for TNB, and the posterior probability that RE is the best cross given the LMH model was 0.985 and 0.991 for NBA and TNB, respectively (Table 5).

Discussion

The main advantage of the multiple population repeatability model proposed by García-Cortés and Toro (2006) is that it provides specific estimates of the additive variance component (and heritability) for each population of origin and, therefore, it can detect differences in their genetic variation. The additive variance (and heritability) of the E variety was markedly higher than that of the other two populations (R and T) in all models. The estimates of the E population were higher than those obtained in previous studies of other Iberian pig populations (Pérez-Enciso and Gianola, 1992; García-Casco *et al.*, 2012), and the estimates for R and T were closer to such published estimates; however, estimates of heritability over 0.10 are common in white pig (Bidanel, 2011; Putz *et al.*, 2015; Ogawa *et al.*, 2018) and in Iberian populations (Fernández *et al.*, 2008). The results of our study should be confirmed based on a larger database; however, given the results of our study, it is plausible that the response to selection in the E population under an appropriate breeding scheme to improve litter size might be greater than it would be in the other lines. Thus, the E variety is a good candidate for inclusion in a pyramidal scheme for the improvement of litter size.

Table 3. Posterior mean (and posterior SD) estimates [‘of?’] the additive (σ_{aE}^2 , σ_{aR}^2 and σ_{aT}^2), permanent environmental (σ_p^2), and residual variance (σ_e^2) components and population specific heritabilities (h_E^2 , h_R^2 , h_T^2) for NBA and TNB for each of three varieties of Iberian pig (Entrepelado (E), Retinto (R) and Torbiscal (T)) analyzed with full model with direct line, maternal and heterosis effect

	NBA	TNB
σ_{aE}^2	0.665 (0.123)	0.717 (0.126)
σ_{aR}^2	0.401 (0.088)	0.439 (0.091)
σ_{aT}^2	0.371 (0.106)	0.418 (0.115)
σ_p^2	0.361 (0.051)	0.371 (0.053)
σ_e^2	4.020 (0.048)	4.029 (0.048)
h_E^2	0.131 (0.022)	0.140 (0.022)
h_R^2	0.084 (0.017)	0.090 (0.017)
h_T^2	0.078 (0.021)	0.086 (0.022)

NBA = number born alive; TNB = total number born.

Table 4. Posterior mean (and posterior SD) estimates of the contrast between three varieties of Iberian pig for the direct line (L_E , L_R and L_T) and maternal (M_E , M_R and M_T) effects, and the heterosis (H_{ER} , H_{ET} and H_{RT}) effects on NBA and TNB under the full model with direct line, maternal and heterosis effects

		NBA	TNB
Direct line	L_E v. L_R	−0.684 (0.317)	−0.876 (0.327)
	L_E v. L_T	0.114 (0.324)	−0.024 (0.334)
	L_R v. L_T	0.798 (0.210)	0.852 (0.217)
Maternal	M_E v. M_R	0.443 (0.123)	0.554 (0.147)
	M_E v. M_T	0.450 (0.144)	0.533 (0.117)
	M_R v. M_T	0.007 (0.104)	−0.021 (0.107)
Heterosis	H_{ER}	0.653 (0.098)	0.661 (0.099)
	H_{ET}	0.600 (0.129)	0.622 (0.131)
	H_{RT}	0.690 (0.092)	0.666 (0.093)

L_E , L_R and L_T are the direct line effects for Entrepelado, Retinto and Torbiscal, respectively. M_E , M_R and M_T are the maternal line effects for Entrepelado, Retinto and Torbiscal. H_{ER} , H_{ET} and H_{RT} are the heterosis effects between Entrepelado and Retinto, Entrepelado and Torbiscal and Retinto and Torbiscal, respectively. NBA=number born alive; TNB=total number born.

The results of the comparison of models indicated the importance of the direct line, maternal and heterosis effects based on both measures (LogCPO and DIC) because the LMH model had the best fit; however, the extent of this relevance was mixed. For example, the best model was followed closely by a group of models that share the effects of heterosis (LH, MH and H), which is reinforced by a posterior probability of a heterosis effect > 0 that was > 0.999 for all models and traits. Therefore, there was clear evidence of heterosis between each of the three potential crosses and small differences among them. A significant degree of heterosis was expected because it is common in crosses between pig populations (Haley *et al.* 1995; Cassady *et al.*, 2002; García-Casco *et al.*, 2012). The results were surprising, however, because previous studies (Fabuel *et al.*, 2004) have suggested that the genetic distance between E and R is lower

Table 5. Posterior mean (and SD) of the differences of the crosses (Dif.) with the Iberian pig population with the best performance (RE) and the posterior probability (Prob.) of being the best cross under the LMH model for NBA and TNB

	NBA		TNB	
	Dif.	Prob.	Dif.	Prob.
EE	−0.995 (0.196)	0.000	−1.09 (0.202)	0.000
ER	−0.443 (0.123)	0.000	−0.554 (0.125)	0.000
ET	−0.902 (0.175)	0.000	−1.000 (0.178)	0.000
RE	—	0.985	—	0.991
RR	−0.754 (0.189)	0.000	−0.777 (0.193)	0.000
RT	−0.471 (0.178)	0.003	−0.516 (0.183)	0.002
TE	−0.452 (0.164)	0.003	−0.464 (0.169)	0.003
TR	−0.463 (0.198)	0.009	−0.536 (0.203)	0.004
TT	−1.560 (0.226)	0.000	−1.607 (0.232)	0.000

EE = Entrepelado × Entrepelado; ER = Entrepelado × Retinto; ET = Entrepelado × Torbiscal; RE = Retinto × Entrepelado; RR = Retinto × Retinto; RT = Retinto × Torbiscal; TE = Torbiscal × Entrepelado; TR = Torbiscal × Retinto; TT = Torbiscal × Torbiscal; LMH = full model with direct line; maternal and heterosis effects. NBA = number born alive; TNB = total number born.

than the distance between either of them with T. Therefore, we expected a lower degree of heterosis between E and R than in crosses involving T.

The superiority of the LMH model over MH and the posterior distribution of the contrast between direct line effects in the LMH model confirmed the differences in the direct line effects on litter size. Numerous studies have reported differences in reproductive performance among swine populations (Bidanel, 2011); thus, it is not surprising that this variability occurs among the Iberian varieties because the genetic diversity in Iberian pig populations is as high as it is among white pig populations (Martínez *et al.*, 2000; Fabuel *et al.*, 2004). Over centuries, the Iberian pig has evolved adapting to harsh environmental conditions, and producers have subjected populations to ‘empirical’ selection in which adipogenic capacity and morphological criteria have played an important role. In addition, the small size of the herds and the degree of isolation between populations are other important factors. In general, the RR variety had the best line effects for prolificacy, and the E and T varieties had similar performances.

Furthermore, the best fit of LMH compared to the LH model and the posterior distribution of maternal effects provides evidence of its importance. The results indicated clearly the superiority (posterior probability > 0.99) of the maternal effect of the E variety over the R and T varieties, which was supported by the significant differences between reciprocals (RE v. ER and TE v. ET). In quantitative genetics, the importance of maternal effects is well recognized (Wilham, 1972; Wolf and Wade, 2009) and is typically found in growth traits in non-prolific species; however, few studies have reported on the maternal effects of sows on the litter size of their daughters. In addition, the estimates of the maternal effect on NBA and TNB were small (Southwood and Kennedy, 1990; Ferraz and Johnson, 1993; Irgang *et al.*, 1994), which differs

considerably from the results of our study, where the estimates of the differences between the maternal effects of the Dickerson's model of the E variety compared to those of the R and T varieties were significant. Given the information available, the possible cause of that phenomena cannot be disentangled; however, recent studies on the genetic determinism of litter size in prolific species suggest a very complex panorama, which might permit speculation about a possible reason. The influence of maternal care on the reproductive performance of adult females has been investigated in rats (Cameron, 2011), which suggests that maternal behavior is transmitted to the next generation through epigenetic modifications such as methylation and histone acetylation, which is reflected in the variability in the expression of the estrogen receptor alpha. In addition, that phenomena is consistent with recent evidence of the effects of DNA methylation (Hwang *et al.*, 2017) and the presence of imprinted genes (Coster *et al.*, 2012) on litter size in pigs. Furthermore, apparently epistasis is a very important source of genetic variation in litter size in mice (Peripato *et al.*, 2004) and swine (Noguera *et al.*, 2009). Another possibility is the effects of mitochondrial DNA, the variation of which has recently been associated with the maturation of oocytes in bovids (Srirattana *et al.*, 2017) and with the reproductive performance of daughters in pigs (Tsai *et al.* 2016).

Despite the uncertainty about the biological basis of the results of this study, it has identified the clear advantage of crossbred over purebred individuals in litter size traits in the Iberian breed. Therefore, the implementation of a pyramidal scheme to provide crossbred sows to the producers is strongly recommended. Among the varieties investigated, the R population had the greatest direct line effect, and the E population had the greatest maternal effect and can provide a distinct advantage if it is used as the maternal line. The recommended cross for the practical implementation of a pyramidal scheme is R as the paternal and E as the maternal lines, which is supported by the posterior probability of being the best cross for NBA and TNB.

Acknowledgements

The authors gratefully acknowledge INGA FOOD SA and its veterinarians Emilio Magallón, M.J. García-Santana, Lourdes Muñoz, Pilar Díaz and Manuel Ramos for their cooperation with the experimental protocol and their technical support. The work was partially funded by the Center for Industrial Technological Development (CDTI) via grant IDI-20170304 and by grant CGL-2016-80155 from the Ministry of Economy, Industry and Competitiveness (MINECO), Spain. Preliminary results of this study were presented at the World Congress of Genetics Applied to Livestock Production (Noguera *et al.*, 2018).

 L. Varona 0000-0001-6256-5478

Declaration of interest

The authors declare that they have no competing interests.

Ethics statement

The research ethics committee of the Institute of Agrifood Research and Technology (IRTA) approved all of the management and experimental procedures involving live animals, which were performed in accordance with the Spanish Policy of Animal Protection RD1201/05, which complies with the European Union Directive 86/609 for the protection of animals used in experimentation.

Software and data repository resources

None of the data were deposited in an official repository. FORTRAN code is available from the corresponding author under reasonable request.

Supplementary material

To view supplementary material for this article, please visit <https://doi.org/10.1017/S1751731119001125>.

References

- Barea R, Nieto R, Vitari F, Domeneghini C and Aguilera JF 2011. Effects of pig genotype (Iberian v. Landrace x Large White) on nutrient digestibility, relative organ weight and small intestine structure at two stages of growth. *Animal* 5, 547–557.
- Bidanel JP 2011. Biology and genetics of reproduction. In *The genetics of the pig*, 2nd edition (ed. MF Rothschild and A Ruvisky), pp. 218–241. CAB International, Wallingford, UK.
- Boletín Oficial del Estado 2014. Real Decreto 4/2014, de 10 de enero, por el que se aprueba la norma de calidad para la carne, el jamón, la paleta y la caña de lomo ibérico. BOE-A-2014-318.
- Cameron NM 2011. Maternal programming of reproductive function and behavior in the female rat. *Frontiers in Evolutionary Neuroscience* 3, 10.
- Cassady JP, Young LD and Leymaster KA 2002. Heterosis and recombination effects on pig reproductive traits. *Journal of Animal Science* 80, 2303–2315.
- Coster A, Madsen O, Heuven HC, Dibbitts B, Groenen MA, van Arendonk JA and Bovenhuis H 2012. The imprinted gene DIO3 is a candidate gene for litter size in pigs. *PLoS One* 7, e31825.
- Dekkers J, Mather PK and Knol EF 2011. Genetic improvement of the pig. In *The genetics of the pig*, 2nd edition (ed. MF Rothschild and A Ruvisky), pp. 390–425. CAB International, Wallingford, UK.
- Dickerson GE 1969. Experimental approaches in utilizing breed resources. *Animal Breeding Abstracts* 37, 191–202.
- Fabuel E, Barragán C, Silió L, Rodríguez MC and Toro MA 2004. Analysis of genetic diversity and conservation priorities in Iberian pigs based on microsatellite markers. *Heredity* 93, 104–113.
- Fernández A, Rodríguez J, Zuzárrregui J, Rodríguez MC and Silió L 2008. Genetic parameters for litter size and weight at different parities in Iberian pigs. *Spanish Journal of Agricultural Research* 6, 98–106.
- Ferraz JBS and Johnson RK 1993. Animal model estimation of genetic parameters and response to selection for litter size and weight, growth, and backfat in closed seedstock populations of Large White and Landrace swine. *Journal of Animal Science* 71, 850–858.
- Gandemer G 2009. Dry cured ham quality as related to lipid quality of raw material and lipid changes during processing: A review. *Grasas y Aceites* 60, 297–307.
- García-Casco JM, Fernández A, Rodríguez MC and Silió L 2012. Heterosis for litter size and growth in crosses of four strains in Iberian pig. *Livestock Science* 147, 1–8.
- García-Cortés LA and Toro MA 2006. Multibreed analysis by splitting the breeding values. *Genetics Selection Evolution* 38, 601–615.
- Gelfand AE 1996. Model determination using sampling-based methods. In *Markov chain Monte Carlo in practice* (ed. WR Gilks, S Richardson and DJ Spiegelhalter), pp. 145–161. Chapman & Hall, New York, NY, USA.

Gelfand AE and Smith AFM 1990. Sampling-based approaches to calculating marginal densities. *Journal of American Statistical Association* 85, 398–409.

Haley CS, Lee GJ and Ritchie M 1995. Comparative reproductive performance in Meishan and Large White pigs and their crosses. *Animal Science* 2, 259–267.

Hayman BI 1954. The theory and analysis of diallel crosses. *Genetics* 39, 789–809.

Hwang JH, An SM, Kwon S, Park DH, Kim TW, Kang DG, Yu GE, Kim IS, Park HC, Ha J and Kim CW 2017. DNA methylation patterns and gene expression associated with litter size in Berkshire pig placenta. *PLoS ONE* 12, e0184539.

Ibañez-Escriche N, Magallón E, González E, Tejeda JF and Noguera JL 2016. Genetic parameters and crossbreeding effects of fat deposition and fatty acid profiles in Iberian pig lines. *Journal of Animal Science* 94, 28–37.

Irgang R, Favero JA and Kennedy BW 1994. Genetic parameters for litter size of different parities in Duroc, Landrace and Large White sows. *Journal of Animal Science* 72, 2237–2246.

Martínez AM, Delgado JV, Rodero A and Vega-Pla JL 2000. Genetic structure of the Iberian pig breed using microsatellites. *Animal Genetics* 31, 295–301.

Noguera JL, Ibañez-Escriche N, Magallón E, Varona L 2018. Genetic parameters and crossbreeding effects of prolificacy in three commercial lines of Iberian pigs. *Proceedings of the World Congress on Genetics Applied to Livestock Production* 11, 238.

Noguera JL, Rodríguez C, Varona L, Tomás A, Muñoz G, Ramírez O, Barragán C, Arqué M, Bidanel JP, Amills M, Oviló C and Sánchez A 2009. A bi-dimensional genome scan for prolificacy traits in pigs shows the existence of multiple epistatic QTL. *BMC Genomics* 10, 636.

Ogawa S, Kanta A, Kimata M, Ishii K, Uemoto Y and Satoh M 2019. Estimation of genetic parameters for farrowing traits in purebred Landrace and Large White pigs. *Animal Science Journal* 90, 23–28. <https://doi.org/10.1111/asj.13120>.

Pérez-Enciso M and Gianola D 1992. Estimates of genetic parameters for litter size in six strains of Iberian pigs. *Livestock Production Science* 32, 283–293.

Peripato AC, De Brito RA, Mantioli SR, Pletscher LS, Vaughn TT and Cheverud JM 2004. Epistasis affecting litter size in mice. *Journal of Evolutionary Biology* 17, 593–603.

Putz AM, Tiezzi F, Maltecca C, Gray KA and Knauer MT 2015. Variance component estimates for alternative litter size traits in swine. *Journal of Animal Science* 11, 5153–5163.

Quaas RL 1976. Computing the diagonal elements and inverse of a large numerator relationship matrix. *Biometrics* 32, 949–953.

Quinton VM, Wilton JW, Robinson JA and Mathur PK 2006. Economic weights for sow productivity traits in nucleus pig populations. *Livestock Science* 99, 69–77.

Rodríguez C, Rodríguez J and Silio L 1994. Genetic analysis of maternal ability in Iberian pigs. *Journal of Animal Breeding and Genetics* 111, 220–227.

Serra X, Gil F, Pérez-Enciso M, Oliver MA, Vázquez JM, Gispert M, Díaz I, Moreno F, Latorre R and Noguera JL 1998. A comparison of carcass, meat quality and histochemical characteristics of Iberian (Guadyerbas line) and Landrace pigs. *Livestock Production Science* 56, 215–223.

Serrano MP, Valencia DG, Nieto M, Lázaro R and Mateos GG 2008. Influence of sex and terminal sire line on performance and carcass and meat quality of Iberian pigs reared under intensive production systems. *Meat Science* 78, 420–428.

Silio L, Rodríguez C, Rodríguez J and Toro MA 2001. La selección de cerdos ibéricos. In *Porcino Ibérico: Aspectos claves* (ed. C Buxade and A Daza), pp. 125–159. Mundiprensa, Madrid, Spain.

Southwood OI and Kennedy BW 1990. Estimation of direct and maternal genetic variance for litter size in Canadian Yorkshire and Landrace swine using an animal model. *Journal of Animal Science* 68, 1841–1847.

Spiegelhalter DJ, Best NG, Carlin BP and Van der Linde A 2002. Bayesian measures of model complexity and fit (with discussion). *Journal of Royal Statistical Society: Series B Statistical Methodology* 64, 583–639.

Srirattana K, McCosker K, Schatz T and St. John JC 2017. Cattle phenotypes can disguise their maternal ancestry. *BMC Genetics* 18, 59.

Tsai T, Rajasekar S and St. John JC 2016. The relationship between mitochondrial DNA haplotype and the reproductive capacity of domestic pigs (*Sus scrofa domestica*). *BMC Genetics* 17, 67.

Wilham RL 1972. The role of maternal effects in animal breeding: III Biometrical aspects of maternal effects in animals. *Journal of Animal Science* 35, 1288–1293.

Wolf JB and Wade MJ 2009. What are maternal effects (and what they are not)? *Philosophical Transactions of the Royal Society B* 364, 1107–1115.

Appendix : Partial numerator relationship matrices and their generalized inverses

In order to illustrate the procedure of definition of the partial numerator relationship matrices required for the implementation of the procedure of García-Cortés and Toro (2006), we have defined the following pedigree with purebred and crossbred individuals between three populations (A, B and C):

Given this genealogical information, the partial numerator relationship matrix for the A population is:

Individual	Sire	Dam	Population
1	0	0	A
2	0	0	A
3	0	0	B
4	0	0	B
5	0	0	C
6	0	0	C
7	1	2	A
8	1	4	A × B
9	3	4	B
10	1	6	A × C
11	3	6	B × C
12	5	6	C

$$A_A = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0.5 & 0.5 & 0 & 0.5 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0.5 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0.5 & 0.5 & 0 & 0 & 0 & 0 & 1 & 0.25 & 0 & 0.25 & 0 & 0 \\ 0.5 & 0 & 0 & 0 & 0 & 0 & 0.25 & 0.5 & 0 & 0.25 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0.5 & 0 & 0 & 0 & 0 & 0 & 0.25 & 0.25 & 0 & 0.5 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

The inverse of the non-zero part of this matrix (A_A^{-}) can be calculated following the rules described by García-Cortés and Toro (2006), given that $A_A^{-} = (I - P)'D_X^{-}(I - P)$. P is a matrix that relates progeny to parent and D_X is recursively obtained by the algorithm described by Quaas (1976). Therefore:

