

Anatomical and genetic study of an ancient animal tooth showing brachyodont and hypsodont mixed taxonomical characteristics

Short running title: *DNA identification of ancient animal tooth.*

L.V. Monteagudo *, J.A. Obón, A. Whyte, M. T. Tejedor, J. Whyte, A. Cisneros.

Luis Vicente Monteagudo & María Teresa Tejedor: Departamento de Anatomía, Embriología y Genética. Universidad de Zaragoza (Spain).

Jesus Angel Obón, Jaime Whyte & Ana Cisneros: Departamento de Anatomía e Histología Humanas. Universidad de Zaragoza (Spain).

Ana Whyte: Departamento de Patología Animal. Universidad de Zaragoza (Spain).

*Address for correspondence: Luis V. Monteagudo. Departamento de Anatomía, Embriología y Genética Animal. Universidad de Zaragoza. Miguel Servet 177. 50013-ZARAGOZA (Spain).

e-mail: <monteagu@unizar.es>

Abstract

A non human dental piece was found in a Roman Empire tomb dated in the 3rd century A.C. in Zaragoza (Spain). The morphology of this piece showed mixed brachyodont (carnivores) and hypsodont (herbivores) characteristics. As a result, the taxonomical assignation of the piece was impossible. Therefore, a protocol based up in the DNA sequence of the cytochrome c oxidase subunit 1 mitochondrial region (COI) was applied. For this purpose, a pair of primers able to amplify this region in a large variety of animals was designed. The results point to a species of the Genus *Bos* (Family Bovidae). This assignation was later confirmed by the sequencing of a short fragment of the mitochondrial d-loop region. A complete morphological description of the tooth is presented together with the DNA sequence study and comparison protocol.

Keywords: ancient tooth ■ ambiguous taxonomical identification ■ DNA study ■ mitochondrial DNA.

INTRODUCTION

In 1988, a Roman necropolis dated in the 3rd century A.C. was discovered in the city of Zaragoza (Spain) [5]. A non-human dental piece was found in one of the tombs (Fig. 1). As described below, it presented only minimal deteriorations.

The specimen shows mixed brachyodont (short crown, long roots and closed apexes), and hypsodont (no cusps and small crests in the occlusal surface) characteristics. As a consequence, the assignment of the piece to a particular species was unclear, even for experts in animal anatomy.

Therefore, molecular genetics identification was applied, based up on DNA from the internal dentine and on the “DNA Bar Coding” concept (Consortium for the Bar Coding of Life, Smithsonian Institution, 2010, available at <http://www.barcoding.si.edu/DNABarCoding.htm>). DNA Bar coding tries to identify animal specimens mainly by analysing the sequence of the cytochrome c oxidase subunit 1 mitochondrial region (COI). Multiple copies of the mitochondrial genome exist and are protected by the mitochondrial membranes in each cell. Dental pieces provide a resistant support, so that mitochondrial DNA can be recovered from dentine in sufficient amounts for laboratory analysis. The usual approach is based up on consensus Polymerase Chain Reaction (PCR) primers suitable to allow COI amplification and sequencing in a set of species.

We designed PCR primers suitable to amplify and to sequence COI in a wide range of camelids, ruminants and horses. Mitochondrial D-loop analysis was later studied in order to confirm the first results.

CASE REPORT

The tooth presents a short crown. Half-moon shaped small crests are visible in its occlusal surface. Dentine is exposed among the crests. The crown is also crossed by fracture lines running in vestibular to lingual direction. A deeper fracture line follows a mesial to distal direction in the vicinity of the vestibular surface.

The vestibular surface is plain and presents a small defect as a result of a fracture. The mesial and distal surfaces are plain, while the palatal surface is half-moon shaped.

The smallt shows some deterioration in all surfaces (except the mesial one). This results in longitudinal, transversal and oblique fracture lines, especially in the distal surface.

The neck of the tooth is clearly marked. A horizontal fracture line runs its distal and palatal surfaces.

Three independent divergent roots converge into a common radicular trunk close to the neck line.

The mesial-vestibular root is fractured. The distal-vestibular root is conical, has a smooth surface and closed apex. Finally, the palatal root is the largest one. Its vestibular surface is tuberculated in the area surrounding the neck. Its apex is closed.

According to the roots morphology, the tooth is a permanent piece from the left superior dental arch.

The assignment of the piece to a particular the species of origin by means of morphological analysis was unclear, even after consulting with experts in domestic and wild animal anatomy. On one hand, the anatomy of the crown and roots could be compatible with a carnivore or an omnivore (humans, canids, felids, ursids, suids, simians), since it presents a brachyodont aspect due to its short crown, long roots and closed apexes.

However the occlusal surface should point to the opposite assignment. As it presents no cuspids and it shows small crests, a hypsodont ruminant species (bovids, camelids, horses) should be a logical origin for the piece.

MATERIAL AND METHODS

DNA extraction

DNA extraction was performed in the facilities of the Parasite Diseases Section at the Faculty of Veterinary Sciences, in a building different from the one used for PCR and agarose electrophoresis. Mitochondrial DNA from domestic animals had never been studied at this place. The working area was previously irradiated by UV light.

The dentine extraction followed a previously reported protocol [6]. To avoid using cooling systems, the Hedström files were hand operated. For DNA extraction, the dentine was transferred to a 1.5 ml microtube and submitted to two decalcification washes by 0.5 mol/l EDTA (0.5 M pH 8.0), for 5' each. It was then lysed in a buffer containing 10 mM Tris-HCl, 50 mM NaCl, 2% (w/v) sodium dodecylsulfate (SDS) (pH 8.0), and 15 µl of proteinase K (10 mg/ml) at 56 °C for 16 hours.

After spinning, the same volume of Chelex (10% in TE) was added to the supernatant. The mix was incubated at 100°C for 5', vortexed and submitted to 100°C

for 20 minutes. After centrifugation at 13.000 g, the supernatant was used as mould DNA (4 µl/10 µl of final PCR volume).

PCR and sequencing

Consensus PCR primers were designed by means of Primer 3 to amplify a 362 bp segment of COI in a wide range of camelids (Arabian camel, and Bactrian camel), ruminants (cattle, sheep and goat) and horses. The following Genbank entries were used for such purpose: EU159113, EF212038, EF212037, EF507801, EF507800, EF507798, EF507799, AP003423, NC_001941, NC_001640.1, and AF533441. Primers sequences are: Dien-F: 5'-tcgctgattattctcaaccaa-3' and Dien-R: 5'-cctgtacctgctcctgcttc-3'.

Amplification protocol for COI

Different annealing temperatures and Mg⁺⁺ concentrations were checked by using a Multigene Gradient Thermal Cycler (Labnet International Inc.). For purpose of sequencing, we choose the PCR product obtained by using a 3mM final Mg⁺⁺ concentration and the following PCR cycles: 94° C x 5' + 35 x (94°C 30'' + 54.4°C 30''+ 72°C 45'') + 72°C 10'. Studies on COI had never been performed in the laboratory where DNA extraction and PCR amplification were made. After verification in agarose gel, the PCR products were sequenced at the Central Sequencing Service (Universidad de Zaragoza).

Verification by means of mitochondrial d-loop sequencing

Mitochondrial D-loop was partially PCR-amplified by using the primers Ach-F:5'-cctaagactcaaggaagaaactgc -3' and Ach-R: 5'- aacctagagggcattctcactg-3' according to a previously published protocol [1]. The PCR products were sequenced too.

The software Blastn [2] was used to align sequences with those available in the Genbank database.

RESULTS AND DISCUSSION

The odontometric study of the specimen produced the following measurements:

In the crown, the mesiodistal width is 17 mm, while the vestibule-lingual width is 20.05 mm and the distance between the occlusive surface and the cervix is 13 mm.

In the mesiovestibular root, the mesiodistal width is 7 mm, while the vestibule-lingual width is 8.5 mm. Since this root is fractured, its height could not be determined.

In the distovestibular root the mesiodistal width is 5 mm, the vestibule-lingual width is 7 mm and the cervix to apex distance is 20 mm.

In the palatal root the mesiodistal width is 10 mm, the vestibule-lingual width is 12 mm and the cervix to apex distance is 24.5 mm.

In all cases, the diameters were measured at the root bifurcations.

The total height of the specimen from the occlusive surface to the apex is 37.5 mm.

Visual examination of the piece allowed us to disregard a possible human origin, even if the morphological details did not allow us to identify the origin of the specimen.

A reliable 312 bp COI sequence was obtained and deposited at GenBank (Accession number HQ739078). According to Blastn alignment, it is 100% concordant with the sequence of COI in two species: *Bos taurus* (for instance with GenBank entry HM045018.1 and others), and *Bos javanicus* (GenBank entry FJ997262.1). Small differences exist with other *Bos* genus species, such as *Bos primigenius* (309 coincidences, entry GU985279.1), *Bos indicus* (entry AY126697.1, 307 coincidences) and *Bos grunniens* (entry GQ464268.1, 299 coincidences).

Since the alignment with other herbivores' COI sequence (horse, camel, goat, sheep, red deer and roe deer) provides coincidences around 80%, they are disregarded as possible sources of the tooth. The coincidence is similarly low for the Ursidae, Felidae, Suidae and Canidae families.

Only a short reliable 123 bp partial fragment from the d-loop region was obtained (GenBank HQ739079). The species and families tested for COI alignment were also studied for this d-loop segment. Only the genus *Bos* provided high alignment with it. In fact, the Blastn alignment confirmed the bovine origin of the specimen, since the coincidence is 100% with *Bos taurus* (entry AY700556.1 among others), over 99% with *Bos grunniens* (entry AY428641.1 among others), and with *Bos indicus* (FJ800951.1 among others), and 98% with *Bos primigenius* (entry GU434123.1 among others).

Taxa different from *Bos* provided only very little alignment coincidences, usually in short fragments.

In all, the analysis of the DNA sequences obtained in this work points to the genus *Bos* as the most suitable origin for this dental piece.

Among the possible extinct *Bos* species, complete mitochondrial DNA sequence information is only available for the auroch (*Bos primigenius*). Comparison with all the

bovine species (present and extinct) is thus impossible. Moreover, the phylogeny of the *Bos* taxon is still under revision [3].

Different factors affect the morphology and size of teeth. Despite the extinction of many cattle breeds, a wide range of body sizes is still observed. The adult body weight of cows from many local breeds conserved all over the old world is below 300 kg. In the past, such limited weight could have been a very common characteristic [7].

ACKNOWLEDGMENTS

The authors thanksgive Dr Joaquin Quilez for the use of the facilities of the Parasitology laboratory and Dr Catarina Ginja for suggestions on the PCR amplifications.

Conflict of interest

None

References

1. Achilli A, Olivieri A, Pellecchia M, Ubaldi C, Colli L, Al-Zahery N, Accetturo M, Pala M, Hooshiar Kashani B, Perego UA, Battaglia V, Fornarino S, Kalamati J, Houshmand M, Negrini R, Semino O, Richards M, Macaulay V, Ferretti L, Bandelt HJ, Ajmone-Marsan P, Torroni A (2008) Mitochondrial genomes of extinct aurochs survive in domestic cattle. *Current Biol*, 18: R157-R158. Available at < <http://www.sciencedirect.com/science/article/pii/S0960982208000651>>
2. Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res*, 25 : 3389-3402. Available at <<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC146917/>>
3. Bibi F, Vrba ES (2010) Unraveling bovin phylogeny: accomplishments and challenges. *BMC Biology*, 8, 50. Available at < <http://www.biomedcentral.com/1741-7007/8/50>>

4. Consortium for the Bar Coding of Life Smithsonian Institution (consulted November 27th 2012.) Available at <http://www.barcoding.si.edu/DNABarCoding.htm>

5. Galve MP, Baena S, Obón JA, Nieto JL (2009) La necrópolis occidental de Caesaraugusta en el siglo III (Calle Predicadores, 20-30, Zaragoza) Prensas Universitarias de Zaragoza, Ayuntamiento de Zaragoza, Institución Fernando el Católico. Zaragoza, Spain. ISBN 978-84-9251-52-4.

6. Malaver PC, Yunis JJ (2003) Different Dental Tissues as Source of DNA for Human Identification in Forensic Cases. CMJ (Forensics Sciences), 44: 306-309. Available at < <http://www.cmj.hr/2003/44/3/12808723.pdf>>

7. Oklahoma State University Livestock Project. Available at <http://www.ansi.okstate.edu/breeds/cattle/> (consulted November 27th 2012).

Figure legends:

Figure 1. a: Occlusal surface presenting small crests (***) and two lines of fracture: vestibular-lingual (thin arrows) and mesial-distal (thick arrows). b: The mesial-vestibular, the distal-vestibular and the palatal root (MV, DV and P, respectively). All of them are conical shape and present closed apexes. Noticeably, the mesial-vestibular root is fractured. Several fracture lines are observed in the palatal surface (arrows).

