

Review of the Apparent Molar Heat Capacities of NaCl(aq), HCl(aq), and NaOH(aq) and Their Representation Using the Pitzer Model at Temperatures from 298.15 K to 493.15 K

Joan J. Manyà^{a,b} and Michael Jerry Antal, Jr.^c

*Hawaii Natural Energy Institute, School of Ocean and Earth Science and Technology,
University of Hawaii at Manoa, Honolulu, Hawaii 96822 and Thermo-chemical
Processes Group (GPT), Aragón Institute of Engineering Research (I3A), University of
Zaragoza, Maria de Luna 3, E-50018 Zaragoza, Spain.*

^a To whom correspondence should be addressed. E-mail address: joanjoma@unizar.es
Fax: + 34 976761879

^b Aragón Institute of Engineering Research (I3A), University of Zaragoza

^c University of Hawaii at Manoa

Abstract

In this study, a new estimation of the standard-state partial molar heat capacity (C_p^0) of three binary systems [NaCl(aq), HCl(aq), and NaOH(aq)] for temperatures ranging from 298.15 K to 493.15 K is performed. From experimental data (available in the literature to date) corresponding to the apparent molar heat capacities of the above-mentioned electrolytes, a procedure based on the assumption of the Pitzer's ion interaction model was applied in order to calculate these important C_p^0 values at a given temperature. Results obtained for each electrolyte (or ion) were correlated as functions of temperature by commonly-used polynomial equations. Use of the hydrogen ion convention enabled estimates of the temperature-dependent values of C_p^0 of the individual ions $\text{Na}^+(\text{aq})$, $\text{Cl}^-(\text{aq})$, and $\text{OH}^-(\text{aq})$. When used to calculate the temperature dependent values of the Gibbs free energy of formation of $\text{OH}^-(\text{aq})$, the formula for $C_p^0(T)$ of $\text{OH}^-(\text{aq})$ given herein provides good agreement with values derived from independent measurements of the ion constant of water at elevated temperatures. In the specific case of NaCl, a considerable disagreement was observed between the estimations performed using the Archer's model (Archer, D. G. *J. Phys. Chem. Ref. Data* **1992**, 793-829) and some experimental data published years later. This fact can be explained, among other factors, by the dependence of the model parameters on the experimental uncertainty and systematic errors, which can be incurred during calorimetric experiments. For this reason, continuous revisions and updating of the proposed correlations are required when new experimental data become available.

Keywords

Standard-State Partial Molar Heat Capacity; Single Electrolyte; Pitzer's Model; Hydrogen Ion Convention.

Introduction

Knowledge of heat capacities of pure substances is critical in order to perform a wide variety of thermodynamic calculations, many of which involve the effects of changing temperature on other thermodynamic properties such as Gibbs energies and related equilibrium constants, enthalpies, and entropies.

In the present study, a new estimation of the standard-state partial molar heat capacity (Cp^θ) of three binary systems [NaCl(aq), HCl(aq), and NaOH(aq)] for temperatures ranging from 298.15 K to 493.15 K is presented. The Pitzer ion-interaction model has been used to fit the previously published data concerning the apparent molar heat capacities for these aqueous electrolytes.

Assuming the hydrogen ion convention, which states that the conventional standard molar Gibbs energy of formation, entropy and heat capacity of H^+ are all set equal to zero at all temperatures¹; the standard-state partial molar heat capacity of Na^+ (aq), $Cl^-(aq)$, and $OH^-(aq)$ can be estimated from the apparent heat capacity values of the three analyzed electrolytes.

The main experimental data available in the literature for these solutions has been measured by means of mass-flow heat capacity calorimeters. In this type of instrument, convection heat transfer from the calorimetric tubing and the heater to the surroundings has been generally recognized as a principal source of systematic error in the measurements.² This experimental uncertainty motivates the performance of more reliable experimental studies which should be continuously collected and published. In addition, the mathematical procedures applied to the experimental data are gradually improving as a consequence of the use of more sophisticated algorithms and software applications. In our opinion, all of these reasons justify the development of the present study.

Theoretical Background

The apparent molar heat capacity (Cp^ϕ) corresponds to the “excess” heat capacity (per mol of solute) with respect to the pure solvent of the solution. According to several researchers³⁻⁷, the apparent molar heat capacity (in $J mol^{-1} K^{-1}$) for binary systems can be written as a function of experimental heat capacity values as follows:

$$Cp^\phi = M Cp^{exp} + \left[\frac{1000(Cp^{exp} - Cp_w^0)}{m} \right] \quad (1)$$

where M is the molecular weight of the solute, m is the molality of the solution, Cp^{exp} is the specific heat capacity of the solution (experimental value), and Cp_w^0 is the specific heat capacity of pure water, respectively. It can be deduced from the second term in equation 1 that the precision and accuracy of Cp^θ is particularly sensitive to the difference $(Cp^{exp} - Cp_w^0)$, and that this difference is amplified increasingly as m decreases because of the factor $(1/m)$.

The Pitzer's ion interaction approach⁸ has been used with success to represent the properties of both single and multicomponent electrolyte solutions from low to high ionic strengths.³ The basic equation of the ion interaction approach considers the excess free energy to be the sum of long-range electrostatic interactions (given by an extended Debye-Hückel theory) and specific short-range binary and ternary interactions between ions.⁸ For heat capacities and for 1-1 electrolytes; the Pitzer's formalism takes the following form:

$$Cp^\theta = Cp^0 + \nu |z_M z_X| A_J (2b)^{-1} \ln(1 + bI^{1/2}) - 2\nu_M \nu_X RT^2 m \frac{[\beta_{MX}^{(0)J} + 2\beta_{MX}^{(1)J} [1 - (1 + \alpha I^{1/2})] \exp(-\alpha I^{1/2})]}{\alpha^2 I} - 2\nu_M \nu_X RT^2 m^2 C_{MX}^J \quad (2)$$

where ν_M and ν_X denote the stoichiometric coefficients of cations and anions, respectively ($\nu = \nu_M + \nu_X$); I is the molality-based ionic strength defined by equation 3 and z_i are the formal charges of the ions. The parameters α and b are arbitrary constants in the Pitzer equation and are assigned the values of 2.0 and 1.2 kg^{1/2} mol^{1/2}, respectively (for 1-1, 1-2, and 2-1 electrolytes).⁴

$$I = \frac{1}{2} \sum_i m_i z_i^2 \quad (3)$$

In equation 2, A_J (in J kg^{1/2} K⁻¹ mol^{-3/2}) is the molality-based Debye-Hückel coefficient for heat capacities. The value of this coefficient as a function of both temperature and pressure depends upon the dielectric constant and density of water. There are several published works in which the values of A_J are reported^{9, 10} and they are in relatively good agreement (within 3%).

It should be noted that α and b are assumed to be temperature independent. In this way, at fixed pressure and temperature, the apparent molar heat capacity is expressed in terms of four adjustable (temperature and pressure dependent) parameters per electrolyte: Cp^0 , $\beta_{MX}^{(0)J}$, $\beta_{MX}^{(1)J}$, and C_{MX}^J ; which are determined by fitting the equations to experimental Cp^θ data. The extrapolation of the model to infinite dilution corresponds to

the parameter Cp^0 (the standard-state partial molar heat capacity of the aqueous electrolyte).

As an alternative to the Pitzer approach, the more simplified Redlich-Rosenfeld-Meyer (RRM) type equation could be adequate for correlating experimental Cp^θ values up to an ionic strength of approximately 1 mol kg⁻¹.¹¹ For heat capacities, the RRM model can be written as follows:¹²

$$Cp^\theta = Cp^0 + (w)^{3/2} A_j^{1/2} (m)^{1/2} + B_c m \quad (4)$$

where w is a valence factor given by equation 5 and B_c is the only parameter fitted from the data and corresponds to an empirical coefficient which is function of both pressure and temperature.

$$w = \frac{1}{2} (v_M z_M^2 + v_X z_X^2) \quad (5)$$

Procedure Description

In this section, the procedures which we employed to estimate the heat capacity values are described. Special attention is focused on the interpolation of the published values of the molality-based Debye-Hückel coefficient for heat capacities (A_j), the description of nonlinear least squares technique adopted to fit the experimental values of apparent molar heat capacity, and the statistical interpretation of the fitting results.

A_j values

Recent values of the Debye-Hückel coefficients at selected temperatures and pressures were reported by Fernandez et al.¹⁰ in 1997. Nevertheless, only values at 300, 350, 400, and 500 K and at 0.1, 10, 100, and 1000 MPa are available. This fact implies that, in most cases, the A_j value for a given temperature and pressure needs to be calculated according to the equations of Bradley and Pitzer.⁹ The precision of this calculation can affect the reliability of the determined A_j value. Unfortunately, no details concerning the exact determination of A_j were reported in previous works (see, for instance, studies of Hovey et al.³, Saluja et al.⁴, and Magalhaes et al.¹¹).

As an alternative to the direct (and arduous) calculation of A_j , we have developed an interpolation procedure to estimate the values of the Debye-Hückel coefficients from the data reported by Fernandez and coworkers.¹⁰ This procedure involves two stages. In the first-one, the reported A_j values at constant temperature were fitted as a function of pressure by an exponential equation:

$$\frac{A_J}{R} = a_0 + a_1 \exp(a_2 p) \quad (6)$$

where p is the pressure (in MPa), R is the universal gas constant ($8.3145 \text{ J mol}^{-1} \text{ K}^{-1}$), and a_0 , a_1 , and a_2 correspond to the adjustable parameters. This function type, which does not have a physical basis, was chosen among several mathematical expressions because it provides a reasonable representation of the behavior of the data. Table 1 shows the fitting results obtained for each temperature.

R^2_{adj} corresponds to the adjusted regression coefficient (the coefficient of determination, R^2 , adjusted by the degree of freedom¹³). In light of the results reported in Table 1, the agreement between data points and the exponential function was excellent.

The second stage of the interpolation procedure involves modeling of the A_J values at a given pressure value as a function of temperature. At any given pressure, the values of A_J/R at 300, 350, 400, 450, and 500 K are determined by the exponential equations (the parameters of which are reported in Table 1). Finally, these values of A_J/R at constant pressure were fitted by a polynomial function of fourth order in temperature. Using this procedure we obtained excellent agreement between data points and the polynomial function for all cases involved in the present work. An example of this approach for a given pressure of 4.0 MPa is shown in Figure 1.

Least squares curve fitting

The experimental apparent heat capacity values at a given temperature and as a function of molality were fitted by the Pitzer model (see equation 2) using the Levenberg-Marquardt (L-M) algorithm.¹⁴ The Origin Pro v. 8 software was used for this purpose. The goal of the fitting process is to estimate the parameter values which best describe the data. The standard way of finding the best fit is to choose the parameters that would minimize the deviations of the theoretical curve from the experimental points (chi-square minimization). As starting points of the iteration process, the Pitzer parameter values reported by Criss and Millero¹⁵ for several aqueous electrolytes at 298.15 K were selected.

The solution given by the software was analyzed from a statistical point of view. Every fitting process was submitted to an F -test to check the ability of the model. The F -ratio quantifies the relationship between the relative increase in sum-of-squares and the relative increase in degrees of freedom. For a confidence interval of 95%, probability values lower than 0.05 indicate that the null hypothesis can be rejected and,

consequently, the model is significant. In addition, a *t-test* was also performed for each individual parameter (Cp^0 , $\beta_{MX}^{(0)J}$, $\beta_{MX}^{(1)J}$, and C_{MX}^J) to assess its significance for a confidence interval of 95%.

The results of the two tests and the R^2_{adj} values should provide sufficient information about the ability of the Pitzer approach to fit the experimental data for each molality and temperature value.

Results and Discussion

In this section, the results obtained applying the Pitzer approach to previous published data sets for each electrolyte are presented.

NaCl(aq)

Table 2 summarizes the experimental data sources for apparent molar heat capacities of NaCl(aq). All of the experiments were performed using flow calorimeters that have revolutionized the measurement of heat capacities of fluids. Nevertheless, concerns have been raised about the accuracy of this kind of calorimeter because of possible heat losses.¹¹ Several attempts have been made to quantify these losses (e.g., Desnoyers et al.¹⁶), but the results have been inconclusive.

The values of the apparent heat capacities reported by Saluja et al.⁴ differ in a significant way from those calculated using the model of Archer² (differences ranging from 1.35% at 298 K and $m = 0.1076$ to 17.29% at 373 K and $m = 5.9477$). Archer's model was developed in 1992 assuming a revised Pitzer equation (with more than 50 adjustable parameters) and fitting experimental results available at that time. Nevertheless, the more recent experimental data of Saluja and coworkers do not agree with Archer's model probably because of the excessive dependence of the adjusted parameters to the experimental uncertainty of data points evaluated by Archer.²

An example of application of the Pitzer model to fit a single given data set is presented in Figure 2, that displays the satisfactory fit of the Pitzer model to the apparent molar heat capacity as a function of molality for data reported by Saluja et al.⁴ at 348.15 K.

A summary of the results obtained from the fitting process is given in Table 3. The data corresponding to the work of Perron and coworkers¹⁹ are not reported in Table 3 as a consequence of the poor agreement between Pitzer model and their experimental data. This fact could be due to the inaccuracy of the experimental measurements performed

by Perron et al.¹⁹ The last sentence is based on the comparison of the Cp^0 values obtained by Perron et al.¹⁹ and Tanner and Lamb¹⁸ at a temperature value of 318.15 K (see Figure 3b). This inconsistency between experimental data points from different sources is not present when data from Tanner and Lamb¹⁸ and Saluja et al.⁴ are compared (see Figure 3a). The parameters not reported correspond to the cases in which the $prob(t)$ value obtained for each individual parameter was higher than 0.05 (not significant parameters assuming a confidence interval of 95%). It is interesting to note that the model was significant in all cases. Only the results obtained from experiments reported by Smith-Magowan and Wood in 1981¹⁷ exhibit p -values ($Prob. > F$) near to 0.05.

It is well-known that the Pitzer model is adequate for correlating experimental Cp^0 values up to relatively high ionic strengths values.¹¹ However, an additional estimation of the Pitzer parameters was performed using the experiments reported by Saluja et al.⁴ and performed at relatively low molalities (up to 1 mol kg⁻¹). Table 4 reports the results obtained for the Cp^0 parameter using only the low-molality data reported by Saluja and coworkers. The results obtained are relatively in agreement with the values reported in Table 3 and, for this reason, the ability of the Pitzer model to correlate experimental NaCl data up to approximately 6 mol kg⁻¹ is reconfirmed here.

The ability of the alternative approach, the RRM type equation, has also been checked in this work for the apparent heat capacity values of the NaCl(aq) reported by Saluja et al.⁴ at low molalities. The correlation results are given in Table 5. The RRM model seems to be appropriate in light of the correlating results shown in Table 5. This model can be an interesting option in order to obtain a preliminary Cp^0 value when the number of data points available is less than 5 (minimal data points required to perform a correlation by means of the Pitzer formalism).

Figure 4 shows the evolution of the Cp^0 values reported in Table 3 as a function of temperature. The apparent disagreement of the values calculated from the data of Smith-Magowan and Wood¹⁷ could be due to both the relatively poor fit of the experimental data by the Pitzer model and the additional experimental uncertainty related to the operation at high pressures (see Table 2). The values displayed in Figure 4 were submitted to a fitting process using several polynomial functions assuming as negligible the pressure effects. This assumption may compromise the accuracy of the procedure. The effect of pressure on the apparent molar heat capacity can be expressed by the following equation:

$$\left(\frac{\partial Cp^\phi}{\partial p}\right)_T = -T\left(\frac{\partial^2 V^\phi}{\partial T^2}\right)_p \quad (7)$$

The problem of solving this equation is the difficulty of determining second derivatives from experimental measurements. From the literature, there are clearly large differences in precision of the volumetric data sets, as well as in data coverage of temperature, pressure, and concentration, and these proved to be troublesome in evaluating an overall fit of the volumetric data.²⁰ In addition to this, it should be kept in mind that one principal objective of estimating the standard-state heat capacity (at a given temperature value) is the goal of calculating the heat capacity change for reactions involving aqueous ions. In this sense, Puigdomenech and coworkers¹ considered that pressure effects can be neglected for temperatures lower than 573 K

From data displayed in Figure 4, the apparent best results ($R^2_{adj} = 0.9914$) were obtained with the following equation:

$$Cp^0 (NaCl) = -5.90873760962963 \cdot 10^4 + \frac{3.4028457831 \cdot 10^6}{T} + 3.764418314936 \cdot 10^2 T - 1.1379818529 T^2 + 1.6607597 \cdot 10^{-3} T^3 - 9.4980168897 \cdot 10^{-7} T^4 \quad (8)$$

The black solid line in Figure 4 corresponds to this equation. Obviously, the usefulness of equation 8 is limited to the applied temperature range. Extrapolations of this function will not provide reliable solutions. Figure 4 also shows the pressure-dependent Cp^0 values estimated using the revised Helgeson-Kirkham-Flowers (HKF) equations of state for aqueous species.²¹⁻²³ All of the calculations reported here (related to the revised HKF model) have been performed using the CHNOSZ software package (which can be freely downloaded from the project website at <http://www.chnosz.net>), which has been developed at U. C. Berkeley for the R software environment.²⁴ For the Cp^0 estimations made in the present study, the thermodynamic properties of pure water were calculated using an implementation in R code of the IAPWS-95 formulation.²⁵ The electrostatic properties of the pure solvent were calculated using the equations of Archer and Wang.²⁶

Figure 4 indicates that the values calculated using the HKF model at 17.7 MPa agree reasonably well with the Cp^0 parameters obtained from the experimental data reported by Smith-Magowan and Wood¹⁷ at the same pressure. This fact confirms the effect of the pressure on the value of standard-state partial molar heat capacity. Nevertheless, it is also noted that equation 8 predicts in a reasonable way the Cp^0 values when the effect of pressure is not considered (when several data points measured at different pressures are

combined). According to Puigdomenech and coworkers¹, the measurement of aqueous solution heat capacities should be given a high priority, especially at high temperature, in order to validate the parameters of the equation of state-based models.

HCl(aq)

The same above-mentioned procedure was applied in order to estimate the temperature dependence of the standard-state partial molar heat capacity values (Cp^0) for the aqueous hydrochloric acid system. Table 6 summarizes the experimental data sources for apparent molar heat capacities of HCl(aq). It is interesting to note that results reported by Patterson and coworkers²⁷ were obtained using a scanning calorimeter (not a flow calorimeter). The results of the fitting process using the Pitzer model are reported in Table 7. Some results obtained from the data published by Tremaine et al.²⁸ and Allred and Woolley²⁹ are not reported in Table 7 as a consequence of the bad fitting results obtained. The experimental uncertainty could be higher in the data reported in relatively old studies because of the systematic errors that have been identified in subsequent studies.^{2,11}

The results reported in Table 7 concerning the steady-state heat capacity values (Cp^0) are displayed in Figure 5. We must remark that results obtained from Patterson and coworkers²⁷ are in relatively good agreement with those obtained from other researchers who used Picker flow calorimeters. The values displayed in Figure 5 were submitted to a fitting process using several polynomial functions assuming as negligible the pressure effects. The best results ($R^2_{adj} = 0.9986$) were obtained using the following equation:

$$Cp^0(HCl) = 1.773367390171106 \cdot 10^5 - \frac{1.3758183265 \cdot 10^{-7}}{T} - 9.097252470855 \cdot 10^2 \cdot T + \quad (9)$$

$$2.315748398 \cdot T^2 - 2.9169397 \cdot 10^{-3} \cdot T^3 + 1.4503729301 \cdot 10^{-6} \cdot T^4$$

The black solid line in Figure 5 corresponds to equation 9. In this case, the revised HKF model values depart from the corresponding Cp^0 values obtained from the experimental data reported by Sharygin and Wood³⁰ at 28 MPa. This fact emphasizes the need for additional experimental data to refine the current HKF parameters.

At this point, the steady-state heat capacity of individual ions (Na^+ and Cl^-) at a given temperature can be calculated assuming the hydrogen ion convention:

$$Cp^0(Cl^-) = Cp^0(HCl) \quad (10)$$

$$Cp^0(Na^+) = Cp^0(NaCl) - Cp^0(Cl^-) \quad (11)$$

The values calculated from equation 11 were then fitted by means of the equation 12 with a relatively high degree of precision ($R^2_{adj} = 0.9210$; $R^2 = 0.9895$).

$$Cp^0 (Na^+) = -5.42899712329456710^5 + \frac{4.023267891510^{-7}}{T} + 2.901241692242910^3 T - 7.6728859593 T^2 + 1.0044970910^{-2} T^3 - 5.209547720410^{-6} T^4 \quad (12)$$

It is interesting to compare the average Cp^0 values (average value between the Cp^0 at 298.15 K and the Cp^0 at a given temperature value) calculated using equations 10-11 with those reported by Barner and Scheuerman³¹ and determined assuming the “entropy correspondence principle” enunciated by Criss and Cobble.³² These authors observed that the “absolute” entropies of many aqueous ions at elevated temperatures could be correlated with their “absolute” entropies at 298.15 K, provided that the entropy of $H^+(aq)$ could be assigned a temperature-dependent, nonzero “absolute” reference value. Figure 6 shows this comparison for both $Na^+(aq)$ and $Cl^-(aq)$ ions. The large differences observed for the average heat capacity values for individual ions could be due, among other factors, to the inherent assumptions of the two approaches: the “hydrogen ion convention” assumed in the present work and the “entropy correspondence principle” adopted by Criss and Cobble.³² The relatively good agreement observed for the aqueous NaCl electrolyte at low temperatures could be an indicator that the differences for individual ions can be explained by the assumption of different conventions. Nevertheless, the larger disagreement observed when the temperature increases could be related to the fact that the approach developed by Criss and Cobble is less accurate at temperatures higher than approximately 400 K.

NaOH(aq)

For the aqueous NaOH electrolyte Table 8 summarizes the experimental data sources for apparent molar heat capacities which have been considered in this work. Table 9 reports the results obtained for the Pitzer model parameters.

The apparent heat capacities at 298.15 K reported by Magalhaes et al.¹¹, Patterson et al.²⁷, and Allred and Woolley²⁹ are compared in Figure 7. Results obtained by Allred and Woolley are clearly in disagreement with those reported by other authors. This fact could be explained by the higher systematic errors inherent to the calorimetric measurements performed by Allred and Woolley in 1981. As has been mentioned above, the accuracy of a Picker flow calorimeter depends to a great extent on the correct quantification of the heat losses. It is reasonable to think that this correction process has been improved in recent years and, for this reason, the most recent experiments could provide the most reliable results. In any case, the standard-state partial molar heat capacity (Cp^0) reported in Table 9 and calculated from the data set reported by Allred

and Woolley²⁹ ($-95.16 \text{ J mol}^{-1} \text{ K}^{-1}$) is not in serious disagreement with the same parameter estimated from the data of Magalhaes et al.¹¹ ($-99.12 \text{ J mol}^{-1} \text{ K}^{-1}$).

In addition, Figure 7 also shows that the data published by Patterson et al.²⁷ and experimentally obtained using a fixed-cell scanning calorimeter (equipped with a constant-volume cell) are in excellent agreement with those reported by Magalhaes et al.¹¹ This fact could confirm the ability of this type of calorimeter to provide reliable heat capacity results for electrolyte solutions. However, the calorimeter used by Patterson and coworkers was calibrated³⁵ assuming the values reported by Archer² for the aqueous NaCl electrolyte and estimated from experimental data obtained using Picker flow calorimeters. Unfortunately, no additional calibrations with other liquid standards (as suggested by Cerdeirina et al.³⁶) were performed in order to check the accuracy of the instrument. For this reason, we believe that the high similarity between the results obtained from the two types of calorimeters is expected.

The results reported in Table 9 concerning the standard-state heat capacity values (C_p^0) for the aqueous NaOH electrolyte are displayed in Figure 8. The black solid line in the figure corresponds to the fitted curve according to equation 13. The fit quality was excellent ($R^2_{adj} = 0.9999$).

$$C_p^0(\text{NaOH}) = -4.19505400752324 \cdot 10^4 + \frac{1.9615565652 \cdot 10^6}{T} + 2.959385831059 \cdot 10^2 T - 0.95475222715 T^2 + 1.4644429 \cdot 10^{-3} T^3 - 8.7369274529 \cdot 10^{-7} T^4 \quad (13)$$

From analyzing Figure 8, it can be deduced that, at temperatures higher than 400 K, the revised HKF model values significantly depart from the corresponding C_p^0 values.

The standard-state heat capacity of the individual aqueous OH^- ion as a function of temperature can be calculated assuming the hydrogen ion convention and the heat capacity values for $\text{Na}^+(\text{aq})$ estimated in the present study:

$$C_p^0(\text{OH}^-) = C_p^0(\text{NaOH}) - C_p^0(\text{Na}^+) \quad (14)$$

The results provided by equation 14 are displayed in Figure 9. The evolution of the C_p^0 values as a function of temperature can be reproduced by the following equation ($R^2_{adj} = 1.0000$):

$$C_p^0(\text{OH}^-) = -3.79062897306247 \cdot 10^4 + \frac{2.0095760617 \cdot 10^6}{T} + 2.539848770523 \cdot 10^2 T - 0.8025127175 T^2 + 1.22030464 \cdot 10^{-3} T^3 - 7.306388889 \cdot 10^{-7} T^4 \quad (15)$$

The comparison between the average C_p^0 values estimated here and those reported by Barner and Scheuerman³¹ has also been performed for both $\text{OH}^-(\text{aq})$ and $\text{NaOH}(\text{aq})$. Figure 10 shows how the comparative trends are similar to those obtained for aqueous

NaCl electrolyte: a significant disagreement is observed for the individual ion Cp^0 values (probably as a consequence of the different reference values adopted by the two procedures) and a relatively good agreement (but only at low temperatures) is observed for the electrolyte results.

Water ionization

The ionization constant of water (K_w) is a benchmark property in aqueous solution chemistry and has been experimentally obtained over wide ranges of temperature and pressure. Based on these available experimental measurements, several authors have developed models to predict the pK_w value as a function of both temperature and density.³⁷⁻⁴¹ Recently, Bandura and Lvov⁴² reported a new model (based on a semitheoretical approach in which statistical thermodynamics is employed) to successfully fit the experimental data. From these pK_w values, we can estimate $\Delta_f G^0$ value for $\text{OH}^-(\text{aq})$ at a given temperature, according to the following relationship:

$$\Delta_f G^0_{\text{OH}^-} = -RT \ln K_w + \Delta_f G^0_{\text{H}_2\text{O}} \quad (16)$$

with $\Delta_f G^0_{\text{H}^+} \equiv 0$ (hydrogen ion convention) and $\Delta_f G^0_{\text{H}_2\text{O}}$ being given by Wagner and Pruss.²⁵

On the other hand, the $\Delta_f G^0_{\text{OH}^-}$ values can also be estimated according to the second-law extrapolation:

$$\Delta_r G^0(T) = \Delta_r G^0(298.15) - (T - 298.15)\Delta_r S^0(298.15) + \int_{298.15}^T \Delta_r C_p^0 dT - T \int_{298.15}^T \frac{\Delta_r C_p^0}{T} dT \quad (17)$$

$$\Delta_f G^0_{\text{OH}^-}(T) = \Delta_r G^0(T) - \Delta_f G^0_{\text{H}_2\text{O}}(T) \quad (18)$$


Equation 17 can be rewritten assuming an average value of $\Delta_r C_p^0$ for the range of 298.15– T :

$$\Delta_r G^0(T) = \Delta_r G^0(298.15) - (T - 298.15)\Delta_r S^0(298.15) + \Delta_r C_p^0 \left[T - 298.15 - T \ln \left(\frac{T}{298.15} \right) \right] \quad (19)$$

A way to check the reliability of the above-reported Cp^0 values for $\text{OH}^-(\text{aq})$ is to compare the temperature-dependent values of $\Delta_f G^0_{\text{OH}^-}$ calculated from (a) equation 16 using the pK_w values reported by Bandura and Lvov⁴², (b) equation 17 or 19 and equation 18 using the Cp^0 values for $\text{OH}^-(\text{aq})$ provided by equation 15, and (c) equations 18 and 19 using the average heat capacities reported by Barner and Scheuerman.³¹ The data corresponding to the Cp^0 values for the ordinary water have been obtained from the IAPWS-95 formulation²⁵ at 0.1 MPa and saturation pressure (for

temperatures higher than 273.15 K). Table 10 summarizes the results of the comparison process and shows the good agreement of $\Delta_f G^0_{OH^-}$ given by Barner and Scheuerman with those calculated from the ionization constant of water reported by Bandura and Lvov. Likewise, values of $\Delta_f G^0_{OH^-}$ calculated using equations 18 and 19 with C_p^0 values for $OH^-(aq)$ provided by equation 15 agree well with those calculated from the ionization constant of water reported by Bandura and Lvov; whereas the use of the integral form (equation 17) offers poor agreement above 373 K. Special attention should be focused on analyzing these unexpected discrepancies in future studies.

In addition to this, we can also estimate the standard (infinite dilution) molar heat capacity change for the ionization of water, $\Delta_r C_p^0$, from the standard partial molar heat capacities of $NaCl(aq)$, $NaOH(aq)$ and $HCl(aq)$ (given by equations 8, 9, and 13, respectively) by assuming electrolyte additivity:



$$\Delta_r C_p^0(T) = C_{pNaOH(aq)}^0(T) + C_{pHCl(aq)}^0(T) - C_{pNaCl(aq)}^0(T) - C_{pH_2O}^0(T) \quad (20)$$

The data corresponding to the C_p^0 values for the ordinary water have been obtained from the IAPWS-95 formulation²⁵ at 10 MPa. Table 11 reports the results of these calculations and shows the relatively good agreement between the $\Delta_r C_p^0$ values calculated from the two equilibrium approaches. This fact seems to confirm the accuracy of estimations of C_p^0 for all of electrolytes analyzed here. Table 11 also shows that results obtained from the present study are in agreement (especially at temperatures below 373.15 K) with those recently reported by Schrödle and coworkers.⁴³ Exact agreement at higher temperatures would not be expected because of the uncertainties that inevitably arise from the different procedures used in the extrapolation to infinite dilution. In addition to this, the results obtained for the $\Delta_r C_p^0$ values by using the revised HKF equations show a significant disagreement with the values obtained using equations 8, 9, 13, and 20 (see Table 11). This discrepancy increases at high temperatures as the differences among C_p^0 predicted values increase, especially for $HCl(aq)$ and $NaOH(aq)$ (see Figures 5 and 8).

Conclusions

In the present work, the experimental data available in literature corresponding to the apparent molar heat capacity, Cp^o , of three aqueous electrolytes (NaCl, HCl, and NaOH) were fitted using a Pitzer's ion-interaction model. As a result of this mathematical approach, the standard-state heat capacity values have been estimated as a function of temperature for each electrolyte.

The new estimated Cp^o values for both molecules and ions have been correlated as a function of temperature (from 298.15 K to approximately 493.15 K) by means of relatively simple equations (assuming the pressure effects as negligible). The use of these mathematical expressions can be very useful in order to perform thermodynamic calculations involving temperature changes.

In addition, the average Cp^o values estimated in the present study for NaCl(aq) and NaOH(aq) are clearly in disagreement (especially at temperatures higher than 400 K) with those calculated assuming the ion "absolute" entropy values reported by Barner and Scheuerman in 1973. In our opinion, this finding is very important because many thermodynamic studies were based on the assumption of the values available in the Barner and Scheuerman's handbook. Further comparative analyses would be interesting to confirm this observed trend.

In the specific case of NaCl, a significant disagreement was observed between experimental data reported by Saluja and coworkers⁴ in 1995 and the estimations performed using the Archer's model² (in which the parameters were adjusted from experimental data points reported before 1992). This fact can be related, among other factors, to the critical dependence of the model parameters on the experimental uncertainty. For this reason, continuous updates of the proposed models are required when new experimental data becomes available. The experimental uncertainty is related to the systematic errors that can be introduced during the calorimetric measurements. In this sense, it would be interesting to perform further studies using both flow and differential scanning calorimeters in order to gain a better understanding of the peculiarities of each type of instrument. Probably the performance of a round robin study for a given pure liquid substance will be needed in the near future.

On the other hand, further studies focused on evaluating and overall fit of volumetric data are also needed in order to determine the pressure effects on the standard-state heat capacities and, consequently, to improve the accuracy of the heat capacity predictions as a function of temperature.

Additional thermodynamic calculations, based on the ionization of water, were performed to check the accuracy of the new C_p^0 values. The good agreement of $\Delta_f G^0_{OH^-}$ values with those calculated from the ionization constant of water reported by Bandura and Lvov⁴² corroborates the usefulness of equation 15 to estimate the temperature-dependent C_p^0 values for the $OH^-(aq)$ ion. Also, a comparison of estimated $\Delta_r C_p^0$ values, which represent two different equilibrium approaches (the ionization of water and the NaOH/HCl neutralization), shows satisfactory agreement. This last fact gives credence to the validity and reliability of the estimations of the standard-state molar heat capacities of NaOH(aq), HCl(aq), and NaCl(aq) made in this paper.

Acknowledgement

The authors acknowledge and thank the helpful comments and suggestions of Dr. Gerard Nihous (HNEI). We thank two anonymous reviewers for their helpful and constructive comments. JJM acknowledges the financial support provided by the European Social Found and the Spanish Ministry of Science and Innovation (“Juan de la Cierva” contract and “José Castillejo” mobility program). MJA acknowledges support from the Coral Industries Endowment of the University of Hawaii and thanks his colleagues in HNEI for their continued interest and encouragement.



European Social Found

Nomenclature

A_J = molality-based Debye-Hückel coefficient for heat capacities ($\text{J kg}^{1/2} \text{K}^{-1} \text{mol}^{-3/2}$)

B_c = adjustable parameter of the RRM model ($\text{J kg mol}^{-1} \text{K}^{-1}$)

b = arbitrary constant of the Pitzer equation ($1.2 \text{ kg}^{1/2} \text{mol}^{1/2}$)

Cp^0 = Standard-state partial molar heat capacity ($\text{J mol}^{-1} \text{K}^{-1}$)

Cp^θ = apparent molar heat capacity ($\text{J mol}^{-1} \text{K}^{-1}$)

Cp^{exp} = specific heat capacity of the solution ($\text{J g}^{-1} \text{K}^{-1}$)

Cp_w^0 = specific heat capacity of water ($\text{J g}^{-1} \text{K}^{-1}$)

C_{MX}^J = ion interaction parameter for Pitzer's equation

I = molality-based ionic strength (mol kg^{-1} for 1-1 electrolytes)

K_w = Ionization constant of water (–)

M = molecular weight of the solute (g mol^{-1})

m = molality of the solute (mol kg^{-1})

m^0 = molality value at the reference state (1.0 mol kg^{-1})

p = pressure (MPa)

R = universal gas constant ($8.3145 \text{ J mol}^{-1} \text{K}^{-1}$)

R^2 = coefficient of determination

R^2_{adj} = adjusted regression coefficient

T = temperature (K)

V^θ = apparent partial molar volume ($\text{cm}^3 \text{mol}^{-1}$)

w = valence factor used in the RRM equation

z = formal charge of an ion

Greek Symbols

α = arbitrary constant of the Pitzer equation ($2.0 \text{ kg}^{1/2} \text{mol}^{1/2}$)

$\beta_{MX}^{(0)J}$, $\beta_{MX}^{(1)J}$ = ion interaction parameters for Pitzer's equation

$\Delta_f G^0$ = standard-state Gibbs free energy of formation (J mol^{-1})

$\Delta_r G^0$ = standard-state Gibbs free energy of reaction (J mol^{-1})

$\Delta_r S^0$ = standard-state entropy change for the reaction ($\text{J mol}^{-1} \text{K}^{-1}$)

$\Delta_r C_p^0$ = standard-state heat capacity change for the reaction ($\text{J mol}^{-1} \text{K}^{-1}$)

ν_M = stoichiometric coefficients of a cation

ν_X = stoichiometric coefficients of an anion

Literature Cited

- (1) Puigdomenech, I.; Rard, J. A.; Plyasunov, A. V.; Grenthe, I. Temperature Corrections to Thermodynamic data and Enthalpy Calculations. In *Modelling in Aquatic Chemistry*; Grenthe, I., Puigdomenech, I., Eds.; OECD Nuclear Energy Agency: 1997; pp 427.
- (2) Archer, D. G. Thermodynamic Properties of the NaCl + H₂O System II. Thermodynamic Properties of NaCl(aq), NaCl·2H₂O, and Phase Equilibria. *J. Phys. Chem. Ref. Data* **1992**, *21*, 793-829.
- (3) Hovey, J. K.; Hepler, L. G.; Tremaine, P. R. Thermodynamics of aqueous aluminate ion: standard partial molar heat capacities and volumes of tetrahydroxyluminate(1-)(aq) from 10 to 55.degree.C. *J. Phys. Chem.* **1988**, *92*, 1323-1332.
- (4) Saluja, P. P. S.; Jobe, D. J.; Leblanc, J. C.; Lemire, R. J. Apparent Molar Heat-Capacities and Volumes of Mixed Electrolytes - [NaCl(aq)+CaCl₂(aq)], [NaCl(aq)+MgCl₂(aq)], and [CaCl₂(aq)+MgCl₂(aq)]. *J. Chem. Eng. Data* **1995**, *40*, 398-406.
- (5) Hovey, J. K. Thermodynamics of Hydration of a 4+ Aqueous Ion: Partial Molar Heat Capacities and Volumes of Aqueous Thorium(IV) from 10 to 55 °C. *J. Phys. Chem. B* **1997**, *101*, 4321-4334.
- (6) Ballerat-Busserolles, K.; Ford, T. D.; Call, T. G.; Woolley, E. M. Apparent molar volumes and heat capacities of aqueous acetic acid and sodium acetate at temperatures from T = 278.15 K to T = 393.15 K at the pressure 0.35MPa. *J. Chem. Thermodyn.* **1999**, *31*, 741-762.
- (7) Jones, J. S.; Ziemer, S. P.; Brown, B. R.; Woolley, E. M. Apparent molar volumes and apparent molar heat capacities of aqueous magnesium nitrate, strontium nitrate, and manganese nitrate at temperatures from 278.15 K to 393.15 K and at the pressure 0.35 MPa. *J. Chem. Thermodyn.* **2007**, *39*, 550-560.
- (8) Pitzer, K. S. Thermodynamics of electrolytes. I. Theoretical basis and general equations *J. Phys. Chem.* **1973**, *77*, 268-277.
- (9) Bradley, D. J.; Pitzer, K. S. Thermodynamics of electrolytes. 12. Dielectric properties of water and Debye-Huckel parameters to 350.degree.C and 1 kbar. *J. Phys. Chem.* **1979**, *83*, 1599-1603.

- (10) Fernandez, D. P.; Goodwin, A. R. H.; Lemmon, E. W.; Sengers, J. M. H. L.; Williams, R. C. A formulation for the static permittivity of water and steam at temperatures from 238 K to 873 K at pressures up to 1200 MPa, including derivatives and Debye-Huckel coefficients. *J. Phys. Chem. Ref. Data* **1997**, *26*, 1125-1166.
- (11) Magalhaes, M. C. F.; Konigsberger, E.; May, P. M.; Hefter, G. Heat Capacities of Concentrated Aqueous Solutions of Sodium Sulfate, Sodium Carbonate, and Sodium Hydroxide at 25 °C. *J. Chem. Eng. Data* **2002**, *47*, 590-598.
- (12) Perron, G.; Desnoyers, J. E.; Millero, F. J. Apparent Molal Volumes and Heat Capacities of Some Sulfates and Carbonates in Water at 25 °C. *Can. J. Chem.* **1975**, *53*, 1134-1138.
- (13) Weisberg, S. *Applied Linear Regression*; John Wiley & Sons: Hoboken, New Jersey, 2005.
- (14) Marquardt, D. An Algorithm for Least-Squares Estimation of Nonlinear Parameters. *SIAM Journal on Applied Mathematics* **1963**, *11*, 431-441.
- (15) Criss, C. M.; Millero, F. J. Modeling the Heat Capacities of Aqueous 1-1 Electrolyte Solutions with Pitzer's Equations. *J. Phys. Chem.* **1996**, *100*, 1288-1294.
- (16) Desnoyers, J. E.; de Visser, C.; Perron, G.; Picker, P. Reexamination of the Heat Capacities Obtained by Flow Microcalorimetry. Recommendation for the Use of a Chemical Standard. *J. Solution Chem.* **1976**, *5*, 605-616.
- (17) Smith-Magowan, D.; Wood, R. H. Heat capacity of aqueous sodium chloride from 320 to 600 K measured with a new flow calorimeter. *J. Chem. Thermodyn.* **1981**, *13*, 1047-1073.
- (18) Tanner, J. E.; Lamb, F. W. Specific heats of aqueous solutions of NaCl, NaBr, and KCl: Comparisons with related thermal properties. *J. Solution Chem.* **1978**, *7*, 303-316.
- (19) Perron, G.; Fortier, J.; Desnoyers, J. E. The apparent molar heat capacities and volumes of aqueous NaCl from 0.01 to 3 mol kg⁻¹ in the temperature range 274.65 to 318.15 K. *J. Chem. Thermodyn.* **1975**, *7*, 1177-1184.
- (20) Pabalan, R. T.; Pitzer, K. S. Apparent molar heat capacity and other thermodynamic properties of aqueous potassium chloride solutions to high temperatures and pressures. *J. Chem. Eng. Data* **1988**, *33*, 354-362.

- (21) Helgeson, H. C.; Kirkham, D. H.; Flowers, G. C. Theoretical Prediction of the Thermodynamic Behavior of Aqueous-Electrolytes at High-Pressures and Temperatures .4. Calculation of Activity-Coefficients, Osmotic Coefficients, and Apparent Molal and Standard and Relative Partial Molal Properties to 600-Degrees-C and 5 Kb. *Am. J. Sci.* **1981**, *281*, 1249-1516.
- (22) Shock, E. L.; Helgeson, H. C. Calculation of the Thermodynamic and Transport-Properties of Aqueous Species at High-Pressures and Temperatures - Correlation Algorithms for Ionic Species and Equation of State Predictions to 5-Kb and 1000-Degrees-C. *Geochim. Cosmochim. Acta* **1988**, *52*, 2009-2036.
- (23) Tanager, J. C.; Helgeson, H. C. Calculation of the Thermodynamic and Transport-Properties of Aqueous Species at High-Pressures and Temperatures - Revised Equations of State for the Standard Partial Molal Properties of Ions and Electrolytes. *Am. J. Sci.* **1988**, *288*, 19-98.
- (24) Dick, J. Calculation of the relative metastabilities of proteins using the CHNOSZ software package. *Geochem. Trans.* **2008**, *9*, 10.
- (25) Wagner, W.; Pruss, A. The IAPWS formulation 1995 for the thermodynamic properties of ordinary water substance for general and scientific use. *J. Phys. Chem. Ref. Data* **2002**, *31*, 387-535.
- (26) Archer, D. G.; Wang, P. M. The dielectric constant of water and Debye-Huckel limiting law slopes. *J. Phys. Chem. Ref. Data* **1990**, *19*, 371-411.
- (27) Patterson, B. A.; Call, T. G.; Jardine, J. J.; Origlia-Luster, M. L.; Woolley, E. M. Thermodynamics for ionization of water at temperatures from 278.15 K to 393.15 K and at the pressure 0.35 MPa: apparent molar volumes of aqueous KCl, KOH, and NaOH and apparent molar heat capacities of aqueous HCl, KCl, KOH, and NaOH. *J. Chem. Thermodyn.* **2001**, *33*, 1237-1262.
- (28) Tremaine, P. R.; Sway, K.; Barbero, J. A. The apparent molar heat capacity of aqueous hydrochloric acid from 10 to 140°C. *J. Solution Chem.* **1986**, *15*, 1-22.
- (29) Allred, G. C.; Woolley, E. M. Heat capacities of aqueous acetic acid, sodium acetate, ammonia, and ammonium chloride at 283.15, 298.15, and 313.15 K: ΔC_p° for ionization of acetic acid and for dissociation of ammonium ion. *J. Chem. Thermodyn.* **1981**, *13*, 155-164.

- (30) Sharygin, A. V.; Wood, R. H. Volumes and heat capacities of aqueous solutions of hydrochloric acid at temperatures from 298.15 K to 623 K and pressures to 28 MPa. *J. Chem. Thermodyn.* **1997**, *29*, 125-148.
- (31) Barner, H. E.; Scheuerman, R. V. *Handbook of Thermochemical Data for Compounds and Aqueous Species*; Wiley-Interscience: New York, 1973.
- (32) Criss, C. M.; Cobble, J. W. The Thermodynamic Properties of High Temperature Aqueous Solutions. IV. Entropies of the Ions up to 200° and the Correspondence Principle. *J. Am. Chem. Soc.* **1964**, *86*, 5385-5390.
- (33) Simonson, J. M.; Mesmer, R. E.; Rogers, P. S. Z. The enthalpy of dilution and apparent molar heat capacity of NaOH(aq) to 523 K and 40 MPa. *J. Chem. Thermodyn.* **1989**, *21*, 561-584.
- (34) Conti, G.; Gianni, P.; Papini, A.; Matteoli, E. Apparent Molar Heat-Capacity and Relative Enthalpy of Aqueous NaOH between 323 K and 523 K. *J. Solution Chem.* **1988**, *17*, 481-497.
- (35) Ballerat-Busserolles, K.; Origlia, M. L.; Woolley, E. M. Calibration of a fixed-cell temperature-scanning calorimeter to measure precise solution heat capacities from 275 to 398 K at 0.35 MPa. *Thermochim. Acta* **2000**, *347*, 3-7.
- (36) Cerdeiriña, C. A.; Míguez, J. A.; Carballo, E.; Tovar, C. A.; de la Puente, E.; Romani, L. Highly precise determination of the heat capacity of liquids by DSC: calibration and measurement. *Thermochim. Acta* **2000**, *347*, 37-44.
- (37) Marshall, W. L.; Franck, E. U. Ion product of water substance, 0 – 1000 °C, 1–10,000 bars. New international formulation and its background. *J. Phys. Chem. Ref. Data* **1981**, *10*, 295.
- (38) Pitzer, K. S. Self-ionization of water at high temperature and thermodynamic properties of ions. *J. Phys. Chem.* **1982**, *86*, 4704-4708.
- (39) Tanger, J. C.; Pitzer, K. S. Calculation of the ionization constant of H₂O to 2273 K and 500 MPa. *AIChE J.* **1989**, *35*, 1631-1638.
- (40) Chen, X.; Izatt, R. M.; Oscarson, J. L. Thermodynamic data for ligand interaction with protons and metal ions in aqueous solutions at high temperatures. *Chem. Rev.* **1994**, *94*, 467-517.
- (41) Tawa, G. J.; Pratt, L. R. Theoretical calculation of the water ion product K_w . *J. Am. Chem. Soc.* **1995**, *117*, 1625-1628.
- (42) Bandura, A. V.; Lvov, S. N. The ionization constant of water over wide ranges of temperature and density *J. Phys. Chem. Ref. Data* **2006**, *35*, 16-30.

- (43) Schrödle, S.; Königsberger, E.; May, P. M.; Hefter, G. Heat capacities of aqueous solutions of sodium hydroxide and water ionization up to 300 °C at 10 MPa. *Geochim. Cosmochim. Acta* **2008**, 72, 3124-3138.

Table 1. Parameters of the exponential function given by equation 6

	Temperature (K)				
	300	350	400	450	500
a_0	1.14073	1.03078	2.13600	2.61389	2.76623
a_1	2.75999	5.02168	7.14257	12.91512	29.51689
a_2	-0.00316	-0.00291	-0.00465	-0.00668	-0.00999
R^2adj	1.0000	0.9979	0.9999	0.9968	0.9985

Table 2. Sources of experimental data of C_p^θ of NaCl(aq)

Reference	Instrument	Molality range	Operating conditions	
			Temperature range (K)	Pressure (MPa)
Saluja et al. (1995) ⁴	Flow heat capacity microcalorimeter	0.1-6.0	298.15-371.82	0.60
Smith-Magowan and Wood (1981) ¹⁷	Heat flow calorimeter	0.1-3.0	320.55-497.73	10.00-17.70
Tanner and Lamb (1978) ¹⁸	Twin heat flow calorimeter	0.04-1.25	298.15-358.15	0.10
Perron et al. (1975) ¹⁹	Flow microcalorimeter	0.02-0.95	288.15-318.15	0.10

Table 3. Summarized results for the NaCl(aq) using the ion-interaction (Pitzer) approach

Source	T (K)	p (MPa)	Cp^0 (J mol ⁻¹ K ⁻¹)	$\beta_{MX}^{(0)J}$ (kg mol ⁻¹ K ⁻²)	$\beta_{MX}^{(1)J}$ (kg ² mol ⁻² K ⁻²)	C_{MX}^J (kg ² mol ⁻² K ⁻²)	$Prob.>F$	R^2_{adj}
Saluja et al. (1995) ⁴	298.15	0.6	-86.22 ± 1.20	-3.02 10 ⁻⁵ ± 4.07 10 ⁻⁶	-4.51 10 ⁻⁵ ± 1.35 10 ⁻⁵	2.49 10 ⁻⁵ ± 7.11 10 ⁻⁶	1.79 10 ⁻⁹	0.9995
Saluja et al. (1995) ⁴	323.15	0.6	-56.87 ± 0.59	-5.09 10 ⁻⁶ ± 1.70 10 ⁻⁶	6.69 10 ⁻⁶ ± 5.66 10 ⁻⁶	-3.25 10 ⁻⁶ ± 2.97 10 ⁻⁶	4.24 10 ⁻¹⁰	0.9998
Saluja et al. (1995) ⁴	348.15	0.6	-59.25 ± 0.56	-2.65 10 ⁻⁶ ± 1.39 10 ⁻⁶	Not significant	Not significant	3.95 10 ⁻¹⁰	0.9998
Saluja et al. (1995) ⁴	373.15	0.6	-76.13 ± 0.96	-1.12 10 ⁻⁵ ± 2.41 10 ⁻⁶	-1.98 10 ⁻⁵ ± 8.00 10 ⁻⁶	1.06 10 ⁻⁵ ± 4.21 10 ⁻⁶	1.73 10 ⁻⁹	0.9994
Tanner and Lamb (1978) ¹⁸	298.15	0.1	-87.64 ± 1.38	-2.58 10 ⁻⁵ ± 1.15 10 ⁻⁵	Not significant	Not significant	5.00 10 ⁻¹⁵	0.9904
Tanner and Lamb (1978) ¹⁸	318.15	0.1	-70.74 ± 1.33	-4.27 10 ⁻⁵ ± 7.64 10 ⁻⁶	-1.79 10 ⁻⁴ ± 5.24 10 ⁻⁵	8.20 10 ⁻⁵ ± 2.26 10 ⁻⁵	3.60 10 ⁻⁹	0.9955
Tanner and Lamb (1978) ¹⁸	338.15	0.1	-59.08 ± 0.41	-4.29 10 ⁻⁶ ± 2.00 10 ⁻⁶	-2.19 10 ⁻⁵ ± 1.32 10 ⁻⁵	7.87 10 ⁻⁶ ± 5.74 10 ⁻⁶	2.14 10 ⁻⁷	0.9995
Tanner and Lamb (1978) ¹⁸	358.15	0.1	-64.32 ± 6.42	Not significant	Not significant	Not significant	1.23 10 ⁻³	0.9762
Smith-Magowan and Wood (1981) ¹⁷	320.60	17.7	-43.84 ± 1.75	Not significant	Not significant	Not significant	2.58 10 ⁻²	0.9970
Smith-Magowan and Wood (1981) ¹⁷	358.40	17.7	-53.52 ± 3.39	Not significant	Not significant	Not significant	4.49 10 ⁻²	0.9885
Smith-Magowan and Wood (1981) ¹⁷	397.97	17.7	-83.12 ± 6.53	Not significant	Not significant	Not significant	5.04 10 ⁻²	0.9750
Smith-Magowan and Wood (1981) ¹⁷	452.56	17.7	-182.96 ± 14.24	-4.17 10 ⁻⁵ ± 2.98 10 ⁻⁵	-1.53 10 ⁻⁴ ± 1.29 10 ⁻⁴	7.83 10 ⁻⁵ ± 6.52 10 ⁻⁵	5.16 10 ⁻²	0.9645
Smith-Magowan and Wood (1981) ¹⁷	491.05	17.7	-287.11 ± 3.15	-5.08 10 ⁻⁵ ± 5.60 10 ⁻⁶	-1.98 10 ⁻⁴ ± 2.44 10 ⁻⁵	1.01 10 ⁻⁴ ± 1.22 10 ⁻⁵	7.22 10 ⁻³	0.9991
Smith-Magowan and Wood (1981) ¹⁷	497.73	17.7	-336.96 ± 21.57	-6.62 10 ⁻⁵ ± 3.73 10 ⁻⁵	-2.41 10 ⁻⁴ ± 1.62 10 ⁻⁴	1.24 10 ⁻⁴ ± 8.17 10 ⁻⁵	4.27 10 ⁻²	0.9731

Table 4. C_p^0 values of NaCl(aq) estimated from the low-molality experimental data of Saluja et al.⁴ and their comparison with values deduced from the full data set corresponding to the same work

$T(K)$	C_p^0 values correlated from experimental data ($\text{J mol}^{-1} \text{K}^{-1}$)	Difference respect to the values reported in Table 3 (%)
298.15	-87.63 ± 0.54	-1.64
323.15	-56.00 ± 1.63	1.53
348.15	-60.26 ± 0.89	-1.70
373.15	-79.90 ± 2.23	-4.95

Table 5. Summarized results for the NaCl(aq) data reported by Saluja and coworkers⁴ ($m \leq 1 \text{ mol kg}^{-1}$) using the RRM type equation

T (K)	Cp^0 ($\text{J mol}^{-1} \text{K}^{-1}$)	Difference respect to the values reported in Table 3 (%)	B_c ($\text{J kg mol}^{-2} \text{K}^{-1}$)	Prob.>F	R^2_{adj}
298.15	-80.16 ± 1.67	-8.52	8.70 ± 0.66	$1.90 \cdot 10^{-8}$	0.9910
323.15	-59.19 ± 0.37	5.70	1.80 ± 0.15	$1.02 \cdot 10^{-11}$	0.9992
348.15	-63.46 ± 1.00	5.31	-2.97 ± 0.40	$1.36 \cdot 10^{-8}$	0.9932
373.15	-82.66 ± 1.23	3.45	-6.09 ± 0.49	$9.96 \cdot 10^{-9}$	0.9912

Table 6. Sources of experimental data of Cp^θ of HCl(aq)

Reference	Instrument	Operating conditions		
		Molality range	Temperature range (K)	p (MPa)
Patterson et al. (2001) ²⁷	Fixed-cell scanning calorimeter (NanoDSC model 6100, Calorimetry Sciences Corporation)	0.015-0.500	298.15-393.15	0.35
Sharigyn and Wood (1997) ³⁰	Flow Picker-type calorimeter	0.2-6.0	302.14-523.92	27.97
Tremaine et al. (1986) ²⁸	Flow Picker-type calorimeter	0.09-1.00	298.15-412.61	0.10-0.50
Allred and Wooley (1981) ²⁹	Picker flow microcalorimeter	0.05-0.20	298.15-313.15	0.10

Table 7. Summarized results for the HCl(aq) using the ion-interaction (Pitzer) approach

Source	T (K)	p (MPa)	Cp^0 (J mol ⁻¹ K ⁻¹)	$\beta_{MX}^{(0)J}$ (kg mol ⁻¹ K ⁻²)	$\beta_{MX}^{(1)J}$ (kg ² mol ⁻² K ⁻²)	C_{MX}^J (kg ² mol ⁻² K ⁻²)	$Prob.>F$	R^2_{adj}
Patterson et al. (2001) ²⁷	298.15	0.35	-124.84 ± 1.58	-4.55 10 ⁻⁵ ± 3.17 10 ⁻⁵	-7.72 10 ⁻⁴ ± 6.98 10 ⁻⁴	Not significant	8.94 10 ⁻⁵	0.9470
Patterson et al. (2001) ²⁷	303.15	0.35	-120.60 ± 1.70	-4.77 10 ⁻⁵ ± 3.31 10 ⁻⁵	Not significant	Not significant	1.13 10 ⁻⁴	0.9411
Patterson et al. (2001) ²⁷	308.15	0.35	-117.36 ± 1.57	-4.40 10 ⁻⁵ ± 2.96 10 ⁻⁵	Not significant	2.31 10 ⁻⁴ ± 2.07 10 ⁻⁴	1.02 10 ⁻⁴	0.9510
Patterson et al. (2001) ²⁷	313.15	0.35	-114.79 ± 1.52	-4.09 10 ⁻⁵ ± 2.78 10 ⁻⁵	Not significant	2.16 10 ⁻⁴ ± 1.95 10 ⁻⁵	1.02 10 ⁻⁴	0.9555
Patterson et al. (2001) ²⁷	318.15	0.35	-114.23 ± 1.61	-4.53 10 ⁻⁵ ± 2.84 10 ⁻⁵	-7.89 10 ⁻⁴ ± 7.14 10 ⁻⁴	2.43 10 ⁻⁴ ± 1.99 10 ⁻⁴	1.16 10 ⁻⁴	0.9558
Patterson et al. (2001) ²⁷	323.15	0.35	-113.88 ± 1.50	-4.35 10 ⁻⁵ ± 2.58 10 ⁻⁵	-7.32 10 ⁻⁴ ± 6.48 10 ⁻⁴	2.27 10 ⁻⁴ ± 1.81 10 ⁻⁴	1.04 10 ⁻⁴	0.9650
Patterson et al. (2001) ²⁷	328.15	0.35	-114.33 ± 1.60	-4.64 10 ⁻⁵ ± 2.65 10 ⁻⁵	-7.83 10 ⁻⁴ ± 6.68 10 ⁻⁴	2.43 10 ⁻⁴ ± 1.86 10 ⁻⁴	1.18 10 ⁻⁴	0.9646
Patterson et al. (2001) ²⁷	333.15	0.35	-114.26 ± 1.35	-3.87 10 ⁻⁵ ± 2.18 10 ⁻⁵	-6.31 10 ⁻⁴ ± 5.48 10 ⁻⁴	1.98 10 ⁻⁴ ± 1.53 10 ⁻⁴	8.45 10 ⁻⁵	0.9757
Patterson et al. (2001) ²⁷	338.15	0.35	-114.63 ± 1.09	-3.20 10 ⁻⁵ ± 1.71 10 ⁻⁵	-4.94 10 ⁻⁴ ± 4.29 10 ⁻⁴	Not significant	5.48 10 ⁻⁵	0.9851
Patterson et al. (2001) ²⁷	343.15	0.35	-116.10 ± 1.15	-3.18 10 ⁻⁵ ± 1.74 10 ⁻⁵	-4.71 10 ⁻⁴ ± 4.38 10 ⁻⁴	1.52 10 ⁻⁴ ± 1.22 10 ⁻⁴	5.96 10 ⁻⁵	0.9852
Patterson et al. (2001) ²⁷	348.15	0.35	-117.26 ± 0.80	-2.42 10 ⁻⁵ ± 1.18 10 ⁻⁵	-3.40 10 ⁻⁴ ± 2.96 10 ⁻⁴	1.11 10 ⁻⁴ ± 8.24 10 ⁻⁵	2.82 10 ⁻⁵	0.9932
Patterson et al. (2001) ²⁷	353.15	0.35	-119.50 ± 0.94	-2.68 10 ⁻⁵ ± 1.35 10 ⁻⁵	-3.75 10 ⁻⁴ ± 3.40 10 ⁻⁴	1.22 10 ⁻⁴ ± 9.48 10 ⁻⁵	3.85 10 ⁻⁵	0.9915
Patterson et al. (2001) ²⁷	358.15	0.35	-121.93 ± 0.92	-2.76 10 ⁻⁵ ± 1.28 10 ⁻⁵	-3.83 10 ⁻⁴ ± 3.21 10 ⁻⁴	1.25 10 ⁻⁴ ± 8.96 10 ⁻⁵	3.53 10 ⁻⁵	0.9929
Patterson et al. (2001) ²⁷	363.15	0.35	-124.60 ± 0.95	-2.80 10 ⁻⁵ ± 1.30 10 ⁻⁵	-3.86 10 ⁻⁴ ± 3.25 10 ⁻⁴	1.26 10 ⁻⁴ ± 9.08 10 ⁻⁵	3.70 10 ⁻⁵	0.9930
Patterson et al. (2001) ²⁷	368.15	0.35	-127.86 ± 1.01	-3.16 10 ⁻⁵ ± 1.34 10 ⁻⁵	-4.50 10 ⁻⁴ ± 3.36 10 ⁻⁴	1.45 10 ⁻⁴ ± 9.36 10 ⁻⁵	3.99 10 ⁻⁵	0.9930
Patterson et al. (2001) ²⁷	373.15	0.35	-130.90 ± 0.89	-3.04 10 ⁻⁵ ± 1.15 10 ⁻⁵	-4.20 10 ⁻⁴ ± 2.88 10 ⁻⁴	1.36 10 ⁻⁴ ± 8.04 10 ⁻⁵	2.98 10 ⁻⁵	0.9951
Patterson et al. (2001) ²⁷	378.15	0.35	-135.14 ± 1.06	-3.49 10 ⁻⁵ ± 1.33 10 ⁻⁵	-4.49 10 ⁻⁴ ± 3.33 10 ⁻⁴	1.60 10 ⁻⁴ ± 9.28 10 ⁻⁵	3.98 10 ⁻⁵	0.9939
Patterson et al. (2001) ²⁷	383.15	0.35	-139.01 ± 1.07	-3.66 10 ⁻⁵ ± 1.31 10 ⁻⁵	-5.23 10 ⁻⁴ ± 3.39 10 ⁻⁴	1.67 10 ⁻⁴ ± 9.18 10 ⁻⁵	3.90 10 ⁻⁵	0.9944

Table 7 (continued)

Source	T (K)	p (MPa)	Cp^0 (J mol ⁻¹ K ⁻¹)	$\beta_{MX}^{(0)J}$ (kg mol ⁻¹ K ⁻²)	$\beta_{MX}^{(1)J}$ (kg ² mol ⁻² K ⁻²)	C_{MX}^J (kg ² mol ⁻² K ⁻²)	$Prob.>F$	R^2_{adj}
Patterson et al. (2001) ²⁷	388.15	0.35	-143.55 ± 1.10	-4.01 10 ⁻⁵ ± 1.31 10 ⁻⁵	-5.76 10 ⁻⁴ ± 3.28 10 ⁻⁴	1.84 10 ⁻⁴ ± 9.16 10 ⁻⁵	3.87 10 ⁻⁵	0.9948
Patterson et al. (2001) ²⁷	393.15	0.35	-148.76 ± 1.29	-4.59 10 ⁻⁵ ± 1.50 10 ⁻⁵	-6.98 10 ⁻⁴ ± 3.77 10 ⁻⁴	2.18 10 ⁻⁴ ± 1.05 10 ⁻⁴	5.05 10 ⁻⁵	0.9936
Sharigyn and Wood (1997) ³⁰	302.14	27.97	-113.13 ± 2.39	-9.10 10 ⁻⁶ ± 7.10 10 ⁻⁶	-2.51 10 ⁻⁵ ± 2.26 10 ⁻⁵	Not significant	7.79 10 ⁻³	0.9967
Sharigyn and Wood (1997) ³⁰	373.71	27.97	-118.02 ± 0.69	Not significant	6.03 10 ⁻⁶ ± 4.26 10 ⁻⁶	-3.24 10 ⁻⁶ ± 2.25 10 ⁻⁶	2.52 10 ⁻³	0.9999
Sharigyn and Wood (1997) ³⁰	448.70	27.97	-206.34 ± 1.28	-2.66 10 ⁻⁵ ± 2.05 10 ⁻⁶	-6.91 10 ⁻⁵ ± 6.77 10 ⁻⁶	3.68 10 ⁻⁵ ± 3.56 10 ⁻⁶	4.45 10 ⁻⁵	0.9998
Sharigyn and Wood (1997) ³⁰	523.92	27.97	-377.11 ± 5.80	-4.85 10 ⁻⁵ ± 6.79 10 ⁻⁶	-1.23 10 ⁻⁴ ± 2.24 10 ⁻⁵	6.59 10 ⁻⁵ ± 1.18 10 ⁻⁵	4.02 10 ⁻⁴	0.9992
Tremaine et al. (1986) ²⁸	298.15	0.1	-127.03 ± 1.22	-1.40 10 ⁻⁵ ± 8.09 10 ⁻⁶	-8.40 10 ⁻⁵ ± 6.44 10 ⁻⁵	3.57 10 ⁻⁵ ± 2.63 10 ⁻⁵	4.80 10 ⁻¹⁰	0.9875
Tremaine et al. (1986) ²⁸	398.25	0.3	-130.02 ± 8.46	3.18 10 ⁻⁵ ± 2.64 10 ⁻⁶	2.46 10 ⁻⁴ ± 1.85 10 ⁻⁴	-1.04 10 ⁻⁴ ± 7.80 10 ⁻⁵	4.07 10 ⁻⁶	0.9733
Tremaine et al. (1986) ²⁸	412.61	0.5	-153.42 ± 2.32	Not significant	4.26 10 ⁻⁵ ± 6.36 10 ⁻⁶	Not significant	6.14 10 ⁻⁹	0.9951
Allred and Woolley (1981) ²⁹	298.15	0.1	-125.43 ± 1.22	-4.19 10 ⁻⁵ ± 2.48 10 ⁻⁵	-9.94 10 ⁻³ ± 3.05 10 ⁻³	1.04 10 ⁻³ ± 4.14 10 ⁻⁴	9.54 10 ⁻⁴	0.9970

Table 8. Sources of experimental data of Cp^θ of NaOH(aq)

Reference	Instrument	Molality range	Operating conditions	
			Temperature range (K)	Pressure (MPa)
Magalhaes et al. (2002) ¹¹	Picker flow calorimeter (Sodev, Canada)	0.02-1.00	298.15	0.10
Patterson et al. (2001) ²⁷	Fixed-cell scanning calorimeter (NanoDSC model6100, Calorimetry Sciences Corporation)	0.015-0.500	298.15-393.15	0.35
Simonson et al. (1989) ³³	Picker flow calorimeter	0.10-4.08	323.04-522.79	7.00
Conti et al. (1988) ³⁴	Picker flow calorimeter	0.05-1.50	325.25-521.55	0.10-4.00
Allred and Wooley (1981) ²⁹	Picker flow microcalorimeter	0.05-0.20	298.15-313.15	0.10

Table 9. Summarized results for the aqueous electrolyte NaOH using the ion-interaction (Pitzer) approach

Source	T (K)	p (MPa)	Cp^0 (J mol ⁻¹ K ⁻¹)	$\beta_{MX}^{(0)J}$ (kg mol ⁻¹ K ⁻²)	$\beta_{MX}^{(1)J}$ (kg ² mol ⁻² K ⁻²)	C_{MX}^J (kg ² mol ⁻² K ⁻²)	Prob.>F	R^2_{adj}
Magalhaes et al. (2002) ¹¹	298.15	0.1	-99.12 ± 0.44	-3.68 10 ⁻⁵ ± 4.84 10 ⁻⁶	Not significant	Not significant	7.08 10 ⁻¹⁴	0.9995
Patterson et al. (2001) ²⁷	298.15	0.35	-102.42 ± 1.41	-6.33 10 ⁻⁵ ± 2.73 10 ⁻⁵	-6.55 10 ⁻⁴ ± 6.33 10 ⁻⁴	1.96 10 ⁻⁴ ± 1.80 10 ⁻⁵	1.42 10 ⁻⁴	0.9912
Patterson et al. (2001) ²⁷	303.15	0.35	-88.86 ± 1.35	-5.20 10 ⁻⁵ ± 2.53 10 ⁻⁵	Not significant	Not significant	1.78 10 ⁻⁴	0.9904
Patterson et al. (2001) ²⁷	308.15	0.35	-78.49 ± 1.29	-4.44 10 ⁻⁵ ± 2.35 10 ⁻⁵	Not significant	Not significant	2.18 10 ⁻⁴	0.9902
Patterson et al. (2001) ²⁷	313.15	0.35	-69.80 ± 1.25	-3.67 10 ⁻⁵ ± 2.20 10 ⁻⁵	Not significant	Not significant	2.69 10 ⁻⁴	0.9897
Patterson et al. (2001) ²⁷	318.15	0.35	-63.30 ± 1.18	-3.08 10 ⁻⁵ ± 2.01 10 ⁻⁵	Not significant	Not significant	3.04 10 ⁻⁴	0.9904
Patterson et al. (2001) ²⁷	323.15	0.35	-58.13 ± 1.16	-2.60 10 ⁻⁵ ± 1.91 10 ⁻⁵	Not significant	Not significant	3.62 10 ⁻⁴	0.9904
Patterson et al. (2001) ²⁷	328.15	0.35	-54.19 ± 1.06	-2.10 10 ⁻⁵ ± 1.69 10 ⁻⁵	Not significant	Not significant	3.64 10 ⁻⁴	0.9919
Patterson et al. (2001) ²⁷	333.15	0.35	-51.13 ± 0.98	-1.62 10 ⁻⁵ ± 1.51 10 ⁻⁵	Not significant	Not significant	3.62 10 ⁻⁴	0.9931
Patterson et al. (2001) ²⁷	338.15	0.35	-49.22 ± 1.03	-1.57 10 ⁻⁵ ± 1.55 10 ⁻⁵	Not significant	Not significant	4.53 10 ⁻⁴	0.9924
Patterson et al. (2001) ²⁷	343.15	0.35	-48.22 ± 1.01	-1.48 10 ⁻⁵ ± 1.47 10 ⁻⁵	-1.05 10 ⁻⁴ ± 3.42 10 ⁻⁵	Not significant	4.73 10 ⁻⁴	0.9930
Patterson et al. (2001) ²⁷	348.15	0.35	-47.83 ± 1.08	-1.30 10 ⁻⁵ ± 1.18 10 ⁻⁵	-8.27 10 ⁻⁵ ± 3.57 10 ⁻⁵	2.24 10 ⁻⁵ ± 1.16 10 ⁻⁵	5.72 10 ⁻⁴	0.9921
Patterson et al. (2001) ²⁷	353.15	0.35	-47.93 ± 0.96	Not significant	-3.56 10 ⁻⁵ ± 3.09 10 ⁻⁵	Not significant	4.58 10 ⁻⁴	0.9940
Patterson et al. (2001) ²⁷	358.15	0.35	-48.51 ± 0.95	Not significant	4.68 10 ⁻⁵ ± 2.96 10 ⁻⁵	Not significant	4.35 10 ⁻⁴	0.9944
Patterson et al. (2001) ²⁷	363.15	0.35	-49.93 ± 0.91	Not significant	5.95 10 ⁻⁵ ± 5.74 10 ⁻⁵	-2.14 10 ⁻⁵ ± 7.79 10 ⁻⁶	3.74 10 ⁻⁴	0.9952
Patterson et al. (2001) ²⁷	368.15	0.35	-52.08 ± 0.92	Not significant	1.11 10 ⁻⁴ ± 2.71 10 ⁻⁵	-3.50 10 ⁻⁵ ± 7.70 10 ⁻⁶	3.53 10 ⁻⁴	0.9954
Patterson et al. (2001) ²⁷	363.15	0.35	-49.93 ± 0.91	Not significant	5.95 10 ⁻⁵ ± 5.74 10 ⁻⁵	-2.14 10 ⁻⁵ ± 7.80 10 ⁻⁶	3.74 10 ⁻⁴	0.9952
Patterson et al. (2001) ²⁷	368.15	0.35	-52.08 ± 0.92	Not significant	1.11 10 ⁻⁴ ± 2.71 10 ⁻⁵	-3.50 10 ⁻⁵ ± 7.70 10 ⁻⁶	3.53 10 ⁻⁴	0.9954

Table 9 (*continued*)

Source	T (K)	p (MPa)	Cp^0 (J mol ⁻¹ K ⁻¹)	$\beta_{MX}^{(0)J}$ (kg mol ⁻¹ K ⁻²)	$\beta_{MX}^{(1)J}$ (kg ² mol ⁻² K ⁻²)	C_{MX}^J (kg ² mol ⁻² K ⁻²)	Prob.>F	R^2_{adj}
Patterson et al. (2001) ²⁷	373.15	0.35	-54.70 ± 0.91	Not significant	1.01 10 ⁻⁴ ± 2.62 10 ⁻⁵	-3.28 10 ⁻⁵ ± 7.44 10 ⁻⁶	3.12 10 ⁻⁴	0.9958
Patterson et al. (2001) ²⁷	378.15	0.35	-57.70 ± 0.81	Not significant	1.61 10 ⁻⁴ ± 2.26 10 ⁻⁵	-5.02 10 ⁻⁵ ± 4.45 10 ⁻⁵	2.17 10 ⁻⁴	0.9969
Patterson et al. (2001) ²⁷	383.15	0.35	-61.17 ± 0.72	Not significant	1.79 10 ⁻⁴ ± 1.75 10 ⁻⁴	Not significant	1.49 10 ⁻⁴	0.9978
Patterson et al. (2001) ²⁷	388.15	0.35	-65.17 ± 0.62	Not significant	1.81 10 ⁻⁴ ± 1.65 10 ⁻⁴	-5.66 10 ⁻⁵ ± 4.71 10 ⁻⁵	9.76 10 ⁻⁵	0.9984
Patterson et al. (2001) ²⁷	393.15	0.35	-68.41 ± 0.42	1.13 10 ⁻⁵ ± 4.65 10 ⁻⁶	3.71 10 ⁻⁴ ± 1.08 10 ⁻⁴	-1.14 10 ⁻⁴ ± 3.06 10 ⁻⁵	3.81 10 ⁻⁵	0.9993
Simonson et al. (1989) ³³	323.04	7.00	-59.91 ± 1.20	-3.32 10 ⁻⁵ ± 3.80 10 ⁻⁶	-5.99 10 ⁻⁵ ± 1.32 10 ⁻⁵	3.30 10 ⁻⁵ ± 6.88 10 ⁻⁶	4.49 10 ⁻⁴	0.9994
Simonson et al. (1989) ³³	373.07	7.00	-46.92 ± 1.66	-8.40 10 ⁻⁶ ± 3.92 10 ⁻⁶	Not significant	Not significant	1.44 10 ⁻³	0.9982
Simonson et al. (1989) ³³	423.09	7.00	-96.32 ± 6.89	Not significant	3.77 10 ⁻⁵ ± 3.44 10 ⁻⁵	Not significant	7.13 10 ⁻³	0.9892
Simonson et al. (1989) ³³	522.79	7.00	-449.63 ± 23.87	-1.75 10 ⁻⁵ ± 1.71 10 ⁻⁵	-9.23 10 ⁻⁵ ± 9.07 10 ⁻⁵	Not significant	2.95 10 ⁻³	0.9719
Conti et al. (1988) ³⁴	325.25	0.10	-64.43 ± 10.69	-6.12 10 ⁻⁵ ± 5.88 10 ⁻⁵	-2.47 10 ⁻⁴ ± 3.66 10 ⁻⁵	Not significant	1.55 10 ⁻²	0.8124
Conti et al. (1988) ³⁴	373.97	0.12	-46.10 ± 5.12	Not significant	-1.35 10 ⁻⁵ ± 1.33 10 ⁻⁵	Not significant	6.30 10 ⁻³	0.9371
Conti et al. (1988) ³⁴	422.45	0.50	-85.78 ± 3.20	2.42 10 ⁻⁵ ± 1.04 10 ⁻⁵	1.75 10 ⁻⁴ ± 6.49 10 ⁻⁵	-7.92 10 ⁻⁵ ± 2.92 10 ⁻⁵	1.22 10 ⁻⁴	0.9882
Conti et al. (1988) ³⁴	470.55	1.50	-182.85 ± 4.72	Not significant	3.52 10 ⁻⁵ ± 7.73 10 ⁻⁶	Not significant	3.51 10 ⁻⁵	0.9893
Conti et al. (1988) ³⁴	521.55	4.00	-436.03 ± 11.67	Not significant	4.50 10 ⁻⁵ ± 1.55 10 ⁻⁶	Not significant	3.65 10 ⁻⁵	0.9865
Allred and Woolley (1981) ²⁹	298.15	0.1	-95.16± 4.56	-3.12 10 ⁻⁴ ± 1.42 10 ⁻⁴	-2.35 10 ⁻² ± 1.55 10 ⁻²	3.65 10 ⁻³ ± 2.15 10 ⁻³	1.54 10 ⁻⁷	0.9850
Allred and Woolley (1981) ²⁹	313.15	0.1	-57.86± 5.04	Not significant	7.13 10 ⁻³ ± 1.20 10 ⁻³	Not significant	1.54 10 ⁻⁷	0.9850

Table 10. Values of $\Delta_f G^0_{OH^-}$ at various temperatures at 0.1 MPa ($T < 273.15$ K) or saturation pressure ($T > 273.15$ K)

Temperature (K)	$\Delta_f G^0_{OH^-}$ (kJ mol ⁻¹)			
	Using pK_w values reported in ref. 42 and eq. 15	Using heat capacity data from the present work and eq. 18 and 17	Using heat capacity data from the present work and eq. 16 and 17	Using heat capacity data from ref. 31 and eq. 18 and 17
323.15	-156.20	-157.13	-157.30	-157.15
348.15	-149.92	-150.87	-151.53	-150.93
373.15	-143.42	-144.26	-145.72	-144.36
423.15	-129.44	-129.84	-134.86	-130.29
473.15	-114.44	-112.88	-122.86	-114.94

Table 11. Standard molar heat capacity change (in $\text{J mol}^{-1} \text{K}^{-1}$) for the ionization of water at 10 MPa

Temperature (K)	323.15	348.15	373.15	423.15	473.15
This study from equilibrium: $\text{H}_2\text{O} = \text{H}^+ + \text{OH}^-$	-185.90	-178.96	-183.48	-213.09	-293.27
This study from equilibrium: $\text{NaOH} + \text{HCl} = \text{NaCl} + \text{H}_2\text{O}$	-187.02	-180.63	-180.17	-203.87	-298.27
Difference (%)	0.60	0.93	-1.80	-4.33	1.71
Ref. 43	-186 ± 5	n. a.	-182 ± 5	-224 ± 10	-314 ± 26
Revised HKF model	-175.25	-170.47	-179.21	-229.63	-355.20

Caption for Figures

Figure 1. A/R values as a function of temperature at a given pressure of 4.0 MPa.

Figure 2. Apparent molar heat capacity of NaCl(aq) versus molality at 348.15 K (squares: experimental data from Saluja et al.⁴; line: Pitzer approach performed in this study).

Figure 3. Apparent molar heat capacity of NaCl(aq) versus molality at (a) 298.15 K (\square , experimental data from Saluja et al.⁴; \bullet , data from Tanner and Lamb¹⁸) and (b) 318.15 K (\triangle , data from Perron et al.¹⁹; \bullet , data from Tanner and Lamb¹⁸).

Figure 4. Estimated values of Cp^0 for NaCl(aq) using the Pitzer approach to correlate experimental Cp^0 values obtained by Saluja and coworkers⁴ (\square), Tanner and Lamb¹⁸ (∇), and Smith-Magowan and Wood¹⁷ (\circ). The black solid line corresponds to the equation 7. The red, blue, and green solid lines correspond to the Cp^0 values estimated using the revised HKF model at 0.1 MPa, 4 MPa, and 17.7 MPa, respectively.

Figure 5. Estimated values of Cp^0 for HCl(aq) using the Pitzer approach to correlate experimental Cp^0 values obtained by Patterson and coworkers²⁷ (\circ), Sharygin and Wood³⁰ (∇), and Tremaine and coworkers²⁸ (\triangle). The black solid line corresponds to the equation 8. The red, blue, and green solid lines correspond to the Cp^0 values estimated using the revised HKF model at 0.1 MPa, 4 MPa, and 28 MPa, respectively.

Figure 6. Average partial molar standard-state heat capacity for aqueous (a) Na⁺, (b) Cl⁻, and (c) NaCl (\blacksquare , values calculated in this work; \blacktriangle , values reported by Barner and Scheuerman³¹).

Figure 7. Apparent molar heat capacity of NaOH(aq) versus molality at 298.15 K (\square : experimental data from Magalhaes et al.¹¹; \circ : data from Patterson et al.²⁷; \triangle : data from Allred and Woolley²⁹; solid lines: Pitzer model for each data set using parameters reported in Table 9).

Figure 8. Estimated values of Cp^0 for NaOH(aq) using the Pitzer approach to correlate experimental Cp^0 values obtained by Patterson and coworkers²⁷ (\square), Simonson and coworkers³³ (\triangle), Conti and coworkers³⁴ (\circ), Allred and Woolley²⁹ (\diamond), and Magalhaes and coworkers¹¹ (∇). The black solid line corresponds to the equation 12. The red and blue solid lines correspond to the Cp^0 values estimated using the revised HKF model at 0.1 MPa and 4 MPa, respectively.

Figure 9. Estimated values of Cp^0 for the aqueous OH^- ion (■, calculated values from equation 10; solid line, equation 14).

Figure 10. Average partial molar standard-state heat capacity for aqueous (a) OH^- , and (b) NaOH (■, values calculated in this work; ▲, values reported by Barner and Scheuerman³¹).




Figure 1. A_j/R values as a function of temperature at a given pressure of 4.0 MPa.




Figure 2. Apparent molar heat capacity of NaCl(aq) versus molality at 348.15 K (squares: experimental data from Saluja et al.⁴; line: Pitzer approach performed in this study).





Figure 3. Apparent molar heat capacity of NaCl(aq) versus molality at (a) 298.15 K (\square , experimental data from Saluja et al.⁴; \bullet , data from Tanner and Lamb¹⁸) and (b) 318.15 K (\triangle , data from Perron et al.¹⁹; \bullet , data from Tanner and Lamb¹⁸).




Figure 4. Estimated values of Cp^0 for NaCl(aq) using the Pitzer approach to correlate experimental Cp^0 values obtained by Saluja and coworkers⁴ (\square), Tanner and Lamb¹⁸ (∇), and Smith-Magowan and Wood¹⁷ (\circ). The black solid line corresponds to the equation 7. The red, blue, and green solid lines correspond to the Cp^0 values estimated using the revised HKF model at 0.1 MPa, 4 MPa, and 17.7 MPa, respectively.




Figure 5. Estimated values of C_p^0 for HCl(aq) using the Pitzer approach to correlate experimental C_p^0 values obtained by Patterson and coworkers²⁷ (○), Sharygin and Wood³⁰ (▽), and Tremaine and coworkers²⁸ (△). The black solid line corresponds to the equation 8. The red, blue, and green solid lines correspond to the C_p^0 values estimated using the revised HKF model at 0.1 MPa, 4 MPa, and 28 MPa, respectively.






Figure 6. Average partial molar standard-state heat capacity for aqueous (a) Na^+ , (b) Cl^- , and (c) $NaCl$ (■, values calculated in this work; ▲, values reported by Barner and Scheuerman³¹).




Figure 7. Apparent molar heat capacity of NaOH(aq) versus molality at 298.15 K (□: experimental data from Magalhaes et al.¹¹; ○: data from Patterson et al.²⁷; △: data from Allred and Woolley²⁹; solid lines: Pitzer model for each data set using parameters reported in Table 9).




Figure 8. Estimated values of Cp^0 for NaOH(aq) using the Pitzer approach to correlate experimental Cp^0 values obtained by Patterson and coworkers²⁷ (\square), Simonson and coworkers³³ (\triangle), Conti and coworkers³⁴ (\circ), Allred and Woolley²⁹ (\diamond), and Magalhaes and coworkers¹¹ (∇). The black solid line corresponds to the equation 12. The red and blue solid lines correspond to the Cp^0 values estimated using the revised HKF model at 0.1 MPa and 4 MPa, respectively.




Figure 9. Estimated values of C_p^0 for the aqueous OH^- ion (■, calculated values from equation 10; solid line, equation 14).




Figure 10. Average partial molar standard-state heat capacity for aqueous (a) OH^- , and (b) NaOH (■, values calculated in this work; ▲, values reported by Barner and Scheuerman³¹).