




RESEARCH ARTICLE OPEN ACCESS

Effect of the Consumption of Lean Red Meat from Beef (Pirenaica Breed) Versus Lean White Meat (Chicken) on the Gut Microbiota: A Randomized Cross-Over Study in Healthy Young Adults

Isabel Rueda-De Torre^{1,2,3} | Julio Plaza-Díaz^{3,4,5,6}  | María L. Miguel-Berges^{1,2,3,7}  | Angel Gil^{3,8,9,10} | Laura Grasa^{2,7,11}  | Luis Mariano Esteban^{12,13} | Sergio Sabroso-Lasa¹⁴ | María M. Campo^{7,15} | Pilar Santolaria^{15,16} | Luis A. Moreno^{1,2,3,7} | Alba M. Santaliestra-Pasías^{1,2,3,7}

¹GENUD (Growth, Exercise, Nutrition and Development) Research Group, Faculty of Health Sciences, Universidad de Zaragoza, Zaragoza, Spain | ²Instituto de Investigación Sanitaria de Aragón (IIS Aragón), Zaragoza, Spain | ³Centro de Investigación Biomédica en Red de Fisiopatología de la Obesidad y Nutrición (CIBEROBN), Madrid, Spain | ⁴ANUT-DSM (Alimentació, Nutrició Desenvolupament i Salut Mental), Departament de Bioquímica i Biotecnologia, Universitat Rovira i Virgili, Reus, Spain | ⁵School of Health Sciences, Universidad Internacional de La Rioja, Logroño, Spain | ⁶Institut D'Investigació Sanitària Pere Virgili (IISPV), Reus, Spain | ⁷Instituto Agroalimentario de Aragón (IA2), Universidad de Zaragoza-CITA, Zaragoza, Spain | ⁸Department of Biochemistry and Molecular Biology II, School of Pharmacy, University of Granada, Granada, Spain | ⁹Institute of Nutrition and Food Technology "José Mataix", University of Granada, Armilla, Granada, Spain | ¹⁰Instituto de Investigación Biosanitaria de Granada (IBS.Granada), Complejo Hospitalario Universitario de Granada, Granada, Spain | ¹¹Departamento de Farmacología, Fisiología y Medicina Legal y Forense, Facultad de Veterinaria, Universidad de Zaragoza, Zaragoza, Spain | ¹²Escuela Politécnica de La Almunia, Universidad de Zaragoza, Zaragoza, Spain | ¹³Biocomputation and Physics of Complex Systems, Universidad de Zaragoza, Zaragoza, Spain | ¹⁴Genetic and Molecular Epidemiology Group (GMEG), Spanish National Cancer Research Centre (CNIO), Madrid, Spain | ¹⁵Departamento de Producción Animal y Ciencia de los Alimentos, Universidad de Zaragoza, Zaragoza, Spain | ¹⁶Instituto Universitario de Ciencias Ambientales, Universidad de Zaragoza, Zaragoza, Spain

Correspondence: María L. Miguel-Berges (mlmiguel@unizar.es)

Received: 23 December 2024 | **Revised:** 3 July 2025 | **Accepted:** 9 July 2025

Funding: The project DIETAPYR2 (Ref. EFA 144/16) was supported by the European Regional Development Fund (Fondo Europeo de Desarrollo Regional, FEDER) at 65% through the INTERREG V-A España-Francia-Andorra Programme (POCTEFA 2014–2020) created by the European Union.

Keywords: diet intervention | gut microbiota | red meat | white meat | young adults

ABSTRACT

Limited evidence exists regarding the impact of meat consumption on the human gut microbiota, with factors such as animal source, production system, and cooking methods often overlooked. This study evaluates the effect of Pirenaica breed beef or conventional chicken-based diets on the gut microbiota of healthy adults. A randomized cross-over controlled trial with two 8-week periods, separated by a 5-week washout, is carried out. Participants consume either Pirenaica breed beef or chicken three times per week with their diet. Stool samples are collected at the beginning and end of each period. Gut microbiota is analyzed via amplification and sequencing of V3–V4 regions of 16S rRNA. Alpha diversity and relative abundances at phylum and genus levels are calculated. Sixteen participants are included (mean age 20.12 ± 2.36 years). Both diets induce modest changes in microbial composition, with no significant differences between groups. At the phylum level, *Bacillota* increased, while *Synergistota*, *Chloroflexota*, and *Pseudomonadota* decreased. Alpha diversity parameters declined significantly after the chicken-based diet, although overall reduction in microbial diversity was observed across both interventions. The consumption of lean red meat or lean white meat as part of habitual diet produces similar effects on the gut microbiota.

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2025 The Author(s). *Molecular Nutrition & Food Research* published by Wiley-VCH GmbH.

1 | Introduction

The gut microbiota has emerged as a key factor in human health, as evidence indicates that intestinal bacteria play a role in disease etiology [1]. Similarly, diet is recognized as a potent modulator of microbiome composition, diversity, and metabolic activity [2]. Understanding how diet influences microbial communities could provide insights into the mechanisms underlying the association between food consumption and health outcomes [3]. However, it is critical to note that the quality of food and its nutrients rather than quantity, should be considered the most important factors [4].

Meat consumption has been in the spotlight in recent years as it has been criticized from ethical, environmental, and health perspectives [5]. Both unprocessed and processed red meat consumption have been reported to be risk factors for cardiovascular diseases and type 2 diabetes mellitus [6]. In this context, it is crucial to understand the biological function of the nutrients included in a variety of types of meat, which have different effects on health [7]. There is no consistent evidence of the association between meat consumption and gut microbiota-related outcomes in humans. A systematic review of meat consumption and the gut microbiota revealed that bacterial composition may be affected differently at lower taxonomic levels, such as genus or species level, or according to overall alpha diversity [8]. Moreover, most of these studies did not consider factors such as animal sources, feeding, processing, or cooking methods, which could be crucial for determining the nutritional profile of meat products [9–11].

In recent years, consumption patterns for animal-derived products have evolved considerably. In addition to their flavor, shape, and tenderness, consumers are particularly interested in their origin, potential health effects, and sustainable production methods [12]. Rural areas contribute to the promotion of local agriculture and livestock farming systems, providing an optimal environment for sustainable meat production [12]. The Pirenaica breed is an autochthonous beef breed closely linked to the Pyrenees Mountains. It follows an extensive husbandry system model and is raised on local feedstuffs and collective grazing lands. Livestock farming systems in some specific areas are based on sustainable and local practices. It is crucial to provide safe and high-quality meat, which will have positive effects on consumers' health.

The present study aimed to evaluate the effect of the consumption of lean red meat from beef (Pirenaica breed) in healthy young adults on their gut microbiota composition, compared with lean white meat (chicken).

2 | Experimental Section

2.1 | Study Design

This was a sub-study in the context of the DIETAPYR2 study (*Innovaciones aplicadas a la cadena productiva pirenaica de vacuno para valorizar una carne identificable por el consumidor*). This was a randomized crossover controlled trial (NCT04832217, clinicaltrials.gov) that included two experimental periods lasting

8 weeks. Participants were randomly assigned either a Pirenaica breed beef or a conventional chicken-based diet. They were instructed to consume the allocated products three times per week within their usual diet. A 5-week washout period was implemented between the two intervention phases to minimize residual effects from the previous diet, and to allow participant's baseline status to be restored before starting the second phase.

Figure 1 shows the study design. Two weeks before the beginning of the study, participants were contacted and informed consent was obtained. At the first visit, medical history, sociodemographic, and lifestyle behavior questionnaires were completed. Additionally, anthropometric, blood pressure (systolic and diastolic blood pressure) and heart rate measurements were taken. Following an overnight 12-h fast, blood was drawn. As a voluntary test of participation, stool samples were collected.

At the initial visit, participants were randomly assigned to a beef (Pirenaica breed) (Intervention Group) or chicken-based diet (Control Group). After the first experimental period, participants attended the second visit in the afternoon and the following morning, during which blood extraction and stool samples were collected under the same conditions. Following the crossover design, participants were switched to an alternative diet, either a chicken or Pirenaica breed-based diet, for the second 8-week period. Detailed experimental explanations were previously published [13].

2.2 | Participants

The participants were healthy young adults over the age of 18 years with no chronic, metabolic, endocrine, or nutrition-related diseases. Volunteers who completed stool sample collection at all four time points were included in this sub-study. Three university accommodation halls, two of them in Huesca (Colegio Mayor Universitario Ramón Acín and Residencia Misioneras del Pilar) and one in Zaragoza (Residencia Baltasar Gracián) (Spain), were the recruitment settings. The study protocol was approved by the Research Ethics Committee of the Government of Aragón (Spain) (N° 17/2018, 11 October 2018). All procedures were performed following the ethical guidelines of the Edinburgh revision of the 1964 Declaration of Helsinki (2024) [14].

2.3 | Dietary Intervention

Participants were urged to follow their usual diet and randomly assigned to consume allocated food products three times per week. The nutritional value of both diets, including protein and fats sources, was similar in both groups. Participants consumed 150 g of boneless chicken or Pirenaica breed beef (200 g with bones). The Pirenaica breed beef was obtained from entire young bulls that were raised through an extensive husbandry farming system for approximately 6 months. They were fed using cereal-based concentrates and cereal straw until they reached approximately 13.5 months of age. The nutritional composition of the Pirenaica breed beef has been previously analyzed [15]. The meat from Pirenaica breed young bulls is lean and is a




FIGURE 1 | Study design. A total of 16 participants were randomly assigned to start with either the Pirenaica breed beef or chicken-based diet for 8 weeks, followed by a 5-week washout period, after which they changed to the alternative diet for another 8 weeks. Stool samples were collected at the beginning and end of each period.

source of potassium and phosphorus, high in protein, zinc, and vitamins B3, B6, and B12, and low in sodium [15]. The chicken meat was sourced from conventional poultry farms. Meat cuts included loin, silverside and brisket for beef and thigh, drumstick, and breast. There were no analysis of macronutrient and micronutrient intakes between accommodation halls since the menus were similar. To ensure the same conditions, the study products were served during lunchtime and each chef in the three student accommodation halls was provided with cooking methods instructions. The cooking methods included braiding, stewing, and grilling for beef and braiding, stewing and roasting for chicken [13]. Dietary assessment was conducted for each participant using a food frequency questionnaire, at the beginning and end of each 8-week period. The questionnaire was previously validated [16, 17]. To evaluate overall dietary habits beyond the assigned products and account for potential variability in total intake, the Diet Quality Index (DQI) [18] was also calculated for each participant. This composite index is widely used to assess overall quality of the diet and includes three components: dietary quality, dietary diversity, and dietary equilibrium. A detailed description of the index is available in the previously published manuscript based on the same cohort [19].

2.4 | Stool Sample Collection

Stool samples were collected by participants at the beginning and end of each intervention period. The samples were frozen at -20°C immediately after collection and stored at -80°C until analysis.

2.5 | Extraction of DNA

Bacterial DNA was extracted from frozen samples using QIAamp Fast DNA Stool Mini Kit (Qiagen, Hilden, Germany) and following the manufacturer's instructions. Fecal samples were mixed with 1 mL InhibitEX buffer in SK38 tubes and processed by using the Precellys 24 homogenizer for 2×30 s at 6500 rpm and 10 s delay between cycles. Lysis was completed at 95°C for 5 min. Finally, DNA was eluted in 40 μL elution ATE buffer. Once DNA was extracted, DNA concentrations were measured with Qubit 4.0 fluorometer (Invitrogen) and dsDNA HS (high sensitivity) Assay Kit (Invitrogen). DNA purity was assessed by measuring the A260/A280 with NanoDrop ND-1000 Spectrophotometer V3.0.1 (Thermo Scientific Waltham, MA, USA) [20].

2.6 | Sequence Analysis and Bioinformatics

The extracted DNA was amplified by PCR using specific primers for 16S, targeting the V3 and V4 hypervariable regions of the bacterial 16S rRNA gene [21], as well as a primer for dimer cleanup. Using the Illumina Novaseq 6000 platform and 250 PE, the libraries were sequenced.

The primer sequences for 341F and 806R are commonly used to amplify the V3-V4 region of the bacterial 16S rRNA gene. The 341F forward primer sequence is 5'-CCTACGGGAGGCAGCAG-3' and the 806R reverse primer sequence is 5'-GGACTACHVGGGTWTCTAAT-3'. To confirm that there was no contamination, a negative control containing water was obtained. Using Illumina bcl2fastq2 Conversion

Software v2.20, the raw sequences were demultiplexed, and raw data was imported into QIIME 2 2023.7 open-source software [22] using the q2-tools-import script which uses Paired-end with Quality input format. The denoising process was performed with DADA2 [23], which uses a quality-aware model of Illumina amplicon errors to obtain a distribution of sequence variances, each differing by one nucleotide. To truncate and trim the forward reads at positions 240 and 7, the q2-dada2-denoise script was used following the retrieval of the quality scores. After truncating the reverse reads at position 240, we trimmed them at position 7. We removed chimeras using the consensus filter, which detects chimeras in samples individually and removes those found in a sufficient number of samples. In addition, forward and reverse reads are merged during this step. We constructed phylogenies using FASTTREE2 (via q2-phylogeny) [24], using all amplicon sequence variants (ASVs) aligned with MAFFT [25] via q2-alignment. With a similarity threshold of 99%, a naive Bayes taxonomy classifier was employed (via q2-feature-classifier) [26] against the SILVA 16S V3-V4 v138_99 [27]. The data filtering process excluded samples with fewer than 10,000 reads. The diversity of the samples was examined using the vegan library of R programming language version 4.2.2 (The R Foundation for Statistical computing, Vienna, Austria) [28]. In the present study, alpha diversity indices such as Fisher [29], Shannon [30], Simpson, inverse Simpson [31], and species richness indices, as well as Pielou's evenness index were examined [32].

2.7 | Functional Profiles

To explore the functional profiles of the sequenced samples, the PICRUST2 tool was employed [33]. This method infers metagenomic content by placing identified phylotypes into a reference phylogenetic tree built from approximately 20,000 full-length 16S rRNA gene sequences obtained from the Integrated Microbial Genomes (IMG) database. Functional characteristics of these genomes were performed using Clusters of Orthologous Groups of Proteins (COG) and the Enzyme Commission numbers (EC) databases. To reconstruct metabolic pathways, EC numbers were mapped to MetaCyc reactions, allowing the inference of MetaCyc pathway abundances. These were calculated as the harmonic mean of the contributing key reactions within each sample. Final pathway predictions were adjusted for 16S rRNA gene copy number and scaled to the relative abundance of each phylotype, providing estimated abundances for each predicted gene family [20].

2.8 | Statistical Analysis

The sample size for the global study was calculated assuming a two-tailed alpha error of 0.05, with a statistical power of 90% and a 20% loss to follow-up; the required sample size was 60 participants. Due to the exploratory nature of the results of gut microbiota data, combined with the lack of previous research and limitations in reproducibility across studies, it proved challenging to perform an a priori power calculation for this sub-study. The power calculation was performed for the estimated differences between the intervention and the control groups, along with the variance calculation of the crossover study design. Sociodemo-

graphic data are reported as mean \pm standard deviation (SD) for continuous variables or as the number of cases and percentages for categorical variables. Baseline differences were calculated with the Student's t-test for continuous variables or the Chi² test for categorical variables. All quantitative data were tested for normality assumptions with the Shapiro-Wilk test. The gut microbiota composition was analyzed for the relative abundance at the phylum and genus levels. The updated taxonomic designations for *Actinobacteria*, *Bacteroidetes*, *Chloroflexi*, *Firmicutes*, *Proteobacteria*, and *Synergistetes* are referred to as *Actinomycetota*, *Bacteroidota*, *Chloroflexota*, *Bacillota*, *Pseudomonadota*, and *Synergistota*, respectively [34]. To assess differences between the beginning and the end of the experimental period in the relative abundance of bacteria (phylum and genus levels), as well as for the alpha diversity indices, a paired Wilcoxon signed rank test was applied. A linear mixed-effects model for repeated measures was applied to evaluate intervention effects. The model included the intervention/control condition, period, sequence, and DQI values as fixed effects, with participants as random effects. The dependent variable was defined as the difference between the post- and pre-intervention values for each individual. This was to account for within-subject correlations and repeated measures over time. This approach is particularly suitable for crossover studies with small sample sizes, as it improves statistical power and controls for individual variability [35]. We conducted a robust linear mixed model analysis using the nlme package as it employs modern, efficient linear algebra methods, avoiding the constraints of homogeneity of variances, sphericity, and other distributional assumptions. To better illustrate the control and intervention effects, the intercept was set to zero, and the random forest displayed the beta coefficients along with their 95% confidence intervals. Statistical analyses were conducted in R Studio (Integrated Development Environment for R. Posit Software, PBC, Boston, MA) with the nlme and forest libraries and the Statistical Package for the Social Sciences (SPSS Version 25; SPSS, Chicago, IL, USA), with visualizations generated in GraphPad Prism (Version 9). Based on a Rivera-Pinto analysis, it is possible to identify microbial signatures, that is, groups of microbes that can predict particular phenotypes of interest. Based on an individual's unique microbiota, these microbial signatures may be used to diagnose, prognosticate, or predict therapeutic responses. To identify microbial signatures, we modelled the response variable as well as selected the taxa that yield the highest level of classification or prediction accuracy. As part of the Rivera-Pinto method and Selbal algorithm, we evaluated specific signatures at the phylum and genus levels to select a sparse model that adequately explains the response variable. The microbial signatures are calculated using geometric means based on data collected from two groups of taxa. The name implies that these groups are those with relative abundances, or balances, that are related to the response variable of interest [36].

3 | Results

Among the 52 participants who initially agreed to participate, 47 completed the study with acceptable follow-up compliance. Data from 16 participants who collected the stool samples at all four study points were ultimately analyzed for this sub-study. The mean age of participants was 20.16 years (\pm 2.43), 68.8% females. The average body mass index (BMI) was

23.75 kg/m² (\pm 2.68). No significant differences in sociodemographic characteristics, including age, BMI, or parental education, were observed between participants who began the study with the Pirenaica breed beef-based diet and those who started with the chicken-based diet. The DQI was assessed at the beginning and end of each intervention period. Baseline DQI values were comparable between groups. However, a significant reduction was observed after chicken-based diet ($p = 0.009$). The overall variation between both groups was also statistically significant ($p = 0.0025$) (see Table S1). Additionally, the baseline gut microbiota profile for all participants is provided in Table S2.

Table 1 presents the phylum-based abundances and alpha diversity indices, including the median with interquartile range before and after each intervention period. Variation and mean differences between the experimental periods were also considered. After 8 weeks of intervention, the Pirenaica breed beef-based diet group exhibited a modest decrease in the relative abundance of the phyla *Bacteroidota*, *Chloroflexota*, *Pseudomonadota*, and *Synergistota*, along with an increase in the relative abundance of *Actinomycetota*. The dominant phylum, *Bacillota*, remained relatively stable. However, these changes did not reach statistical significance ($p < 0.05$), with the exception of *Chloroflexota* ($p = 0.043$), suggesting stability across in response to the dietary interventions. Additionally, no significant changes were observed in alpha diversity measures, including species richness, Fisher's alpha, Shannon, Pielou's evenness, Simpson, and the Inverse Simpson indices (Figure 2). A decrease in the relative abundance of the phyla *Actinomycetota*, *Bacteroidota*, *Chloroflexota*, *Pseudomonadota*, and *Synergistota* was observed after the chicken-based diet, with statistically reductions in *Chloroflexota* ($p = 0.003$) and *Synergistota* ($p = 0.002$). Remarkably, the chicken-based diet group exhibited a significant decrease in richness ($p = 0.005$), Fisher ($p = 0.005$) and Shannon indices ($p = 0.034$). When comparing the variations observed before and after each intervention in both dietary groups, no statistically significant differences were observed at the phylum level or in the alpha diversity indices. Figure 3 shows the relative abundances at the phylum level of stool samples provided by all participants at the beginning and end of the two periods of the study.

At the genus level (see Table S3), a significant increase in the relative abundance of *Blautia* ($p = 0.044$) was observed after the Pirenaica breed beef-based diet, while *Coproccoccus* ($p = 0.044$), *Eubacterium hallii* group ($p = 0.049$), and *Roseburia* ($p = 0.013$) showed a significant decrease. The chicken-based diet group exhibited significant decreases in the relative abundance of *Lachnospiraceae* NK4A136 group ($p = 0.010$), *Lachnospira* ($p = 0.049$), *Eubacterium eligens* group ($p = 0.009$), *Adlercreutzia* ($p = 0.006$), *GCA-900066575* ($p = 0.039$), *Saccharofermentans* ($p = 0.047$), *Thermovirga* ($p = 0.003$), and *Bacteroides* ($p = 0.048$). Conversely, significant increases were observed in *Sphingomonas* ($p = 0.005$), *Family XIII AD30II* group ($p = 0.004$), and *Ruminococcaceae* UCG-009 ($p = 0.034$). Interestingly, *Blautia* also showed a significant increase in this groups ($p = 0.034$).

Table 2 summarizes the findings from the linear mixed model analysis, which assessed the effects of dietary interventions (Pirenaica breed-beef or chicken-based diets), period, sequence, and DQI values at the phylum-level abundances and alpha diver-

sity indices. After adjusting for DQI, both dietary interventions continued significantly influenced specific phyla, with consistent directional trends. Both diets were associated with an increase in *Bacillota*, and reductions in *Chloroflexota* and *Synergistota*, with all changes remaining statistically significant ($p < 0.005$). Regarding alpha diversity, both interventions led to significant reductions in richness, Fisher's index, and the Shannon index; a significant reduction in the Inverse Simpson index was observed only following the chicken-based diet. Significant period-related effects were observed for several variables. During the second period, the relative abundance of *Bacillota* decreased ($p = 0.005$) and significant increases in *Synergistota* ($p < 0.001$) and *Chloroflexota* ($p < 0.001$). Alpha diversity indices showed notable variability between periods, with significant increases recorded in richness ($p = 0.007$), Fisher's index ($p = 0.007$), Shannon ($p = 0.030$) and the Inverse Simpson index ($p = 0.015$) during the second period compared to the first. A sequence effect was observed only for the Shannon index ($p = 0.015$) showed significant reductions associated with the sequence in which diets were administered.

3.1 | Bacteria Metabolic Pathways

The predicted metabolic profiles of the sequenced samples were assessed, focusing on pathways related to amino acid and polyamine metabolism, lipid metabolism, energy production, and fermentation. Overall, the chicken-based diet was associated with a broader reduction in microbial functional capacity, while the Pirenaica breed beef-based diet induced more limited changes. Following the Pirenaica breed beef-based diet, microbial functional activity remained relatively stable, except to L-methionine biosynthesis pathways ($p = 0.026$) and L-methionine biosynthesis II ($p = 0.017$). The chicken-based diet induced significant reductions in pathways related to aromatic amino acid biosynthesis ($p = 0.026$), fatty acid biosynthesis initiation ($p = 0.010$), and gluconeogenesis ($p = 0.007$). Moreover, reduced activity was observed in L-isoleucine degradation ($p = 0.015$) and fatty acid biosynthesis ($p = 0.004$) (see Table S3).

3.2 | Rivera-Pinto Analysis

Based on the geometric mean of data derived from two groups of taxa whose relative abundances correlate with the response variable, the Rivera-Pinto method [36] is used to determine a microbial signature. In the chicken-based diet group, *Coproccoccus*, *Eubacterium eligens* group, and *Eubacterium hallii* group genera were most closely associated with the baseline (Figure 4A). A higher balance score was associated with a significant relative abundance of the *Coproccoccus*, *Eubacterium eligens* group, and *Eubacterium hallii* group genera. This was in comparison to the *Sphingomonas*, *Streptococcus*, *Collinsella*, and *Ruminoclostridium* 5 genera. The *Coproccoccus* 2 and *Agathobacter* genera, as well as the *Bacteroidota* phylum, were most closely associated with baseline in the Pirenaica breed beef-based diet group (Figure 4B). Significant relative abundances of *Coproccoccus* 2, *Bacteroidota*, and *Agathobacter* were associated with a higher balance score. Comparatively to the *Phascolarctobacterium* and *Lachnospira* genera and the family *Ruminococcaceae*. In the comparison between the chicken-based diet group and Pirenaica breed

TABLE 1 | Relative abundances at phylum level and alpha diversity indices in the intervention and control groups.

	Pirenaica breed beef-based diet group, $n = 16$					Chicken-based diet group, $n = 16$					Mean differences in the variations between both groups		
	Before		After		Δ	p^a		Before		Δ	p^a	Mean difference	CI 95%
	Before	After	Before	After				Before	After				
Sequences	213,560 (203,841–217,903)	212,831 (209,374–216,142)	592.5	0.959				206,240 (203,103–209,348)	209,374 (201,483–214,603)	2447.6	0.301	1855.125	–19,360.693; 23,070.9429
Phylum													
<i>Actinomyces</i>	1.86 (1.26–2.56)	2.11 (1.43–2.57)	–0.02	0.938				3.05 (2.63–3.57)	2.54 (1.81–3.53)	–0.36	0.501	–0.35	–1.93; 1.24
<i>Bacteroidota</i>	0.04 (0.02–0.08)	0.01 (0.00–0.11)	–0.02	0.222				0.03 (0.01–0.12)	0.02 (0.00–0.05)	–0.03	0.109	–0.01	–0.05; 0.03
<i>Chloroflexota</i>	0.00 (0.00–0.59)	0.00 (0.00–0.00)	–0.36	0.043				1.25 (0.16–1.53)	0.00 (0.00–0.16)	–0.82	0.003	–0.46	–1.15; 0.22
<i>Bacillota</i>	97.45 (94.87–98.07)	97.49 (96.84–98.35)	1.21	0.605				92.08 (90.76–94.71)	96.44 (93.60–97.40)	1.98	0.134	0.77	–3.02; 4.56
<i>Pseudomonadota</i>	0.10 (0.02–0.75)	0.07 (0.00–0.11)	–0.25	0.112				0.77 (0.15–1.07)	0.29 (0.10–0.68)	0.48	0.733	0.73	–0.57; 2.02
<i>Synergistota</i>	0.00 (0.00–0.62)	0.00 (0.00–0.00)	–0.41	0.068				1.51 (0.32–2.14)	0.00 (0.00–0.00)	–1.18	0.002	–0.77	–1.65; 0.11
Alpha diversity indices													
Richness	71.00 (66.00–79.25)	70.00 (66.25–75.00)	–5.13	0.737				98.00 (79.75–101.00)	71.00 (62.25–91.75)	–16.75	0.005	–11.36	–25.06; 1.80
Fisher's index	11.26 (10.33–12.84)	11.08 (10.38–12.02)	–1.01	0.717				16.56 (12.95–17.17)	11.26 (9.64–15.33)	–3.24	0.005	–2.23	–4.87; 0.40
Shannon index	3.40 (3.30–3.55)	3.32 (3.11–3.50)	–0.12	0.098				3.53 (3.40–3.68)	3.41 (3.05–3.65)	–0.18	0.034	–0.07	–0.22; 0.09
Pielou's evenness	0.79 (0.78–0.81)	0.78 (0.76–0.82)	–0.02	0.109				0.77 (0.76–0.80)	0.79 (0.73–0.80)	–0.01	0.877	0.01	–0.001; 0.04
Simpson index	0.93 (0.92–0.95)	0.92 (0.90–0.94)	–0.02	0.148				0.93 (0.92–0.95)	0.93 (0.88–0.95)	–0.02	0.148	–0.002	–0.02; 0.02
Inverse Simpson	14.82 (12.30–19.36)	12.53 (10.09–16.90)	–1.85	0.134				14.81 (11.80–20.51)	15.00 (8.63–18.87)	–1.83	0.098	0.01	–2.74; 2.77

Abbreviation: SD, standard deviation.

Note: Data are expressed as median and interquartile range; Δ : mean change (final-beginning) of each diet group.^aMultivariate contrast between the beginning and end of each corresponding food product (Pirenaica breed beef, $n = 16$ or chicken, $n = 16$).^bMultivariate contrasts between the variations (final-beginning) of both diet groups (Pirenaica breed beef or chicken). Significant differences ($p < 0.05$).




FIGURE 2 | Alpha diversity indices of the intervention and control groups. The data are expressed as the median and interquartile range according to the Wilcoxon test. Richness and Fisher's index, which quantify the relationship between the number and abundance of species, were calculated. For microbial diversity and evenness, Shannon, Pielou, Simpson, and inverse Simpson's indices were performed. Pirenaica breed beef-based diet group ($n = 16$); chicken-based diet group ($n = 16$). Box plots represent the interquartile range (25th–75th percentile); the line inside within each box indicates the median, and the whiskers extend to minimum and maximum values. * Indicates statistically significant differences ($p < 0.05$).

beef-based diet group at baseline, *Gracilibacter*, *Negativibacillus*, *Subdoligranulum*, and *Eubacterium hallii* group genera and the *Actinomycetota* phylum were most closely associated with the chicken-based diet group (Figure 4C). A higher balance score was associated with a significant relative abundance of *Gracilibacter*, *Negativibacillus*, *Subdoligranulum*, and *Eubacterium hallii* group genera and the *Actinomycetota* phylum. This was compared to the *Coprococcus* 2, *Lachnospiraceae* ND3007 group, and CAG56 genera. At the end of the intervention, the *Pseudomonadota* phylum and *Ruminococcaceae* UCG013 genus were most closely associated with the chicken meat group (Figure 4D). Significant relative abundances of *Pseudomonadota* phylum and *Ruminococcaceae* UCG013 genus were associated with a higher balance score. Comparatively to the *Ruminococcaceae* NK4A214 group, *Ruminococcaceae* uncultured, *Butyricicoccus*, *Negativibacillus*, and *Collinsella* genera.

4 | Discussion

In this study, the effects of an intensive intervention involving the inclusion of lean red meat from beef (Pirenaica breed) or lean white meat on gut microbiota composition and structure

were examined. The gut microbiota influences critical processes such as digestion, nutrient absorption, and immune modulation; dysbiosis has been associated to various metabolic, autoimmune, and gastrointestinal diseases. While meat consumption has long been regarded as a risk factor for cardiometabolic diseases, its relationship with the microbial ecosystem is increasingly recognized [37]. Most studies do not focus on meat intake but rather, more broadly on dietary patterns with or without meat and their impact on gut microbiota. The positive effects of certain food groups on the gut microbiota are well-documented. For instance, dairy products, among animal-origin foods, have been demonstrated to promote gut health by increasing the abundance of probiotic bacteria as *Bifidobacterium* and *Lactobacillus* [38]. Similarly, dietary fiber from fruits, vegetables and whole grains may influence the gut microbiota composition and function through the increase of specific genera within the phyla *Bacillota* and *Actinomycetota* [39, 40]. In contrast, the effect of meat consumption on gut microbiota has typically been examined in the context of a Western dietary pattern, characterized by increased animal fats, refined sugars, and animal proteins [41]. This dietary pattern leads to an increase in *Bacillota* and *Pseudomonadota*, while decreasing *Bacteroidota*, as well as the genera *Prevotella* and *Bifidobacterium* [42].




FIGURE 3 | Relative abundances at the phylum level by all participants at the beginning and end of the two periods of the study. Pirenaica breed beef-based diet group ($n = 16$); chicken-based diet group ($n = 16$). The stacked bar plots represent the composition of microbial communities for each sample.

TABLE 2 | Effects of the type of intervention, period, sequence, and Diet Quality Index (DQI) on gut microbiota composition and alpha diversity between both groups.

	Pirenaica breed beef-based diet		Chicken-based diet		Period		Sequence		Diet Quality Index	
	Coefficient	<i>p</i>	Coefficient	<i>p</i>	Coefficient	<i>p</i>	Coefficient	<i>p</i>	Coefficient	<i>p</i>
Phylum										
<i>Actinomycetota</i>	−4.01	0.098	−3.82	0.110	1.23	0.131	−0.31	0.692	0.07	0.155
<i>Bacteroidota</i>	0.04	0.632	0.04	0.632	0.01	0.687	−0.03	0.328	0.00	0.441
<i>Chloroflexota</i>	−1.22	0.049	−1.05	0.083	1.26	0.000	−0.17	0.392	0.00	0.928
<i>Bacillota</i>	12.99	0.013	11.39	0.025	−5.12	0.005	0.90	0.571	−0.18	0.074
<i>Pseudomonadota</i>	−3.68	0.119	−2.87	0.214	0.34	0.644	−0.41	0.597	0.08	0.134
<i>Synergistota</i>	−1.81	0.027	−1.78	0.028	1.61	0.000	−0.02	0.927	0.00	0.777
Alpha diversity indices										
Richness	−40.09	0.023	−41.44	0.019	21.61	0.007	−11.49	0.051	0.47	0.161
Fisher's index	−7.82	0.024	−8.04	0.020	4.23	0.007	−2.15	0.060	0.09	0.166
Shannon index	−0.66	0.013	−0.65	0.014	0.18	0.030	−0.22	0.015	0.01	0.049
Pielou's evenness	−0.06	0.211	−0.06	0.264	−0.01	0.665	−0.02	0.235	0.00	0.233
Simpson index	−0.08	0.056	−0.07	0.077	0.02	0.083	−0.02	0.094	0.00	0.144
Inverse Simpson	−5.94	0.195	−4.23	0.348	3.49	0.015	−2.73	0.096	0.05	0.606

Note: Data are expressed as coefficients and *p* values; the coefficients represent the estimated effect of dietary intervention (Pirenaica breed beef or chicken), period, sequence, and DQI on the relative abundances at the phylum level and alpha diversity indices.

Abbreviation: DQI, Diet Quality Index.

^aLinear mixed model for repeated measures ($n = 16$ per group) within individual to evaluate the effect of the type of intervention, period, sequence, and DQI. Significant differences ($p < 0.05$).




FIGURE 4 | Group balances are presented in an overview. Box plots illustrate the distribution of balance scores for before and after intervention in the chicken-based diet group (A), before and after intervention in the Pirenaica breed beef-based diet group (B), at the beginning of both diets (C), and after both diets (D). Pirenaica breed beef-based diet group ($n = 16$); chicken-based diet group ($n = 16$).

4.1 | Meat Consumption and Relative Abundance at the Phylum and Genus Levels

Our results revealed modest changes in microbial composition at the phylum level following both dietary interventions, with no statistically significant differences observed between the Pirenaica breed beef and chicken-based diet groups. Both diets were associated with a slight increase in the relative abundance of *Bacillota*, the dominant phylum, and a reduction in *Bacteroidota*. Notably, *Pseudomonadota* decreased in both groups, although its relative abundance remained consistently higher in participants who received the chicken-based diet. These trends were further supported by the linear mixed model analysis, accounting for type of dietary intervention, sequence, period, and DQI effects. After adjustments, both diet continued to influence microbial composition, with increases in *Bacillota* and reductions in *Synergistota* and *Chloroflexota* observed across interventions. These findings suggest that both meat diets exerted comparable, directional effects on the gut microbiota at the phylum level, regardless of diet quality (Figure 5).

At the genus level, *Blautia* increased in response to both interventions. This genus has previously associated with higher meat consumption [43, 44], and in some studies, with less favorable metabolic profile, although its remains unclear [8]. *Coprococcus*, *Roseburia*, and *Lachnospira* decreased after the interventions. These genera are well-established producers of short-chain fatty acids (SCFAs), particularly butyrate, and their abundance has been reported to decline in meat rich diets [45]. Although some genera changed differently depending on the meat consumed, no significant differences were found between groups, suggesting a similar overall impact on microbial composition.

Few studies have directly explored the effect of the type of meat consumed on the gut microbiota composition. In Chinese volunteers, bacterial taxa responses varied from our findings [46]. The relative abundances at both phylum and genus levels did not significantly change with meat-based diet replacement. After 2 weeks of a beef and chicken based diets, *Pseudomonadota* showed the most significant change during the meat replacement, increasing after a beef-based diet and decreasing




FIGURE 5 | Forest plots showing the effects of the intervention type at the phylum level abundances and alpha diversity indices. Relative abundances at the phylum level for the Pirenaica breed beef-based diet (A), relative abundance at the phylum level for the chicken-based diet (B), alpha diversity indices for the Pirenaica breed beef-based diet (C), and alpha diversity indices for the chicken-based diet (D). Data are presented as estimated coefficients and 95% confidence intervals obtained from linear mixed-effects models. Analysis was conducted using repeated measures ($n = 16$ per group). A negative coefficient indicates a reduction in the corresponding variable following the intervention, while a positive coefficient indicates an increase.

after a chicken-based diet [46]. An increase in *Pseudomonadota* abundance has been associated with dysbiosis, characterized by reduced mucous production in the gut barrier and low-grade inflammation [47]. At the genus level, our findings differ from this study, with reported increases in *Lachnospira*, *Lachnospiraceae* group NK4A136, and *Ruminococcus* 2 detected after consuming the beef-based diet [46]. A study compared the effect of consuming pork versus chicken within a healthy dietary pattern for 10 days in adults; a reduction in the abundance of *Bacteroidota* and the family *Bacteroidaceae* but an increase in the family *Christensenellaceae*, independent of pork or poultry intakes was observed [48]. However, no other types of meat, seafood, or whole eggs were allowed [48], and the short periods limits comparison with our findings. In an observational study, meat preferences caused significant variations in the gut microbiota composition between pork eaters and chicken eaters. Pork eaters showed a greater relative abundance of *Bacillota* and a lower relative abundance of *Bacteroidota* than chicken eaters [49]; these observations may not be attributed to red meat intake

in the absence of controlled full-feeding diets. Similarly, after 5 days of consuming red meat or mushrooms as part of an uncontrolled diet, the relative abundances of *Bacteroidota* and *Bacillota* decreased and increased, respectively [45]. However, it may be challenging to compare our results with those of previous studies due to methodological differences, including shorter interventions periods and the absence of washout periods between interventions.

4.2 | Meat Consumption and Alpha Diversity

After consuming both diets, there was a decrease in microbial diversity and richness, assessed by the Fisher, Shannon, and Simpson indices. However, statistically significant decreases were only observed following the chicken-based diet group, particularly in species richness, Fisher and the Shannon indices. This pattern suggests that the Pirenaica breed beef-based diet may exert a comparatively milder impact on overall microbial diversity

than the chicken-based diet. The analysis of intervention type, period, sequence, and DQI effects highlighted complex interactions modulating the gut microbiota. Notably, DQI significantly declined following the chicken-based diet, which may partially explain the observed microbial shifts. However, after adjusting for DQI, it did not emerge as significant predictor of alpha diversity parameters dietary habits during this phase may not have influenced responses. Interestingly, diversity indices increased during the second intervention period, pointing toward potential temporal or adaptive host-related responses. No consistent sequence effect was observed, supporting the adequacy of the crossover design and the effectiveness of the washout period. A previous study showed that the addition of unprocessed or processed lean red meat to a healthy lacto-ovo vegetarian diet for 3 weeks did not modify the overall microbiota structure, as assessed by metric values such as the Chao1, ACE, Shannon, and inverse Simpson indices [50]. The Chao and Shannon indices showed lower richness and diversity in pork eaters than in the chicken-eaters [49]. Furthermore, a randomized controlled trial revealed that fried meat (including chicken and fish) consumption decreased microbial community richness in overweight adults. At the phylum level, the ratio of *Bacillota*/*Bacteroidota* decreased in both the fried meat and control groups, with a significantly lower decrease in the control group, along with differences at the genus level [51]. High-heat cooking methods, such as frying and grilling, produce higher amounts of advanced glycation end-products [52]. Therefore, it is possible that the type of meat and the cooking method could modify the gut microbiota composition and structure.

4.3 | Functional Profiles

Regarding microbial metabolic pathways, a greater reduction was observed following the chicken-based diet, particularly in aromatic amino acid biosynthesis and gluconeogenesis pathways. However, the overall pattern of functional response was similar between both dietary interventions. These findings highlight the value of integrating microbial functional analysis in dietary intervention studies. Previous intervention comparing meta-rich diets to vegetarian diets have not reported difference in microbial functionally after 4 weeks of intervention [43].

Our study assessed the effects of different types of meat consumption on gut microbiota, taking into consideration an extensive meat production system. Young bulls from Pirenaica beef, raised under extensive husbandry conditions, is naturally lean (1.75–2.28 g fat/100 g), high in high-quality protein (23–24 g/100 g), and rich in micronutrients such as potassium, phosphorus, selenium, zinc, and B vitamins (B3, B6, B12), while being low in sodium [15]. These attributes support its inclusion in balanced into dietary patterns and justify its selection as a model of lean red meat for gut microbiota research. The results from this intervention have already been published, showing similar body composition parameters, lipid profiles, and fatty acid values after both intervention diets [13]. These findings suggest that the integration of lean red meat from beef (Pirenaica breed) into a healthy dietary pattern could positively influence not only the gut microbiota but

also some health indicators. Overall, 8 weeks of intensive nutritional intervention provided sufficient time for changes in the gut microbiota to occur. Indeed, studies have reported significant changes in microbiota composition after 3 weeks of intervention [53]. In this study, a 5-week washout period was established to reduce the influence of the previous intervention. The absence of a sequence effect across phylum-level abundances and alpha diversity parameters indicates that this period was adequate to re-establish baseline microbial conditions. However, the presence of period effect may reflect temporal adaptation, or that the diet consumed during this interval could have interacted with the outcomes.

4.4 | Strengths and Limitations

The study design (crossover clinical trial) is a notable strength as it provides a high level of evidence for comparisons of the outcomes between diets. The inclusion of an institutionalized young adult population minimized lifestyle variability, enhancing the result's validity. Moreover, the control of cooking methods across university accommodation halls is another strength to be considered. Furthermore, other interventions do not take into account the nutritional composition of the meat studied. In our study, the Pirenaica breed beef sourced from extensive husbandry practices yields a unique nutritional profile influenced by breed, feeding practices, sex, and age at slaughter [15].

The study also presented several limitations. It is important to note that this sub-study required voluntary participation, which may have limited the final sample size. To address this limitation, a linear mixed-effects model was applied for repeated measurements. This approach accounted for individual-level variability and enhanced the reliability of the statistical analysis, thereby improving the statistical power to detect effects. Although dietary intake was assessed using a validated food frequency questionnaire and overall dietary quality was considered using the DQI, the potential influence of unmeasured or residual dietary factors on gut microbiota parameters cannot be entirely excluded. Additionally, carry-over, period, or sequence effects could influence the effect of the intervention. The reported results are relevant to a particular type of meat, the Pirenaica breed beef, although this meat could be similar to others from similar husbandry conditions in extensive mountain production systems. However, when comparing our results with other studies, there is a lack of specificity in defining the term “meat”. The varied uses of these methods add to research evidence that is inconsistent and controversy [8].

5 | Conclusion

Overall, our results suggest that lean red meat from beef (Pirenaica breed) responds comparable to lean white meat from chickens in terms of gut microbiota composition and structure in healthy young adults. A more comprehensive analysis of the gut microbial community and its correlation with health parameters should be further considered.

Acknowledgments

The authors express their gratitude to the participants for their collaboration. We also appreciate Residencia Colegio Mayor Universitario Ramón Acín, Residencia Universitaria Misioneras del Pilar (Huesca), and Residencia Baltasar Gracian (Zaragoza) for their support and for providing the facilities to have access to the participants. We acknowledge the dedication of the kitchen teams from the residences that made it possible to carry out the study, and the Escuela de Hostelería y Turismo de Huesca for their collaboration and for designing the recipes used.

Consent

Informed consent was obtained from all subjects involved in the study.

Conflicts of Interest

The authors declare no conflicts of interest.

Data Availability Statement

Data available in article supplementary material.

References

1. A. Vijay and A. M. Valdes, "Role of the Gut Microbiome in Chronic Diseases: A Narrative Review," *European Journal of Clinical Nutrition* 76 (2022): 489–501.
2. R. K. Singh, H.-W. Chang, D. Yan, et al., "Influence of Diet on the Gut Microbiome and Implications for Human Health," *Journal of Translational Medicine* 15 (2017): 73.
3. C. J. Walsh, C. M. Guinane, P. W. O'Toole, and P. D. Cotter, "Beneficial Modulation of the Gut Microbiota," *FEBS Letters* 588 (2014): 4120–4130.
4. C. García-Montero, O. Fraile-Martínez, A. M. Gómez-Lahoz, et al., "Nutritional Components in Western Diet Versus Mediterranean Diet at the Gut Microbiota–Immune System Interplay. Implications for Health and Disease," *Nutrients* 13 (2021): 699.
5. S. Agarwal, K. R. McCullough, and V. L. Fulgoni, "Nutritional Effects of Removing a Serving of Meat or Poultry From Healthy Dietary Patterns—A Dietary Modeling Study," *Nutrients* 15 (2023): 1717.
6. W. Shi, X. Huang, C. M. Schooling, and J. V. Zhao, "Red Meat Consumption, Cardiovascular Diseases, and Diabetes: A Systematic Review and Meta-Analysis," *European Heart Journal* 44 (2023): 2626–2635.
7. K. Albracht-Schulte, T. Islam, P. Johnson, and N. Moustaid-Moussa, "Systematic Review of Beef Protein Effects on Gut Microbiota: Implications for Health," *Advances in Nutrition* 12 (2021): 102–114.
8. Y. Wang, C. N. Uffelman, R. E. Bergia, et al., "Meat Consumption and Gut Microbiota: A Scoping Review of Literature and Systematic Review of Randomized Controlled Trials in Adults," *Advances in Nutrition* 14 (2023): 215–237.
9. T. A. McAllister, K. Stanford, A. V. Chaves, P. R. Evans, E. Eustaquio De Souza Figueiredo, and G. Ribeiro, *Animal Agriculture* (Elsevier, 2020), 75–98.
10. M. M. Campo, E. Muela, J. L. Olleta, et al., "Influence of Cooking Method on the Nutrient Composition of Spanish Light Lamb," *Journal of Food Composition and Analysis* 31 (2013): 185–190.
11. M. Pistón, A. Suárez, V. Bühl, F. Tissot, J. Silva, and L. Panizzolo, "Influence of Cooking Processes on Cu, Fe, Mn, Ni, and Zn Levels in Beef Cuts," *Journal of Food Composition and Analysis* 94 (2020): 103624.
12. B. Chen, D. Li, D. Leng, H. Kui, X. Bai, and T. Wang, "Gut Microbiota and Meat Quality," *Frontiers in Microbiology* 13 (2022): 951726.
13. A. M. Santaliestra-Pasías, M. L. Miguel-Berges, M. M. Campo, et al., "Effect of the Intake of Lean Red-Meat From Beef (Pirenaica Breed) Versus Lean White-Meat on Body Composition, Fatty Acids Profile and Cardiovascular Risk Indicators: A Randomized Cross-Over Study in Healthy Young Adults," *Nutrients* 14 (2022): 3724.
14. World Medical Association, "World Medical Association Declaration of Helsinki: Ethical Principles for Medical Research Involving Human Subjects," *JAMA* 310 (2013): 2191–2194.
15. M. M. Campo, J. V. Romero, A. Guerrero, et al., "Nutrient Composition of Beef from the Pyrenees," *Journal of Food Composition and Analysis* 133 (2024): 106452.
16. J. M. Martín-Moreno, P. Boyle, L. Gorgojo, et al., "Development and Validation of a Food Frequency Questionnaire in Spain," *International Journal of Epidemiology* 22 (1993): 512–519.
17. C. de la Fuente-Arrillaga, Z. V. Ruiz, M. Bes-Rastrollo, L. Sampson, and M. A. Martínez-González, "Reproducibility of an FFQ Validated in Spain," *Public Health Nutrition* 13 (2010): 1364–1372.
18. K. Vyncke, E. Cruz Fernandez, M. Fajó-Pascual, et al., "Validation of the Diet Quality Index for Adolescents by Comparison With Biomarkers, Nutrient and Food Intakes: The HELENA Study," *British Journal of Nutrition* 109 (2013): 2067–2078.
19. M. L. Miguel-Berges, M. Fajó-Pascual, L. A. Moreno, et al., "Effect of Lean Red Meat From Beef (Pirenaica Breed) Versus Lean White Meat Consumption on Diet Quality: A Randomized-Controlled Crossover Study in Healthy Young Adults," *Nutrients* 15 (2022): 13.
20. J. Plaza-Díaz, M. Manzano, F. J. Ruiz-Ojeda, et al., "Intake of Slow-Digesting Carbohydrates is Related to Changes in the Microbiome and Its Functional Pathways in Growing Rats With Obesity Induced by Diet," *Frontiers in Nutrition* 9 (2022): 992682.
21. A. Klindworth, E. Pruesse, T. Schweer, et al., "Evaluation of General 16S Ribosomal RNA Gene PCR Primers for Classical and Next-Generation Sequencing-Based Diversity Studies," *Nucleic Acids Research* 41 (2013): 1.
22. E. Bolyen, J. R. Rideout, M. R. Dillon, et al., "Reproducible, Interactive, Scalable and Extensible Microbiome Data Science Using QIIME 2," *Nature Biotechnology* 37 (2019): 852–857.
23. B. J. Callahan, P. J. McMurdie, M. J. Rosen, A. W. Han, A. J. A. Johnson, and S. P. Holmes, "DADA2: High-Resolution Sample Inference From Illumina Amplicon Data," *Nature Methods* 13 (2016): 581–583.
24. M. N. Price, P. S. Dehal, and A. P. Arkin, "FastTree 2 – Approximately Maximum-Likelihood Trees for Large Alignments," *PLOS ONE* 5 (2010): 9490.
25. K. Katoh and D. M. Standley, "MAFFT Multiple Sequence Alignment Software Version 7: Improvements in Performance and Usability," *Molecular Biology and Evolution* 30 (2013): 772–780.
26. N. A. Bokulich, B. D. Kaehler, J. R. Rideout, et al., "Optimizing Taxonomic Classification of Marker-Gene Amplicon Sequences With QIIME 2's q2-Feature-Classifer Plugin," *Microbiome* 6 (2018): 90.
27. C. Quast, E. Pruesse, P. Yilmaz, et al., "The SILVA Ribosomal RNA Gene Database Project: Improved Data Processing and Web-Based Tools," *Nucleic Acids Research* 41 (2013): D590–596.
28. P. Dixon, "VEGAN, A Package of R Functions for Community Ecology," *Journal of Vegetation Science* 14 (2003): 927–930.
29. R. A. Fisher, A. S. Corbet, and C. B. Williams, "The Relation Between the Number of Species and the Number of Individuals in a Random Sample of an Animal Population," *Journal of Animal Ecology* 12 (1943): 42.
30. C. E. Shannon, "The Mathematical Theory of Communication. 1963," *M.D. Computing: Computers in Medical Practice* 14 (1997): 306–317.
31. E. H. Simpson, "Measurement of Diversity," *Nature* 163 (1949): 688–688.
32. E. C. Pielou, "The Measurement of Diversity in Different Types of Biological Collections," *Journal of Theoretical Biology* 13 (1966): 131–144.
33. G. M. Douglas, V. J. Maffei, J. R. Zaneveld, et al., "PICRUST2 for Prediction of Metagenome Functions," *Nature Biotechnology* 38 (2020): 685–688.
34. A. Oren and G. M. Garrity, "Valid Publication of the Names of Forty-Two Phyla of Prokaryotes," *International Journal of Systematic*

and *Evolutionary Microbiology* 71 (2021): 005056, <https://doi.org/10.1099/jjsem.0.005056>.

35. J. Verbeeck, M. Geroldinger, K. Thiel, et al., "How to Analyze Continuous and Discrete Repeated Measures in Small-Sample Cross-Over Trials?," *Biometrics* 79 (2023): 3998–4011.

36. J. Rivera-Pinto, J. J. Egozcue, V. Pawlowsky-Glahn, R. Paredes, M. Noguera-Julian, and M. L. Calle, "Balances: A New Perspective for Microbiome Analysis," *mSystems* 3 (2018): e00053-18.

37. L. E. Olofsson and F. Bäckhed, "The Metabolic Role and Therapeutic Potential of the Microbiome," *Endocrine Reviews* 43 (2022): 907–926.

38. H. Aslam, W. Marx, T. Rocks, et al., "The Effects of Dairy and Dairy Derivatives on the Gut Microbiota: A Systematic Literature Review," *Gut Microbes* 12 (2020): 1799533.

39. P. Cronin, S. A. Joyce, P. W. O'Toole, and E. M. O'Connor, "Dietary Fibre Modulates the Gut Microbiota," *Nutrients* 13 (2021): 1655.

40. K. Makki, E. C. Deehan, J. Walter, and F. Bäckhed, "The Impact of Dietary Fiber on Gut Microbiota in Host Health and Disease," *Cell Host & Microbe* 23 (2018): 705–715.

41. I. J. Malesza, M. Malesza, J. Walkowiak, et al., "High-Fat, Western-Style Diet, Systemic Inflammation, and Gut Microbiota: A Narrative Review," *Cells* 10 (2021): 3164.

42. A. Beam, E. Clinger, and L. Hao, "Effect of Diet and Dietary Components on the Composition of the Gut Microbiota," *Nutrients* 13 (2021): 2795, <https://doi.org/10.3390/nu13082795>.

43. E. Kohnert, C. Kreutz, N. Binder, et al., "Changes in Gut Microbiota After a Four-Week Intervention With Vegan vs. Meat-Rich Diets in Healthy Participants: A Randomized Controlled Trial," *Microorganisms* 9 (2021): 727.

44. L. A. Bolte, A. Vich Vila, F. Imhann, et al., "Long-Term Dietary Patterns Are Associated With Pro-Inflammatory and Anti-Inflammatory Features of the Gut Microbiome," *Gut* 70 (2021): 1287–1298.

45. J. Hess, Q. Wang, T. Gould, and J. Slavin, "Impact of *Agaricus bisporus* Mushroom Consumption on Gut Health Markers in Healthy Adults," *Nutrients* 10 (2018): 1402.

46. D. Zhao, K. Shan, Y. Xie, et al., "Body Weight Index Indicates the Responses of the Fecal Microbiota, Metabolome and Proteome to Beef/Chicken-Based Diet Alterations in Chinese Volunteers," *NPJ Biofilms Microbiomes* 8 (2022): 56.

47. N.-R. Shin, T. W. Whon, and J.-W. Bae, "Proteobacteria: Microbial Signature of Dysbiosis in Gut Microbiota," *Trends in Biotechnology* 33 (2015): 496–503.

48. S. Dhakal, Z. Moazzami, C. Perry, and M. Dey, "Effects of Lean Pork on Microbiota and Microbial-Metabolite Trimethylamine-N-Oxide: A Randomized Controlled Non-Inferiority Feeding Trial Based on the Dietary Guidelines for Americans," *Molecular Nutrition & Food Research* 66 (2022): 2101136.

49. J. Shi, D. Zhao, F. Zhao, C. Wang, G. Zamaratskaia, and C. Li, "Chicken-Eaters and Pork-Eaters Have Different Gut Microbiota and Tryptophan Metabolites," *Scientific Reports* 11 (2021): 11934.

50. Y. Wang, S. R. Lindemann, T.-W. L. Cross, M. Tang, C. M. Clark, and W. W. Campbell, "Effects of Adding Lean Red Meat to a U.S.-Style Healthy Vegetarian Dietary Pattern on Gut Microbiota and Cardiovascular Risk Factors in Young Adults: A Crossover Randomized Controlled Trial," *Journal of Nutrition* 153 (2023): 1439–1452.

51. J. Gao, X. Guo, W. Wei, et al., "The Association of Fried Meat Consumption With the Gut Microbiota and Fecal Metabolites and Its Impact on Glucose Homeostasis, Intestinal Endotoxin Levels, and Systemic Inflammation: A Randomized Controlled-Feeding Trial," *Diabetes Care* 44 (2021): 1970–1979.

52. K. Hoy, J. Clemens, and A. Moshfegh, "Estimated Protein Intake From Animal and Plant Foods by U.S. Adults, What We Eat in America, NHANES, 2015–2016," *Current Developments in Nutrition* 5 (2021): 133.

53. J. Foerster, G. Maskarinec, N. Reichardt, et al., "The Influence of Whole Grain Products and Red Meat on Intestinal Microbiota Composition in Normal Weight Adults: A Randomized Crossover Intervention Trial," *PLOS ONE* 9 (2014): 109606.

Supporting Information

Additional supporting information can be found online in the Supporting Information section.

Supporting Information file 1: mnfr70189-sup-0001-SuppMat.docx