

ANEXO I

Bibliografía adicional analizada para el desarrollo del estudio:


- *Habibisaravi R, Navaeinia S, Farnia S, Zarghami M. Alcohol, Cannabinoids, and Opioids Abuse and Dependence Among Psychiatric Inpatients. Iran J Psychiatry Behav Sci. 2015; 9(1):e229.*
- *Katz G, Durst R, Shufman E, Bar-Hamburger R, Grunhaus L. Substance abuse in hospitalized psychiatric patients. Isr Med Assoc J. 2008; 10(10):672-5.*
- *Ponizovsky AM, Rosca P, Haklai Z, Goldberger N. Trends in dual diagnosis of severe mental illness and substance use disorders, 1996-2010, Israel. Drug Alcohol Depend. 2015; 148:203-8.*
- *Guillem E, Pelissolo A, Vorspan F, Bouchez-Arbabzadeh S, Lépine J-P. Sociodemographic profiles, addictive and mental comorbidity in cannabis users in an outpatient specific setting. Encephale. 2009; 35(3):226-33.*
- *Weich L, Pienaar W. Occurrence of comorbid substance use disorders among acute psychiatric inpatients at Stikland Hospital in the Western Cape, South Africa. Afr J Psychiatry (Johannesbg). 2009; 12(3):213-7.*
- *Sinclair JMA, Latifi AH, Latifi AW. Co-morbid substance misuse in psychiatric patients: prevalence and association with length of inpatient stay. J Psychopharmacol (Oxford). 2008; 22(1):92-9.*
- *Ilgen MA, Hu KU, Moos RH, McKellar J. Continuing care after inpatient psychiatric treatment for patients with psychiatric and substance use disorders. Psychiatr Serv. 2008; 59(9):982-8.*
- *Błachut M, Badura-Brzoza K, Jarzab M, Gorczyca P, Hese RT. Dual diagnosis in psychoactive substance abusing or dependent persons. Psychiatr Pol. 2013; 47(2):335-52.*
- *Gual A. Dual diagnosis in Spain. Drug Alcohol Rev. 2007; 26(1):65-71.*
- *Lai HMX, Sitharthan T. Exploration of the comorbidity of cannabis use disorders and mental health disorders among inpatients presenting to all hospitals in New South Wales, Australia. Am J Drug Alcohol Abuse. 2012; 38(6):567-74.*
- *Kwon M, Yang S, Park K, Kim D-J. Factors that affect substance users' suicidal behavior: a view from the Addiction Severity Index in Korea. Ann Gen Psychiatry. 2013; 12(1):35.*
- *Landheim AS, Bakken K, Vaglum P. Impact of comorbid psychiatric disorders on the outcome of substance abusers: a six year prospective follow-up in two Norwegian counties. BMC Psychiatry. 2006; 6:44.*
- *DiClemente CC, Nidecker M, Bellack AS. Motivation and the stages of change among individuals with severe mental illness and substance abuse disorders. J Subst Abuse Treat. 2008; 34(1):25-35.*
- *Dixon L, McNary S, Lehman A. One-year follow-up of secondary versus primary mental disorder in persons with comorbid substance use disorders. Am J Psychiatry. 1997; 154(11):1610-2.*
- *Huang M-C, Yu C-H, Chen C-T, Chen C-C, Shen WW, Chen C-H. Prevalence and identification of alcohol use disorders among severe mental illness inpatients in Taiwan. Psychiatry Clin Neurosci. 2009; 63(1):94-100.*
- *Bonsack C, Camus D, Kaufmann N, Aubert AC, Besson J, Baumann P, et al. Prevalence of substance use in a Swiss psychiatric hospital: interview reports and urine screening. Addict Behav. 2006; 31(7):1252-8.*
- *Dixon L, McNary S, Lehman AF. Remission of substance use disorder among psychiatric inpatients with mental illness. Am J Psychiatry. 1998; 155(2):239-43.*

- *Caton CLM, Hasin DS, Shrout PE, Drake RE, Dominguez B, First MB, et al. Stability of early-phase primary psychotic disorders with concurrent substance use and substance-induced psychosis. Br J Psychiatry. 2007; 190:105-11.*
- *Xie H, McHugo GJ, Fox MB, Drake RE. Substance abuse relapse in a ten-year prospective follow-up of clients with mental and substance use disorders. Psychiatr Serv. 2005; 56(10):1282-7.*
- *Helseth V, Lykke-Enger T, Johnsen J, Waal H. Substance use disorders among psychotic patients admitted to inpatient psychiatric care. Nord J Psychiatry. 2009; 63(1):72-7.*
- *Trocchio S, Chassler D, Storbjörk J, Delucchi K, Witbrodt J, Lundgren L. The association between self-reported mental health status and alcohol and drug abstinence 5 years post-assessment for an addiction disorder in U.S. and Swedish samples. J Addict Dis. 2013; 32(2):180-93.*
- *Kessler RC. The epidemiology of dual diagnosis. Biol Psychiatry. 2004; 56(10):730-7.*


ANEXO II

Gráficas de los resultados obtenidos


VOLUNTARIEDAD




SEXO




SEGUIMIENTO




INGRESOS ANTERIORES EN LA UNIDAD




MOTIVO DE ALTA




NACIONALIDAD




PROCEDENCIA DE LOS EXTRANJEROS




CONSUMO DE TÓXICOS




CONVIVENCIA




CONSUMEN ALCOHOL




CONSUMEN PSICOESTIMULANTES




CONSUMEN THC




CONSUMEN BZD




PROBLEMAS JUDICIALES-LEGALES




PROBLEMAS ECONÓMICOS GRAVES




SITUACIÓN FAMILIAR




ANTECEDENTES PSIQ FAMILIARES



ESTADO CIVIL



¿TRASTORNO DUAL?



DIAGNÓSTICOS PRINCIPALES

