AAC Accepted Manuscript Posted Online 2 January 2019 Antimicrob. Agents Chemother. doi:10.1128/AAC.02373-18 Copyright © 2019 American Society for Microbiology. All Rights Reserved.

1 Screening The Pathogen Box for Identification of New Chemicals Agents With Anti

- 2 Fasciola hepatica Activity
- 3 Claudia Machicado^{a,b*}, Maria Pia Soto^{a‡}, Olga Timoteo^a, Abraham Vaisberg^a, Monica
- 4 Pajuelo^a, Pedro Ortiz^c, Luis A. Marcos^d
- 5 a Laboratorios de Investigación y Desarrollo, Facultad de Ciencias y Filosofía, Universidad
- 6 Peruana Cayetano Heredia, Lima, Peru.
- 7 b Institute for Biocomputation and Physics of Complex Systems, University of Zaragoza,
- 8 Spain.
- 9 c Laboratorio de Inmunología, Facultad de Ciencias Veterinarias, Universidad Nacional de
- 10 Cajamarca, Cajamarca, Peru.
- d Department of Medicine (Infectious Diseases), Stony Brook University, Stony Brook,
 New York, USA; Department of Microbiology and Molecular Genetics, Stony Brook
 University, Stony Brook, New York, USA; Global Health Institute, Stony Brook
 University, Stony Brook, New York, USA
- 15 d
- 16 * Corresponding author: Claudia Machicado, E-mail: <u>claudia.machicado.r@upch.pe</u>.
- 17 Laboratorios de Investigacion y Desarrollo, Facultad de Ciencias y Filosofía, Universidad
- 18 Peruana Cayetano Heredia. Av. Honorio Delgado 430, Lima 31, Peru.
- 19 ‡ Not longer at the Universidad Peruana Cayetano Heredia.
- 20 Running title: Screening of compounds against Fasciola hepatica

Downloaded from http://aac.asm.org/ on January 3, 2019 by guest

1

21 Keywords: Fasciola hepatica, triclabendazole, fasciocidal activity, in vitro screening.

22

23 Abstract

24 Fascioliasis is an infectious parasitic disease distributed globally and caused by the liver 25 flukes Fasciola hepatica or F. gigantica. This neglected tropical disease affects both animals and humans and it represents a latent public health problem due to the significant 26 economic losses related to animal husbandry. For decades, Triclabendazole has been the 27 unique anti-Fasciola drug that can effectively treat this disease. However, triclabendazole 28 29 resistance in Fascioliasis has been more recently reported around the world, and thus the 30 discovery of novel drugs is an urgent need. The aim of this study was to investigate the 31 fasciocidal properties of 400 compounds contained in the Pathogen Box. The first stage of 32 the screening was carried out by measuring the fasciocidal activity on metacercariae at a 33 concentration of 33 μ M of each compound (standard dose). Subsequently the 50% inhibitory concentration (IC₅₀) values of the most active compounds (n=33) were assayed 34 35 on metacercariae and resulted in 13 compounds with $IC_{50} \leq 10~\mu M.$ The second stage queried these compounds at 33 µM on adult flukes where seven showed high mortality 36 rates > 50%. Four hit compounds were selected based on predicted nontoxic properties and 37 IC_{50} values obtained on adult worms resulted < 10 μ M thus representing the best 38 39 fasciocidal compounds tested here. Cytotoxicity assay on four types of cell lines 40 demonstrated that three compounds are nontoxic at its most active concentration. In conclusion, three hit compounds identified in this proof-of-concept study are potential 41 42 candidates in the discovery of new fasciocidal drugs. Further studies are warranted.

43 INTRODUCTION

44 Fasciola hepatica is the etiological agent of fascioliasis, the most widespread trematodiasis 45 that affects both humans and herbivorous mammals such as sheep, cattle, goats and other 46 species (1, 2). In humans, fascioliasis can be acquired by the consumption of contaminated 47 vegetables. Up to 17 million people in 51 countries are estimated to be infected with F. 48 hepatica worldwide and more than 91 million are at risk of infection by this parasite (3, 4). 49 Among all continents, the Andean Region of South America is the most affected by 50 Fasciola where prevalence rates above 10% have been documented (5-8) and national 51 treatment programs are being scaled up.

Triclabendazole (TCBZ) is the most single effective fasciocidal drug, with activity against 52 53 both the infective larvae (Metacercaria) and adult worms, and an efficacy that exceeds 90% 54 in humans after a single oral dose (9, 10). Nonetheless, after decades of successful efficacy, 55 TCBZ resistance has developed in both animals and humans (11). Cases of TCBZ-resistant 56 Fasciola in both animals and humans have been reported in Australia, Europe and Latin 57 America (12-18). The development of TCBZ resistance represents an important public 58 health concern throughout the world that mainly affects animal husbandry and leads to 59 enormous economic losses (19). As a consequence, the discovery of novel effective drugs 60 and vaccines against Fasciola is an urgent need for the global control of fascioliasis. 61 Repurposing of praziquantel (PZQ) as anti-Fasciola drug failed whereas oxfebendazole 62 showed to be an effective drug in animals (20, 21). Currently, there is no other fasciocidal 63 drug in clinical practice for humans, and thus TCBZ remains the unique treatment against 64 this infectious disease.

Open-access drug discovery provides a substantial resource in the research of those

diseases that affect primarily people living in low-resources locations. The Medicines for 66 Malaria Venture (MMV) foundation assembled a set of compounds, called "Malaria Box", 67 which has been tested against various infectious agents including Cryptosporidium parvum 68 69 (22), Plasmodium falciparum (23, 24) Schistosoma mansoni (25, 26), Toxoplasma gondii 70 (27), and mycobacteria (28, 29). Later, a new set of chemical entities was assembled and 71 named the Pathogen Box collection. It contains 400 drug-like compounds that have showed 72 inhibitory activity on various infectious diseases such as haemonchosis, toxoplasmosis, 73 tuberculosis, neosporosis, malaria, sleeping sickness, Chagas, leishmaniasis and trypanosomiasis (30-36). The Pathogen Box has been tested also in fungal diseases caused 74 75 by Cryptococcus neoformans and Candida albicans (37-39). The aim of this study was to identify the fasciocidal activity of 400 compounds contained in the Pathogen Box by in 76 77 vitro testing.

78

65

79 **MATERIALS AND METHODS**

80 Study design. The study was conducted in three stages: (i) bioassays on metacercariae; (ii) 81 bioassays on adult worms; and (iii) cytotoxicity on cell (Figure 1). The best ffasciocidal 82 compounds were selected in each stage to be tested in the next phase based on in vitro 83 biological activity. To complement our knowledge on the active compounds obtained by 84 the experimental assays, computational resources were consulted to describe the chemical 85 properties as well as the in silico toxicology features and biological targets from these 86 active compounds.

88 Drugs and Media

87

89 The Pathogen Box was provided by the MMV agency (Geneva, Switzerland) and 90 manufactured by Evotec (USA). The 400 drug-like molecules were supplied in 96-well 91 plates as stock solutions of 10 mM dissolved in dimethyl sulfoxide (DMSO). Full data of 92 The Pathogen Box compounds is available at https://www.pathogenbox.org (40). TCBZ 93 was purchased from Sigma-Aldrich (Buchs, Switzerland). All of the compounds of The 94 Pathogen Box were dissolved in DMSO (Sigma-Aldrich, Irvine, UK) for drug stock 95 solutions of 200 µM. Additional vials of MMV063404, MMV003270, MMV085210, 96 MMV676380, MMV687730, MMV687251, MMV1030799, MMV690102, MMV1029203, 97 MMV676053, MMV688179, MMV023969 and MMV688921 were manufactured by 98 Evotec (France). RPMI 1640 culture medium (Sigma- Aldrich, St. Louis, US) was used for 99 both stages, metacercariae and adult worm; supplemented with penicillin (100 U/ml) and 100 streptomycin (100 µg/ml) (Sigma-Aldrich, St. Louis, US).

101 Parasites

Metacercariae of *F. hepatica* were obtained following the protocol described by Ortiz et al. (16), at the Immunology and Research Laboratory of the Faculty of Veterinary Sciences of the *Universidad Nacional de Cajamarca* in Peru. Eggs of *F. hepatica* were collected directly from the gallbladder of sheep slaughtered in a popular abattoir in the city of Cajamarca (TCBZ-resistant endemic area for fascioliasis). Miracidia were from Fasciola eggs after incubation for 15 days at 25°C. Afterwards, they were used to infect *Lymnaea sp.* snails (5-6 mm) in a proportion of two miracidia per snail. The infected snails were kept in

plastic containers for 45 to 60 days at room temperature. After this time, the snails were

109

110 stimulated by direct solar exposure and water at 4-8°C to produce metacercaria. Approximately 20,000 metacercariae were obtained for this study and stored in cryovials 111 112 on distilled water at 4-8°C. Adult worms were collected from bile ducts of infected cattle from a slaughterhouse in Lima, Peru; and maintained at 37°C until usage (within 2h). 113 Before its incubation, three washes with phosphate-buffered saline (PBS) (HiMedia, India) 114 115 and one additional with supplemented RPMI were performed to remove host debris. All the incubations, for both metacercariae and adults, were carried out at 37°C with 5% CO₂. 116

117 In vitro screening on metacercariae

118 The 400 compounds were initially tested at 33 µM on F. hepatica metacercariae. Drug 119 stock solutions were diluted in 96-well plates (BD Falcon, US) with RPMI 1640 120 supplemented with antibiotic up to a final volume of 180 µL. In all in vitro assays, positive 121 and negative controls were run in parallel for each assay batch. A range between seven and 122 ten metacercariae were added to each well, previously analyzed microscopically to confirm 123 its viability (microscopic features intact). The metacercariae viability considered some 124 physical properties of the parasite determined by microscopy as described previously (41, 125 42). The MCs viability was surveyed as a function of both damage in membrane and fluke 126 colour (translucence). Therefore, a low viability corresponded to big damage and high 127 translucency. The viability scale was scored as follows: +++, total damage (dead parasite, 128 shattered membrane and mostly translucent); ++, partial damage (partial membrane damage 129 and highly translucent); +, mild damage (partial membrane damage, poorly translucent) and 130 no damage (intact membrane, dark metacercariae, lack of translucency).

131 Positive-control wells contained TCBZ 10 µM whereas F. hepatica metacercariae incubated in the presence of the highest concentration of DMSO served as negative control. 132 Each test was performed in triplicate. Culture plates were incubated at 37°C in a humidified 133 5% CO₂ atmosphere for 72 h. First, metacercariae were evaluated by inverted microscopy 134 (PhotoZoom, Cambridge Instruments) at magnification 10X and 20X at 24, 48 and 72 h 135 136 post drug exposure to determine its viability. Only the compounds that caused, on average, at least 25% of metacercariae mortality at 72 hours were considered for IC_{50} (50%) 137 138 inhibitory concentration) determination. Experiments were run in sets of triplicates. The 139 mean mortality percent of the study compounds were compared to that of DMSO. A 140 standard deviation (SD) was also estimated.

141 In the second part, we determined the IC_{50} of the selected compounds chosen in the 142 previous bioassay. Drugs were tested at concentrations of 2.1, 4.2, 8.4 and 33 μ M using 143 supplemented culture medium. The incubation was done under the conditions described 144 above, by triplicate and considering TCBZ and DMSO as controls. Anti-parasite activity 145 was evaluated at 24, 48 and 72 h post exposure, using the above-mentioned metacercariae 146 viability scale. Viability (mean % of viable parasites) at 72h was considered for the 147 estimation of IC₅₀. IC₅₀ values of test compounds were determined by linear regression 148 analysis using CompuSyn software (Version 3.0.1, 2007; ComboSyn Inc., USA). The linear 149 correlation coefficient (r) was obtained.

150 Assessment of anti-Fasciola activity in vitro on adult worms

151 Those compounds that showed activity $IC_{50} \le 10 \ \mu M$ on metacercaria were subsequently 152 tested in adult stage of *F. hepatica*. In all *in vitro* assays, positive and negative controls 153 were run in parallel for each assay batch. First, the selected compounds were tested at 33

154	μM by triplicate, using drug stock solutions diluted in supplemented RPMI on 6-well plate
155	up to a final volume of 4 ml. Adult worms were thoroughly washed with PBS to remove
156	host debris and then three worms were placed in each well. The incubation was done under
157	the same conditions as those applied in bioassays with metacercariae. Positive control
158	consisted of 50 μM TCBZ and the negative control was DMSO at the highest
159	concentration. The viability of adult flukes was scored after 24 and 48 h using a motility
160	criterion previously described (43) and also color and rigidity criteria previously applied by
161	our team (data not published). Motility was assessed only in adults and not in MC because
162	this latter has no movements. Rigidity was a parameter used to confirm the damage caused
163	by the drug once the incubation time finished. In general, low motility level corresponded
164	to transparent and rigid worms. Those changes were attributed to the damage caused by a
165	drug. The viability scale was determined as follows: (i) worm motility: 3, normal
166	movements; 2, reduced movements; 1, very weak movements and 0, absence of movements
167	(i.e. death of worm); (ii) worm color: +++, dark red; ++, pink; +, slightly transparent and -,
168	totally transparent; and (iii) worm rigidity: -, no rigidity; +, rigidity and ++, cell break when
169	touched. Assessments at 72h post drug exposure were not done because death of worms
170	always occurred \leq 48h. Experiments were run in sets of triplicates. The mean mortality
171	percent and SD of the study compounds were estimated. The selected compounds were
172	those that caused an average mortality $> 50\%$ in adult parasites. Then IC_{50} assays were
173	conducted by testing the selected compounds at five different concentrations 0.31, 0.93,
174	2.78, 8.33 and 25.0 $\mu\text{g/ml}.$ DMSO and TBZ were used as negative and positive controls,
175	respectively. Parasite viability at 24 h were estimated based on survival in DMSO. The
176	IC50s and 95% CI were estimated using GraphPad Prism 7.0 software using a variable
177	slope of the sigmoidal curve from normalized percent activity values and log10-

AAC

transformed concentrations. Top and bottom values were constrained to 100 and 0,
respectively. The fasciocidal activity was determined by considering the adult viability
scale described before.

181 Computational analysis.

182 Evaluation of biological targets of small compounds. To learn about biological targets, 183 those compounds that showed promising anti-fasciola activity in the adult stage as well as 184 TCBZ were entered in the ChEMBL database (https://www.ebi.ac.uk/chembl/) (44). First, 185 the SMILES (Simplified Molecular-Input Line-Entry System) of each selected compound 186 were obtained from the supplementary material provided by the MMV (also available at 187 www.mmv.org). Then the SMILES were entered in ChEMBL and known targets of each 188 compound were retrieved. ChEMBL compares the query compound to a large database of 189 compounds and their targets available from multiple sources including the projects funded 190 by MMV (45). The target name, organism and protein target classification were collected.

In silico cytotoxicity prediction. Lazar (lazy structure–activity relationships), a modular framework for predictive toxicology, was consulted to predict the toxic effects of the selected compounds that showed activity on metacercariae (46-48). Lazar was accessed through <u>https://lazar.in-silico.de/predict</u> and SMILES of each compound were entered. Relevant data including carcinogenicity in rodents, mutagenicity in *Salmonella typhi* and acute toxicity on *Fathead minnow*, Blood Brain Barrier Penetration and the maximum recommended daily dose in humans were predicted.

198 Cell Growth Inhibition Bioassay.

Cytotoxicity of the compounds was evaluated in tumor and non-tumor cell lines using the
sulforhodamine B (SRB) assay method (49, 50). Cell lines tested include BALB/3T3 (Nontumorogenic, BALB/c mouse embryo cells), H460 (human lung large cell carcinoma),
DU145 (human prostate carcinoma) and HT-29 (human colon adenocarcinoma).

203 To determine the cytotoxicity of the compounds, cells were plated into 96-well tissue 204 culture plates and in their corresponding growth medium Dulbecco's Modified Eagle 205 Medium (DMEM) at approximately 10% confluency (BALB/3T3 at 3,500 cells/well, H460 206 at 1,500 cells/well, DU145 at 3,500 cells/well and HT-29 at 3,000 cells/well) and incubated 207 at 37°C in a 5% CO₂ and 95% air humidified atmosphere for 24 h to allow cells to attach. A 208 plate containing each of these cells was fixed in situ with trichloroacetic acid (TCA) in 209 order to obtain the cell values at zero time before adding the compounds. The rest of the 210 plates containing the different cell lines received serial dilutions of the compound to be 211 tested at the following final concentrations: 4, 1, 0.25 and 0.0625 μ g/mL. The plates were 212 then incubated at 37°C in a 5% CO₂ and 95 % air humidified atmosphere for 48 h. The 213 assay was terminated by the addition of cold TCA. TCA treated plates were incubated at 214 4°C for 1 hour and then washed five times with tap water to remove TCA and air dried. 215 Background optical densities were measured in wells incubated with growth medium 216 without cells. TCA-fixed cells were stained for 20 minutes with 0.4% (w/v) SRB dissolved 217 in 1% acetic acid. At the end of the staining period unbound dye was removed by washing 218 four times with 1% acetic acid. After air drying the plates, bound dye was solubilized with 219 10 mM Tris base (pH 10.5) and the absorbance read on an automated plate reader at a 220 wavelength of 550 nm. The GI₅₀ value was defined as the concentration of test sample resulting in a 50% reduction of absorbance as compared with untreated controls that 221

222 received a serial dilution of the solvent in which the test samples were dissolved and was 223 determined by linear regression analysis. The optical density values obtained were used to 224 determine the cell growth and cytotoxicity from each compound.

225 Ethics. This study was approved by the Animal Ethics Committee of the Universidad 226 Peruana Cayetano Heredia (Approval ID Code 41-07-16).

227 RESULTS

In vitro activity of The Pathogen Box determined on F. hepatica metacercariae 228

229 In the first stage of the study, the 400 compounds contained in the Pathogen Box were in 230 vitro screened against F. hepatica metacercaria. A total of 33 compounds showed mean 231 mortality rates above 25% at 33 µM but all these resulted being less active than TCBZ 232 (mortality rate of 90%) as shown on Table 1. Fasciocidal activity of these 33 compounds 233 was then assessed by determining the IC₅₀ values (Table 1). As a result, 13 compounds 234 showed potent inhibitory activities with IC_{50} values between 0.31 μ M and 8.23 μ M and 235 were then assayed in adult worms although its low r values (Table 1).

236 In vitro activity of selected compounds on F. hepatica adult worms and in silico toxicology 237 prediction.

238 The 13 selected compounds listed on Table 2 were assayed at 33 µM in adult worms. Seven 239 compounds produced moderate or high mean mortality rates (> 50%) (Table 2). These 240 were MMV003270, MMV676380, MMV690102, MMV1029203, MMV063404, 241 MMV1030799, and MMV688921. Six compounds showed low mortality rates (<50%) and for that reason these were not considered in the next assays. Before to proceed with the IC_{50} 242 243 assay, in silico safety profiles of the seven selected compounds were predicted by lazar 244 program (Table 1). Whereas MMV003270 and MMV676380 were predicted non-245 carcinogenic and non-tumorigenic compounds, MMV690102 was deemed noncarcinogenic and tumorigenic (Table 1). MMV1029203, MMV063404, MMV1030799, and 246 247 MMV688921 were predicted carcinogenic and tumorigenic substances. Thus, the three 248 deemed non-carcinogenic compounds as well as MMV1029203, a predicted carcinogenic 249 substance that had the highest mean mortality rate (78%), were tested in adult worms. Such 250 four compounds constitute our hit compounds.

251 To determine which of the four hit compounds were most potent at inhibiting the growth of 252 F. hepatica adult worms, the IC₅₀ values were determined. The hit compounds had IC₅₀ 253 values $< 10 \ \mu$ M in adult worms (Table 3, Fig. S1, Table S1). These four hit compounds 254 were tested in the cytotoxicity study on cell cultures.

255 In vitro cytotoxicity on cell lines.

256 Cytotoxicity of the four hit compounds against cell lines was evaluated in culture (Table 3). The GI₅₀ values ranged from 0.95 and >23.73 μ M across the four types of cell lines assayed 257 258 (Table 3). MMV003270, MMV676380, MMV1029203 and TCBZ presented GI_{50} values 259 above its IC₅₀ values thus meaning that these compounds are not toxic at their active 260 concentrations. In one of the four cell lines, MMV690102 had a GI_{50} value below its IC_{50} 261 value thus suggesting that it may cause a level of toxicity in certain cell types at its active 262 concentration (Table 3).

²⁶³ Computational recognition of targets.

²⁶⁴ As a result of the search in the ChEMBL database, a total of 27 targets were recognized for 265 TCBZ whereas MMV003270 resulted to have 19 known target, most of them in humans

266 (Table 4). MMV003270 and TCBZ have common human targets that comprise Nuclear 267 factor erythroid 2-related factor 2, Microtubule-associated protein tau and TAR DNA-268 binding protein 43. According to the data deposited in ChEMBL, MMV003270 targets a 269 number of cytochrome p450 of family 1, 2 and 3. MMV676380 and MMV023969 have 270 identical cell targets that include human glucose transporter and hexose transporter of 271 Plasmodium falciparum and Leishmania mexicana (Table 4). Targets for MMV1029203 272 and MMV676053 also resulted to have known targets including human ferrochelatase and Inosine-5'-monophosphate dehydrogenase of Cryptosporidium parvum, respectively. The 273 274 remaining eight compounds had no known targets according to the ChEMBL database 275 (Table 4).

276

277 DISCUSSION

278 In the present study, the Pathogen Box was queried to identify compounds with in vitro 279 anti-Fasciola activity against both metacercariae and adult worms (Figure 1). We found 13 280 compounds with potent inhibitory activity on metacercariae (IC₅₀ < 10 μ M), meaning that 281 3% of the substances within the Pathogen Box are effective against the infective form of F. 282 hepatica. Two out of the 13 compounds (MMV687730 and MMV687251) had the most 283 potent activity against metacercariae with IC₅₀ values below 1 µM but showed mild effects 284 on adult worms (Tables 1 and 2). Since we were interested in identifying hit compounds 285 that were active on larvae and adult stages, these two compounds were not further studied 286 (Table 2). When assayed on adult worms, seven promising compounds showed mortality 287 rates above 50% (Table 2). As a criterion for hit prioritization during the screening on adult 288 worms, we prepared a list of hit compounds that mostly excluded the predicted

289 carcinogenic/tumorigenic compounds (Table 3). Thus, three (MMV676380, MMV003270 290 and MMV690102) of the seven most promising candidates were included in the list of hit 291 compounds since they were predicted noncarcinogenic agents (Table 1). One additional 292 compound (MMV1029203) that was predicted as a carcinogenic compound was also 293 included due to its very high effect on adult worms. According to our results, the four hit 294 compounds resulted potent inhibitory molecules both on MC and adult stages (Table 3). 295 The cytotoxicity assay revealed that three hit compounds (MMV676380, MMV003270 and 296 MMV1029203) were non-toxic agents at its most active concentrations when assayed on 297 cell lines (Table 3). In contrast, MMV690102 may cause cell cytotoxicity at its most active 298 concentration meaning that it is not a primary candidate for drug development (Table 3). 299 Our results are consistent with previous cytotoxicity assays on HepG2, HL60 and MRC5 as 300 shown on Table 3 (data provided by the MMV as part of the supporting information for the 301 Open Access Malaria Box).

302 Repurposing of hits, using the Pathogen Box, against F. hepatica is highly relevant since 303 TCBZ is the only existing effective drug for which resistance is known (51-53). Previous 304 works tried to repurpose albendazole, nitroxynil and closantel as candidate fasciocidal 305 drugs but treatment failed (54, 55). In the present study, four out of 400 compounds 306 contained in The Pathogen Box showed potent inhibitory activity against the infective form 307 of F. hepatica as well as its adult form (Table 3). Such finding represents a relevant 308 contribution in the identification of dual drug candidates that are able to act against the 309 initial stages of the infective larvae (metacercaria) and adult forms of liver flukes, similar to 310 TCBZ. Additionally, other 13- compounds showed biological activity at $< 20 \mu$ M against 311 metacercaria (Table 1). Since MC represents the initial infective form of parasites, it should

14

312 be primarily controlled through potent compounds such as those identified here (Table 1). 313 Future exploration of The Pathogen Box in newly juvenile metacercaria is desirable given 314 that some compounds may have not penetrated the cyst wall of larvae. By testing compounds on juvenile worms, some additional molecules might be recognized that are 315 316 active in adult worms.

317

318 The four hit compounds identified in this study have been previously characterized against 319 Plasmodium falciparum, Ancylostoma ceylanicum, Trypanosoma cruzi and Leishmania 320 donovani (data provided by the MMV as part of the supporting information for the Open 321 Access Malaria Box). Therefore, a common mechanism of action or target is plausible 322 among the hit compounds across such pathogens. For instance, MMV676380 has 323 previously shown to have a lethal effect on P. falciparum and here was found to be a potent 324 inhibitory compound against F. hepatica (36, 56). Known targets of MMV676380 are the 325 glucose and hexoses transporters suggesting that such mechanism may be affected in both 326 parasites in presence of such compound (Table 4). In the other hand, MMV003270 327 (Zoxazolamine), that is also active against A. ceylanicum, resulted to have 19 targets 328 including three human proteins that are also targeted by TCBZ (Table 4). Two of these 329 proteins are transcription regulators (Nuclear factor erythroid 2-related factor 2 and TAR 330 DNA-binding protein 43) whose disruption may affect the gene expression. Such finding is 331 in accordance with a hypothetical mechanism of action of TCBZ that involves a direct 332 effect of the drug on protein synthesis (11, 57). Similarly, the microtubule-associated protein tau is a known target both of TCBZ and MMV003270. TCBZ is a benzimidazole-333 334 derivative that disrupts the assembly of microtubules in helminths by binding to tubulin

335

Downloaded from http://aac.asm.org/ on January 3, 2019 by guest

336 formation mechanism. Common targets of TCBZ and MMV003270 may be partially 337 explained by the similar scaffold structures. MMV1029203, one of the four hit compounds, 338 targets a human ferrochelatase that is a mitochondrial factor involved in protoheme 339 biosynthesis. This latter is a vital process that exists also in F. hepatica and whose 340 disruption may be lethal. Some known targets of the hit compounds here identified 341 correspond to human proteins which suggests that a level of toxicity may exist in humans. 342 However according to our results with cell lines, the compound concentrations needed to 343 kill F. hepatica (IC_{50}) are considerably less than that to cause cell death (GI_{50}) which means 344 that these are nontoxic (Table 3), except for MMV690102. Although no F. hepatica target 345 is recognized for our hit compounds, the demonstration of inhibitory activity of such 346 chemical agents both in metacercariae and adult forms suggests that common targets may 347 exist in both liver fluke stages. The identification of drug targets becomes an important step 348 that drive the discovery of novel antiparasitic agents administered by various ways (34). 349 For that reason, further studies to identify potential F. hepatica targets of hit compounds are 350 desirable. Such a study should consider the recognition of human homologs in F. hepatica 351 according to our results (Table 4).

molecules (58). Our results suggest that MMV003270 also affects the microtubules

Our study has some limitations. First, TCBZ metabolites (TCBZ-sulfoxide and TCBZsulfone) that are quickly released *in vivo* were not included in this pilot study. However given that TCBZ has a moderate *in vivo* and *in vitro* fasciocidal effects it is suitable as positive control in bioassays (59, 60). A second limitation is that alive *F. hepatica* worms were collected from a local abattoir where some animals may have been infected by various other pathogens or may have been treated with TCBZ. To guarantee the best quality of

Chemotherapy

358 adult worms for bioassays, we performed a quality control on adult fasciolas before using 359 these in the experiments. Thus, only worms that presented intense brown or red color and 360 that have active motility were selected. All the remaining were discarded. A third limitation 361 is the low number of parasites used for the assays, that did not allow performing formal 362 statistical comparisons of activity between TCBZ and test drugs. Obtaining MC and adults 363 was a challenging task since both MC and adult worms were collected from natural 364 reservoirs. Therefore we had limited access to parasites for bioassays. However, our exploratory study aimed to identify fasciocidal compounds, we found that negative controls 365 366 were enough for such purposes.

367 In conclusion, we identified three promising non-cytotoxic drug-like compounds, 368 MMV003270, MMV676380 and MMV1029203, that showed a potent biological activity 369 against *F. hepatica* metacercaria and adult worms. Such compounds represent new lead 370 candidates to potentially become future anti-*F. hepatica* drugs. By acting both on infective 371 form and adult worms, such agents may provide an appropriate treatment against 372 fascioliasis.

373

374

375

376 ACKNOWLEDGMENTS

We thank the Medicines Malaria Ventures foundation for having funded this study and
provided the Pathogen Box. We acknowledge Dr. Jennifer Keiser for her invaluable advice
regarding the preparation of bioassays.

380 **Disclosures:** The authors state they have not conflict of interest to declare.

381

382 **REFERENCES**

383 1. Dalton JP. 1999. Fasciolosis. United Kingdom. Ed. Cab International.

- Marcos LA, Yi P, Machicado A, Andrade R, Samalvides F, Sánchez J, Terashima
 A. 2007. Hepatic fibrosis and Fasciola hepatica infection in cattle. J Helminthol
 81(4):381-6.
- 387 3. Keiser J, Utzinger J. 2009. Food-borne trematodiases. Clin Microbiol Rev
 388 22(3):466-83.
- 4. Fürst T, Keiser J, Utzinger J. 2012. Global burden of human food-borne
 trematodiasis: a systematic review and meta-analysis. Lancet Infect Dis 12(3):21021.
- Mas-Coma, S.; Bargues, MD.; Valero, MA. 2005. Fascioliasis and other plant-borne
 trematode zoonoses. Int J Parasitol 35(11-12):1255-78.
- World Health Organization. 2007. Fact sheet on fascioliasis. Action against worms.
 pp. 1–8. Geneva, WHO.
- Espinoza J, Maco V, Marcos L, Saez S, Neyra V, Terashima A, Samalvides F,
 Gotuzzo E, Chavarry E, Huaman MC, Bargues MD, Valero A, Mas-Coma S. 2007.
 Evaluation of fas2-elisa for the serological detection of Fasciola hepatica infection
 in humans. Am J Trop Med Hyg 76(5).

400	8.	Carmona C, Tort JF. 2017. Fasciolosis in South America: epidemiology and control
401		challenges. J Helminthol 91(2):99-109.
402	9.	Bennett J, Köhler P. 1987. Fasciola hepatica: action in vitro of triclabendazole on
403		immature and adult stage. Exp Parasitol 63(1):49-57.
404	10	. Apt W, Aguilera X, Vega F Miranda C, Zulantay I, Perez C, Gabor M, Apt P. 1995.
405		Treatment of human chronic fascioliasis with triclabendazole: drug efficacy and
406		serologic response. Am J Trop Med Hyg 52:532-535.
407	11	. Fairweather I. 2005. Triclabendazole: new skills to unravel an old(ish) enigma. J
408		Helminthol 79(3):227-34.
409	12	. Overend DJ, Bowen FL. 1995. Resistance of Fasciola hepatica to triclabendazole.
410		Aus Vet J 72:275-276.
411	13	. Mitchell GB, Maris L, Bonniwel MA. 1998. Triclabendazole-resistant liver fluke in
412		Scottish sheep. Vet Rec 143:399.
413	14	. Moll L, Gaasenbeek CP, Vellema P, Borgsteede FH. 2000. Resistance of Fasciola
414		hepatica against triclabendazole in cattle and sheep in the Netherlands. Vet Parasitol
415		91:153-158.
416	15	. Oliveira DR, Ferreira DM, Stival CC, Romero F, Cavagnolli F, Kloss A, Araújo
417		FB, Molento MB. 2008. Triclabendazole resistance involving Fasciola hepatica in
418		sheep and goats during an outbreak in Almirante Tamandare, Paraná, Brazil. Rev

419 Bras Parasitol Vet 17, 149–153.

420	16. Ortiz P, Scarcella S, Cerna C, Rosales C, Cabrera M, Guzman M, Lamenza P,
421	Solana H. 2013. Resistance of Fasciola hepática againts Triclabendazole in cattle in
422	Cajamarca (Peru): A clinical trial and an in vivo efficacy test in sheep. Vet Parasitol
423	195:118-121.
40.4	

- 424 17. Winkelhagen AJ, Mank T, de Vries PJ, Soetekouw R. 2012. Apparent
 425 triclabendazole-resistant human Fasciola hepatica infection, the Netherlands. Emerg
 426 Infect Dis 18(6):1028-9.
- 427 18. Cabada MM, Lopez M, Cruz M, Delgado JR, Hill V, White AC Jr. 2016. Treatment
 428 Failure after Multiple Courses of Triclabendazole among Patients with Fascioliasis
 429 in Cusco, Peru: A Case Series. PLoS Negl Trop Dis 25;10(1):e0004361.
- 430 19. Bekele M, Tesfay, H, Getachew Y. 2010. Bovine Fasciolosis: Prevalence and Its
 431 Economic Loss Due to Liver Condemnation at Adwa Municipal Abattoir. Ejast
 432 North Ethiopia, 39-47.
- 433 20. Michael D, Isaac-Renton J. 1992. Praziquantel failure in the treatment of *Fasciola*434 *hepatica*. Can J Infect Dis 3(1):33-36.
- 435 21. Gomez-Puerta LA, Gavidia C, Lopez-Urbina MT, Garcia HH, Gonzalez AE;
 436 Cysticercosis Working Group in Peru. 2012. Efficacy of a single oral dose of
 437 oxfendazole against Fasciola hepatica in naturally infected sheep. Am J Trop Med
 438 Hyg 86(3):486-8.
- 439 22. Bessoff K, Spangenberg T, Foderaro JE, Jumani RS, Ward GE, Huston CD. 2014.
 440 Identification of Cryptosporidium parvum active chemical series by Repurposing
 441 the open access malaria box. Antimicrob Agents Chemother 58(5):2731-9.

	Scammells PJ, McGowan S. 2015. Screening the Medicines for Malaria Venture
	"Malaria Box" against the Plasmodium falciparum aminopeptidases, M1, M17 and
	M18. PLoS One 10(2):e0115859.
24.	Bowman JD, Merino EF, Brooks CF, Striepen B, Carlier PR, Cassera MB. 2014.
	Antiapicoplast and gametocytocidal screening to identify the mechanisms of action
	of compounds within the malaria box. Antimicrob Agents Chemother 58(2):811-9.
25.	Pasche V, Laleu B, Keiser J. 2018. Screening a repurposing library, the Medicines
	for Malaria Venture Stasis Box, against Schistosoma mansoni. Parasit Vectors
	11(1):298.
26.	Ingram-Sieber K, Cowan N, Panic G, Vargas M, Mansour NR, Bickle QD, Wells
	TN, Spangenberg T, Keiser J. 2014. Orally active antischistosomal early leads
	identified from the open access malaria box. PLoS Negl Trop Dis 8(1):e2610.
27.	Boyom FF, Fokou PV, Tchokouaha LR, Spangenberg T, Mfopa AN, Kouipou RM,
	Mbouna CJ, Donfack VF, Zollo PH. 2013. Repurposing the open access malaria
	box to discover potent inhibitors of Toxoplasma gondii and Entamoeba histolytica.
	Antimicrob Agents Chemother 58(10):5848-54.
28.	Van Voorhis WC, Adams JH, Adelfio R, Ahyong V, Akabas MH, Alano P, Alday
	A, Alemán Resto Y, Alsibaee A, Alzualde A, Andrews KT, Avery SV, Avery VM,
	Ayong L, Baker M, Baker S, Ben Mamoun C, Bhatia S, Bickle Q, Bounaadja L,
	Bowling T, Bosch J, Boucher LE, Boyom FF, Brea J, Brennan M, Burton A,

23. Paiardini A, Bamert RS, Kannan-Sivaraman K, Drinkwater N, Mistry SN,

Caffrey CR, Camarda G, Carrasquilla M, Carter D, Belen Cassera M, Chih-Chien

Accepted Manuscript Posted Online

442

443

444

445

446

447

448

449

450

451

452

453

454

455

456

457

458

459

460

461

462

463

Antimicrobial Agents and Chemotherapy

464	Cheng K, Chindaudomsate W, Chubb A, Colon BL, Colón-López DD, Corbett Y,
465	Crowther GJ, Cowan N, D'Alessandro S, Le Dang N, Delves M, DeRisi JL, Du AY,
466	Duffy S, Abd El-Salam El-Sayed S, Ferdig MT, Fernández Robledo J, Fidock DA,
467	Florent I, Fokou PV, Galstian A, Gamo FJ, Gokool S, Gold B, Golub T, Goldgof
468	GM, Guha R, Guiguemde WA, Gural N, Guy RK, Hansen MA, Hanson KK,
469	Hemphill A, Hooft van Huijsduijnen R, Horii T, Horrocks P, Hughes TB, Huston C,
470	Igarashi I, Ingram-Sieber K, Itoe MA, Jadhav A, Naranuntarat Jensen A, Jensen LT,
471	Jiang RH, Kaiser A, Keiser J, Ketas T, Kicka S, Kim S, Kirk K, Kumar VP, Kyle
472	DE, Lafuente MJ, Landfear S, Lee N, Lee S, Lehane AM, Li F, Little D, Liu L,
473	Llinás M, Loza MI, Lubar A, Lucantoni L, Lucet I, Maes L, Mancama D, Mansour
474	NR, March S, McGowan S, Medina Vera I, Meister S, Mercer L, Mestres J, Mfopa
475	AN, Misra RN, Moon S, Moore JP, Morais Rodrigues da Costa F, Müller J,
476	Muriana A, Nakazawa Hewitt S, Nare B, Nathan C, Narraidoo N, Nawaratna S, Ojo
477	KK, Ortiz D, Panic G, Papadatos G, Parapini S, Patra K, Pham N, Prats S, Plouffe
478	DM, Poulsen SA, Pradhan A, Quevedo C, Quinn RJ, Rice CA, Abdo Rizk M,
479	Ruecker A, St Onge R, Salgado Ferreira R, Samra J, Robinett NG, Schlecht U,
480	Schmitt M, Silva Villela F, Silvestrini F, Sinden R, Smith DA, Soldati T,
481	Spitzmüller A, Stamm SM, Sullivan DJ, Sullivan W, Suresh S, Suzuki BM, Suzuki
482	Y, Swamidass SJ, Taramelli D, Tchokouaha LR, Theron A, Thomas D, Tonissen
483	KF, Townson S, Tripathi AK, Trofimov V, Udenze KO, Ullah I, Vallieres C, Vigil
484	E, Vinetz JM, Voong Vinh P, Vu H, Watanabe NA, Weatherby K, White PM,
485	Wilks AF, Winzeler EA, Wojcik E, Wree M, Wu W, Yokoyama N, Zollo PH, Abla
486	N, Blasco B, Burrows J, Laleu B, Leroy D, Spangenberg T, Wells T, Willis PA.

487 2016. Open Source Drug Discovery with the Malaria Box Compound Collection for 488 Neglected Diseases and Beyond. PLoS Pathog 12(7):e1005763.

489 29. Low JL, Wu ML, Aziz DB, Laleu B, Dick T. 2017. Screening of TB Activities 490 against Nontuberculous Mycobacteria Delivers High Hit Rates. Front Microbiol 491 8:1539.

30. Preston S, Jiao Y, Jabbar A, McGee SL, Laleu SB, Willis P, Wells TNC, Gasser 492 493 RB. 2016. Screening of the 'Pathogen Box' identifies an approved pesticide with 494 major anthelmintic activity against the barber's pole worm. Int J Parasitol Drugs 495 Drug Resist 6(3):329-334.

496 31. Spalenka J, Escotte-Binet S, Bakiri A, Hubert J, Renault JH, Velard F, Duchateau S, 497 Aubert D, Huguenin A, Villena I. 2018. Discovery of New Inhibitors of Toxoplasma gondii via the Pathogen Box. Antimicrob Agents Chemother 62(2). pii: 498 499 e01640-17.

500 32. Jeong J, Kim G, Moon C, Kim HJ, Kim TH and Jang J. 2018. Pathogen Box 501 screening for hit identification against Mycobacterium abscessus. PLoS One 13(4).

502 33. Müller J, Aguado A, Laleu B, Balmer V, Ritler D, Hemphill A. 2017. In vitro 503 screening of the open source Pathogen Box identifies novel compounds with 504 profound activities against Neospora caninum. In J Parasitol. 47(12):801-809.

34. Meier A, Erler H, Beitz E. 2018. Targeting Channels and Transporters in Protozoan 505 506 Parasite Infections. Front Chem 6:88.

507 35. Spangenberg T, Burrows JN, Kowalczyk P, McDonald S, Wells TN, Willis P. 2013.
508 The open access malaria box: a drug discovery catalyst for neglected diseases. PLoS
509 One 8(6):e62906.

- 36. Duffy S, Sykes ML, Jones AJ, Shelper TB, Simpson M, Lang R, Poulsen SA,
 Sleebs BE, Avery VM. 2017. Screening the Medicines for Malaria Venture
 Pathogen Box across Multiple Pathogens Reclassifies Starting Points for OpenSource Drug Discovery. Antimicrob Agents Chemother 61(9).
- 514 37. Vila T, Lopez-Ribot JL. 2016. Screening the Pathogen Box for Identification of
 515 Candida albicans Biofilm Inhibitors. Antimicrob Agents Chemother 61(1).
- 516 38. Mayer FL, Kronstad JW. 2017. Discovery of a Novel Antifungal Agent in the
 517 Pathogen Box. mSphere 2(2).
- 518 39. McCarthy MW, Walsh TJ. 2017. Drugs currently under investigation for the
 519 treatment of invasive candidiasis. Expert Opin Investig Drugs 26(7):825-831.
- 40. The Pathogen Box. Available from URL <u>https://www.pathogenbox.org</u>. [accessed
 April 2017].
- 41. Boray JC. 1969. Experimental fascioliasis in Australia. Adv Parasitol 8: 95-210.
- 42. Valero MA, Mas-Coma S. 2000. Comparative infectivity of Fasciola hepatica
 metacercariae from isolates of the main and secondary reservoir animal host species
 in the Bolivian Altiplano high human endemic region. Folia Parasitol (Praha)
 47(1):17-22.

527 43. Duthaler U, Smith T, Keiser J. 2010. In Vivo and In Vitro Sensitivity of Fasciola
528 hepatica to Triclabendazole Combined with Artesunate, Artemether, or OZ78.
529 Antimicrob Agents Chemother 54(11):4596-604.

530 44. ChEMBL. Available from URL <u>https://www.ebi.ac.uk/chembl/</u> [accessed
531 September 2017].

45. Bento AP, Gaulton A, Hersey A, Bellis LJ, Chambers J, Davies M, Krüger FA,
Light Y, Mak L, McGlinchey S, Nowotka M, Papadatos G, Santos R, Overington
JP. 2014. The ChEMBL bioactivity database: an update. Nucleic Acids Res
42(Database issue):D1083-90.

- 46. lazar. Available from URL <u>https://lazar.in-silico.de/predict</u> [accessed October
 2017].
- 47. Wexler P. (2004) The U.S. National Library of Medicine's Toxicology and
 Environmental Health Information Program. Toxicology 198(1-3):161-8.

48. Maunz A, Gütlein M, Rautenberg M, Vorgrimmler D, Gebele D, Helma C. 2013.
lazar: a modular predictive toxicology framework. Front Pharmacol 4:38.

542 49. Skehan P, Storeng R, Scudiero D, Monks A, McMahon J, Vistica D. Warren JT,
543 Bokesch H, Kenney S, Boyd MR. 1990. New colorimetric cytotoxicity assay for
544 anticancer-drug screening. J Natl Cancer Inst 82(13):1107-12.

545 50. Boyd MR, Paull KD. 1995. Some practical considerations and applications of the
546 National Cancer Institute in vitro anticancer drug discovery screen. Drug Dev Res
547 34:91–109.

548	51. Brennan GP, Fairweather I, Trudgett A, Hoey E, McCoy, McConville M, Meaney
549	M, Robinson M, McFerran N, Ryan L, Lanusse C, Mottier L, Alvarez L, Solana H,
550	Virkel G, Brophy PM. 2007. Understanding triclabendazole resistance. Exp Mol
551	Pathol 82(2):104-9.
552	52. Sargison ND, Scott PR. 2011. Diagnosis and economic consequences of
553	triclabendazole resistance in Fasciola hepatica in a sheep flock in south-east

triclabendazole resistance in Fasciola hepatica in a sheep flock in south-Scotland. Vet Rec 168(6):159.

- 555 53. Sargison N. 2012. Diagnosis of triclabendazole resistance in Fasciola hepatica. Vet
 556 Rec 171(6):151-2.
- 557 54. Novobilský A, Averpil HB, Höglund J. 2012. The field evaluation of albendazole
 and triclabendazole efficacy against Fasciola hepatica by coproantigen ELISA in
 naturally infected sheep. Vet Parasitol 190(1-2):272-6
- 560 55. Novobilský A, Höglund J. 2015. First report of closantel treatment failure against
 561 Fasciola hepatica in cattle. Int J Parasitol Drugs Drug Resist 5(3):172-7.
- 562 56. Tong JX, Chandramohanadas R, Tan KS. 2018. High-Content Screening of the
 563 Medicines for Malaria Venture Pathogen Box for Plasmodium falciparum Digestive
 564 Vacuole-Disrupting Molecules Reveals Valuable Starting Points for Drug
 565 Discovery. Antimicrob Agents Chemother 62(3).
- 566 57. Fairweather I, Boray JC. 1999. Fasciolicides: efficacy, actions, resistance and its
 567 management. Vet J 158(2):81-112.
- 568 58. Lacey E. 1988. The role of the cytoskeletal protein, tubulin, in the mode of action
 and mechanism of drug resistance to benzimidazoles. Int J Parasitol 18(7):885-936.

570	59. Duthaler U, Smith TA, Keiser J. 2010. In Vivo and In Vitro Sensitivity of Fasciola
571	hepatica to Triclabendazole Combined with Artesunate, Artemether, or OZ78.
572	Antimicrob Agents Chemother 54(11):4596–4604.
573	60. Farahnak A, Golmohamdi T, Eshraghian M. 2012. In vitro Effects of
574	Triclabendazole (TCBZ) on the Excretory-Secretory Products (ESP) of Fasciola spp
575	Parasites. Acta Medica Iranica 50(3): 164-7.
576	

577

578

Accepted Manuscript Posted Online

of

Antimicrobial Agents and Chemotherapy

AAC





Table 1. Summary of the chemical compounds that showed the best biological activity against metacercariae

						IC50 μM ⁴			In silico toxicity features 6					
Compound plate code ¹	MMV ID ²	Molecular Formula	Molecular weight (g/mol)	Mean % mortality on MC ³	SD(%)		R	Other infectious microorganisms ⁵	Acute cytotoxicity (Fathead minnow)	Blood Brain Barrier Penetration (Human)	Carcinogenicity (Rodent)	Mutagenicity (Salmonella typhimurium)	Maximum Recommended Daily Dose (Human)	
TCBZ	N.A.	C14H9Cl3N2OS	359.7	100 +	0 †	15*	N.A.	Schistosoma	4.57 (mg/L)	penetrating	non-carcinogenic	non-mutagenic	N.A.	
PAA2	MMV010764	C14H16N4OS2	320.4	22	38.5	24.1	-0.3	Plasmodium	N.A.	N.A.	non-carcinogenic	non-mutagenic	N.A.	
PAF4	MMV676388	C15H14N4O3S	330.4	29	24.7	16.9	0.8	Mycobacterium	254.0 (mg/L)	penetrating	carcinogenic	mutagenic	2.44 (mg/kg_bw/day)	
PAF5	MMV202553	C15H15N3O2	269.3	29	24.7	14.9	0.9	Kinetoplastids	7.58 (mg/L)	penetrating	non-carcinogenic	mutagenic	0.993 (mg/kg_bw/day)	
PAG6	MMV063404	C19H24N3OCI	345.9	54	7.2	5.3	1.0	Mycobacterium	N.A.	penetrating	carcinogenic	mutagenic	N.A.	
PAH6	MMV676539	C20H16N2O3	332.4	17	28.9	24.7	-1.0	Mycobacterium	25.9 (mg/L)	penetrating	carcinogenic	mutagenic	4.05 (mg/kg_bw/day)	
PBD3	MMV637953	C51H40N6O23S6	1435.3	25	9.9	21.8	-0.6	Irypanosoma and Onchocerca	N.A.	penetrating	non-carcinogenic	non-mutagenic	N.A.	
PBD7	MMV019838	C18H10N4OF6	412.3	26	11.6	12.4	0.0	Plasmodium	N.A.	penetrating	non-carcinogenic	mutagenic	N.A.	
PBF4	MMV003270	C7H5N2OCI	168.6	26	25.1	8.2	-0.7	Ancylostoma	6.75 (mg/L)	penetrating	non-carcinogenic	non-mutagenic	15.5 (mg/kg_bw/day)	
PBF6	MMV688853	C19H23N5O2	389.9	25	22.5	31.9	-0.8	Cryptosporidium	N.A.	non-penetrating	non-carcinogenic	mutagenic	N.A.	
PBF11	MMV085210	C22H24N3O3CIS	446.0	40	15.3	2.4	0.8	Plasmodium	N.A.	penetrating	non-carcinogenic	non-mutagenic	1.64 (mg/kg_bw/day)	
PBH10	MMV676380	C18H15N4O3Cl	370.8	33	33.3	1.3	0.1	Plasmodium	132.0 (mg/L)	penetrating	non-carcinogenic	non-mutagenic	101.0 (mg/kg_bw/day)	
PCA2	MMV675997	C24H29N4O2F	424.5	22	38.4	18.1	-0.2	Kinetoplastids	N.A.	penetrating	non-carcinogenic	mutagenic	1.51 (mg/kg_bw/day)	
PCA6	MMV688852	C16H17N5CIF	333.8	29	37.4	17.2	-0.7	Toxoplasma	N.A.	penetrating	non-carcinogenic	mutagenic	N.A.	
PCC2	MMV688508	C19H19N2O4F	358.4	26	3.7	16.9	-0.5	Mycobacterium	N.A.	penetrating	non-carcinogenic	mutagenic	N.A.	
PCC5	MMV687730	C22H32N4O2	384.5	28	13.4	0.4	-0.5	Mycobacterium	N.A.	penetrating	carcinogenic	non-mutagenic	N.A.	
PCC6	MMV687251	C8H9N3O4S2	275.3	30	12.0	0.3	-0.5	Mycobacterium	N.A.	penetrating	non-carcinogenic	non-mutagenic	13.3 (mg/kg_bw/day)	
PCC9	MMV688361	C21H19N5O	357.4	32	11.5	17.2	-0.7	Kinetoplastids	N.A.	penetrating	carcinogenic	mutagenic	N.A.	
PCC10	MMV689029	C26H26N4O4S	490.6	33	19.1	10.5	0.8	Kinetoplastids	N.A.	penetrating	carcinogenic	mutagenic	11.9 (mg/kg bw/day)	
PCD11	MMV1030799	C20H18N4O	330.4	28	11.7	1.5	-0.3	Plasmodium	6.62 (mg/L)	non-penetrating	carcinogenic	mutagenic	N.A.	
PCE5	MMV687146	C19H26N2O	298.4	21	25.8	15.6	0.6	Mycobacterium	N.A.	penetrating	non-carcinogenic	mutagenic	N.A.	
PCE6	MMV687696	C29H28N4O2CIF3	557.0	26	20.6	18.2	-0.7	Mycobacterium	N.A.	non-penetrating	carcinogenic	mutagenic	N.A.	
PCE7	MMV687170	C17H13N4O2CI	340.8	34	25.3	13.1	0.0	Mycobacterium	N.A.	penetrating	carcinogenic	mutagenic	N.A.	
PCE8	MMV690102	C22H23N7O2	417.5	38	15.6	2.1	0.7	Kinetoplastids	N.A.	penetrating	non-carcinogenic	mutagenic	3.27 (mg/kg_bw/day)	
PCE11	MMV1029203	C20H17N5OS	375.5	33	29.7	7.1	-0.4	Plasmodium	100.0 (mg/L)	penetrating	carcinogenic	mutagenic	N.A.	
PCF2	MMV676053	C18H16N3O3CI	357.8	38	12.5	1.9	0.6	Cryptosporidium	194.0 (mg/L)	penetrating	non-carcinogenic	mutagenic	0.991 (mg/kg_bw/day)	
PCF3	MMV688179	C18H16N6OCI2	476.2	35	32.0	3.1	-0.1	Kinetoplastids	4.62 (mg/L)	penetrating	carcinogenic	mutagenic	1.41 (mg/kg_bw/day)	
PCF4	MMV023969	C24H24N4O5	453.0	48	21.8	1.5	0.3	Mycobacterium	N.A.	N.A.	carcinogenic	mutagenic	N.A.	
PCF5	MMV687138	C19H17NO35	339.4	26	11.6	14.7	-0.2	Mycobacterium	524.0 (mg/L)	penetrating	non-carcinogenic	mutagenic	89.7 (mg/kg_bw/day)	
PCF11	MMV688921	C23H18N3O5Cl	451.9	31	43.0	2.4	-0.4	Aedes aegypti -	N.A.	penetrating	carcinogenic	mutagenic	N.A.	
PCG9	MMV688891	C18H11NO4BrF3	442.2	25	10.9	25.7	-0.5	Mycobacterium	N.A.	penetrating	carcinogenic	mutagenic	1.25 (mg/kg_bw/day)	
PDH11	MMV688980	C16H18N3O2F5	335.4	33	38.2	21.2	0.2	Plasmodium	N.A.	penetrating	carcinogenic	mutagenic	N.A.	
PEC8	MMV687765	C25H26N6O	463	28	25.5	20.6	-0.8	Mycobacterium	N.A.	penetrating	non-carcinogenic	mutagenic	N.A.	
PEG9	MMV084864	C17H12N60	316.3	40	18.7	17.3	0.8	Plasmodium	14.2 (mg/l)	penetrating	non-carcinogenic	mutagenic	N.A.	

Table 2. Biological activity of the compounds screened on adult worms.

Compound plate code ¹	MMV ID ²	Mean % Mortality on adults ³	SD (%)
TCBZ	N.A.	100 +	0
PAG6	MMV063404	67	33.3
PBF4	MMV003270	67	0
PBF11	MMV085210	0	0
PBH10	MMV676380	78	19.2
PCC5	MMV687730	11	19.2
PCC6	MMV687251	33	33.3
PCD11	MMV1030799	67	33.3
PCE8	MMV690102	56	19.2
PCE11	MMV1029203	78	19.2
PCF2	MMV676053	0	0
PCF3	MMV688179	22	19.2
PCF4	MMV023969	33	33.3
PCF11	MMV688921	67	33.3

² ID codes assigned by the MMV agency.

³ Measured at 48-hr post drug exposure on adult worms. Results are means and standard deviation from triplicate experiments at a concentration of 33 µM.

† Mean and standard deviation of 6 individual experiments performed in 3 plates Compounds in italics were selected for IC50 on adult worms and cytotoxicity assay on cell lines. N.A. is not applicable

Table 3. Hits compounds selected for their fasciocidal activity as a	a new effective drugs against F. hepatica.

	Compound ID/Drug name ¹	Drug name	Molecular weight	AlogP ²	In vitro fasciocidal assesment 3		In vitro cytotoxicity 4				Cytotoxicity data from other studies 5			
Structure/Smiles'					Adult IC _{so} (µM)	CI 95%	3T3 GI50 (µM)	H460 GI50 (µM)	DU145 GI50 (µM)	HT29 GI50 (μM)	HepG2 CC20 (µM)	HL60 CC50 (µM)	MRC5 CC50 (µM)	
~tacb~	Triclabendazole	6-Chloro-5-(2,3- dichlorophenoxy)-2- (methylthio)-1H- benzo(dlimidazole	359.66	6		15*	22.80	32.62	35.22	37.80				
CSC1=NC2=C(N1)C=C(Cl)C(OC1=CC=CC(Cl)=C1Cl)=C2		benzolajimaazoie									N.A.	N.A.	N.A.	
	MMV003270/Zoxazolamine	2-Amino-5- chlorobenzoxazole	168.58	2.1	9.37	1.45 to 53.88	> 23.73	> 23.73	> 23.73	> 23.73	> 80	N.A.	N.A.	
CIC1=CC(N=C(N)O2)=C2C=C1														
a - O - o - o - o - o - o - o - o - o - o	MMV676380	N-(4-Acetamidophenyl)-3-(5- chloro-2-hydroxyphenyl)-1H- pyrazole-5-carboxamide	370.79	3.7	6.68	4.39 to 10.06	> 10.79	> 10.79	> 10.79	> 10.79	> 80	> 50	N.A.	
CC(=O)Nc1ccc(NC(=O)c2cc([nH]n2)c3cc(Cl)ccc3O)cc1														
	MMV690102	2-N-[1-[4-(4- methoxyphenoxy)phenyl]et hyl]-2-N- methylpyrimido[4,5- d]pyrimidine-2,5,7-triamine	417.46	3.6	2.14	1.16 to 4.82	4.86	0.95	9.58	11.00	2.87	N.A.	5.44	
	MMV1029203	N-methyl-2-[[5-phenyl-2-{2- pyridyl]thieno[3,2- e]pyrimidin-4- yl]amino]acetamide	375.45	3.58	4.32	2.82 to 6.60	>10.65	> 10.65	> 10.65	> 10.65	22	N.A.	N.A.	
D reder arrianed by the MMU agency														

TIP code service and a service

te tests elocytic leukemia MRC5 is fib

_

AAC

Antimicrobial Agents and Chemotherapy Table 4. Potential targets of the 13 hits and TCBZ tested in adult worms assays.

Compound plate code ¹	MMV code ²	Target ³					
		Targets predicted	CHEMBL ID	Preferred name	Organism	Protein target classification	
PBH10	MMV676380	3	CHEMBL2535	Glucose transporter	Homo sapiens	transporter > electrochemical transporter > slc superfamily of solute carriers > slc02 family of hexose and sugar alcohol transporters	
			CHEMBL4697	Hexose transporter 1	Plasmodium falciparum		
DCE11	MMMV1020202	1	CHEMBL3431938	Glucose transporter	Leishmania mexicana	transporter	
PCF2	MMV676053	1	CHEMBL6145	Inosine-5'-monophosphate dehydrogenase, probable	Cryptosporidium parvum	enzyme	
PCF4	MMV023969	3	CHEMBL2535	Glucose transporter	Homo sapiens	transporter > electrochemical transporter > slc superfamily of solute carriers > slc02 family of hexose and sugar alcohol transporters	
			CHEMBL4697 CHEMBL3431938	Hexose transporter 1 Glucose transporter	Plasmodium falciparum Leishmania mexicana	unclassified protein	
PBF4	MMV003270	19	CHEMBL340	Cytochrome P450 3A4	Homo sapiens	family 3 > cytochrome p450 family 3a > cytochrome p450 family 3a > cytochrome p450 family 3a > cytochrome p450 3a4	
			CHEMBL289	Cytochrome P450 2D6	Homo sapiens	enzyme > cytochrome p450 > cytochrome p450 family 2 > cytochrome p450 family 2d > cytochrome p450 2d6	
			CHEMBL3397	Cytochrome P450 2C9	Homo sapiens	enzyme > cytochrome p450 > cytochrome p450 family 2 > cytochrome p450 family 2c > cytochrome p450 2c9	
			CHEMBL3622	Cytochrome P450 2C19	Homo sapiens	enzyme > cytochrome p450 > cytochrome p450 family 2 > cytochrome p450 family 2c > cytochrome p450 2c19	
			CHEMBL3356	Cytochrome P450 1A2	Homo sapiens	enzyme > cytochrome p450 > cytochrome p450 family 1 > cytochrome p450 family 1a > cytochrome p450 1a2	
			CHEMBL4040	MAP kinase ERK2	Homo sapiens	enzyme > kinase > protein kinase > cmgc protein kinase group > cmgc protein kinase mapk family > cmgc protein kinase erk subfamily	

AAC

			CHEMBL2903	Arachidonate 15-lipoxygenase	Homo sapiens	enzyme
			CHEMBL2756	Monoamine oxidase B	Bos taurus	enzyme
			CHEMBL3254	Monoamine oxidase A	Bos taurus	enzyme
			CHEMBL1075094	Nuclear factor erythroid 2- related factor 2	Homo sapiens	unclassified protein
			CHEMBL1293224	Microtubule-associated protein tau	Homo sapiens	unclassified protein
			CHEMBL2362981	TAR DNA-binding protein 43	Homo sapiens	unclassified protein
			CHEMBL1293235	Prelamin-A/C	Homo sapiens	unclassified protein
			CHEMBL1781865	78 kDa glucose-regulated protein	Homo sapiens	unclassified protein
			CHEMBL1977	Vitamin D receptor	Homo sapiens	transcription factor > nuclear receptor > nuclear hormone receptor subfamily 1 > nuclear hormone receptor subfamily 1 group i > nuclear hormone receptor subfamily 1 group i member 1
			CHEMBL1947	Thyroid hormone receptor beta-1	Homo sapiens	hormone receptor subfamily 1 > nuclear hormone receptor subfamily 1 group a > nuclear hormone receptor subfamily 1 group a member 2
			CHEMBL1697668	Solute carrier organic anion transporter family member 1B1	Homo sapiens	transporter > electrochemical transporter > slc superfamily of solute carriers > slc21/slco family of organic anion transporting polypeptides
			CHEMBL1743121	Solute carrier organic anion transporter family member 1B3	Homo sapiens	
			CHEMBL1741193	Chromobox protein homolog 1	Homo sapiens	epigenetic regulator > reader > methyl- lysine/arginine binding protein > chromodomain
TCBZ	N.A.	27	CHEMBL1293278	Geminin	Homo sapiens	unclassified protein
			CHEMBL1075094	Nuclear factor erythroid 2- related factor 2	Homo sapiens	unclassified protein
			CHEMBL1293224	Microtubule-associated protein tau	Homo sapiens	unclassified protein
			CHEMBL1293258	Mothers against decapentaplegic homolog 3	Homo sapiens	unclassified protein
			CHEMBL2362981	TAR DNA-binding protein 43	Homo sapiens	unclassified protein

CHEMBL2146310	Aberrant vpr protein	Human immunodeficiency virus 1	unclassified protein	
CHEMBL2029198	Rap guanine nucleotide exchange factor 4	Homo sapiens	unclassified protein	
CHEMBL6152	Alpha-synuclein	Homo sapiens	unclassified protein	
CHEMBL1293191	Transcriptional regulator ERG	Homo sapiens	unclassified protein	
CHEMBL2007624 CHEMBL1795086 CHEMBL5567	Peripheral myelin protein 22 HSP90 Luciferin 4-monooxygenase	Rattus norvegicus Plasmodium falciparum 3D7 Photinus pyralis	unclassified protein unclassified protein enzyme enzyme	
CHEMBL2007625	[NADP] cytoplasmic	Homo sapiens		
CHEMBL3563	Cruzipain	Trypanosoma cruzi	enzyme > protease > cysteine protease > cysteine protease ca clan > cysteine protease c1a family	
CHEMBL1293248	4'-phosphopantetheinyl transferase ffp	Bacillus subtilis	enzyme	
CHEMBL1795087	Ubiquitin carboxyl-terminal hydrolase 1	Homo sapiens	enzyme	
CHEMBL1293234	Putative fructose-1,6- bisphosphate aldolase	Giardia intestinalis	enzyme	
CHEMBL1293228	Streptokinase A	Streptococcus pyogenes serotype M1	enzyme > kinase	
CHEMBL2524	Alpha-galactosidase A	Homo sapiens	enzyme	
CHEMBL1784	Glucagon-like peptide 1 receptor	Homo sapiens	membrane receptor > family b g protein-coupled receptor > peptide receptor (family b gpcr) > glucagon-like receptor > glucagon-like peptide receptor	
CHEMBL1793	Parathyroid hormone receptor	Homo sapiens	membrane receptor > family b g protein-coupled receptor > peptide receptor (family b gpcr) > parathyroid hormone receptor > parathyroid hormone receptor	
CHEMBL5162	Neuropeptide S receptor	Homo sapiens	membrane receptor > family a g protein-coupled receptor > peptide receptor (family a gpcr) > short peptide receptor (family a gpcr) > neuropeptide receptor	

transcription factor > nuclear receptor > nuclear

P	
gents ar apy	
bial A	
ntimicro Che	

			CHEMBL1293231	Nuclear receptor ROR-gamma	Mus musculus	hormone receptor subfamily 1 > nuclear hormone receptor subfamily 1 group f > nuclear hormone receptor subfamily 1 group f member 3 transcription factor > nuclear receptor > nuclear
			CHEMBL1871	Androgen Receptor	Homo sapiens	hormone receptor subfamily 3 > nuclear hormone receptor subfamily 3 group c > nuclear hormone receptor subfamily 3 group c member 4
			CHEMBL3880	Heat shock protein HSP 90- alpha	Homo sapiens	other cytosolic protein
			CHEMBL6032	Histone-lysine N- methyltransferase, H3 lysine- 9 specific 3	Homo sapiens	epigenetic regulator > writer > protein methyltransferase
			CHEMBL4377	Guanine nucleotide-binding protein G(s), subunit alpha	Homo sapiens	other membrane protein
PAG6	MMV063404	No target	N.A.	N.A.	N.A.	N.A.
PCC5	MMV687730	No target	N.A.	N.A.	N.A.	N.A.
PCC6	MMV687251	No target	N.A.	N.A.	N.A.	N.A.
PCD11	MMV1030799	No target	N.A.	N.A.	N.A.	N.A.
PCE8	MMV690102	No target	N.A.	N.A.	N.A.	N.A.
PBF11	MMV085210	No target	N.A.	N.A.	N.A.	N.A.
PCF3	MMV688179	No target	N.A.	N.A.	N.A.	N.A.
PCF11	MMV688921	No target	N.A.	N.A.	N.A.	N.A.

¹ Coordinates used to identify compounds in each plate

² ID codes assigned by the MMV agency.
 ³ By consulting ChEMBL.
 N.A is not available

AAC