



**Sexual Functioning in Spanish and Portuguese Young Adults: Initial Validation of the Arizona Sexual Experience Scale**

Journal:	<i>International Journal of Sexual Health</i>
Manuscript ID	WIJS-2016-1580.R1
Manuscript Type:	Original Article
Keywords:	Adults, Cross-cultural studies, Sexual function

SCHOLARONE<sup>TM</sup>  
Manuscripts

## SEXUAL FUNCTIONING IN YOUNG ADULTS

**Abstract**

**Objectives:** We conducted an initial validation of the Arizona Sexual Experience Scale (ASEX) and examined the sexual function of Spanish and Portuguese young adults. **Methods:** Participants were 523 Spanish and 595 Portuguese heterosexual young adults who completed a background questionnaire, the ASEX, and measures of sexual sensation seeking, excitation and inhibition. **Results:** The ASEX showed good reliability and validity. Participants reported good sexual function. The most prevalent difficulties in men were related to desire and orgasm, and to orgasm and satisfaction in women. **Conclusions:** Sexual health programs for adolescents should incorporate sexual function given its importance in their sexual life.

*Keywords:* Sexual functioning, Sexual difficulties, Young adults, Validation, Arizona Sexual Experience Scale

## SEXUAL FUNCTIONING IN YOUNG ADULTS

**Sexual Functioning in Spanish and Portuguese Young Adults**

Sexual functioning is an important component of sexual health that relates to interpersonal relationships, sexual well-being and quality of life (Lauman, Paik, & Rosen, 1999; World Health Organization, 2010). This has led to growing interest in the study of sexual functioning and problems associated with sexual dysfunction (DeRogatis, 2008; World Health Organization, 2002). Studies to date show a high prevalence of sexual problems and dysfunctions in general population (Simons & Carey, 2001). Laumann et al. (1999) found that sexual problems affected 31% of men and 43% of women, also finding that women reported higher prevalence of sexual dysfunctions than men did. Among men, the most prevalent problems were premature ejaculation, erectile problems and problems related to orgasm, whereas among women, the most frequent were lack of interest, inability to achieve orgasm, and sexual pain (Casas Aranda et al., 2007; Farmer & Meston, 2007; Gomes & Nobre, 2014; Laumann et al., 1999; Mercer et al., 2003; Peixoto & Nobre, 2015). While some other variables that are related to sexual health, such as sexual risk behaviors and STIs, have been the focus of research in adolescents and young adults for decades, the study of sexual function in this population has received very little attention (O'Sullivan, Brotto, Byers, Majerovich, & Wuest, 2014; O'Sullivan & Majerovich, 2008). In a study conducted in Canada (Fisher & Boroditsky, 2000) with women between 18 and 29 years, 33% of the participants reported lack of sexual desire, 22% had experienced sexual pain, and 31% had trouble reaching orgasm. O'Sullivan and Majerovich (2008), in a study conducted with men and women between 17 and 21 years, found that 32.6% of men or more and 61.4% of women or more had experienced at least one sexual difficulty in their lives. However, the authors found that chronic sexual difficulties were very rare and overall sexual function was positive. In this study, 48.7% of men had had problems maintaining an

## SEXUAL FUNCTIONING IN YOUNG ADULTS

1  
2  
3 erection, 43.6% had suffered premature ejaculation, and 32.6% reported sexual pain during  
4  
5 intercourse. Among the women, 86.7% had had difficulty achieving orgasm, 81.2% had lost  
6  
7 interest in sex, and 75.8% had experienced lubrication problems. Recently, O'Sullivan et al.  
8  
9 (2014), using a sample of adolescents from 16 to 21 years, found that 51.1% of the adolescents  
10  
11 who participated in their study had suffered sexual problems. The most common problems in  
12  
13 women were low sexual interest (26-43%), inability to reach orgasm (18-41%), and low arousal  
14  
15 (16-38%), and in men, premature ejaculation (12-31%), erectile problems (13-28%), and low  
16  
17 sexual interest (13-28%). Overall, these findings suggest high prevalence of sexual difficulties in  
18  
19 adolescents and young adults and also a great variability in the results from different studies. The  
20  
21 results found in studies with adolescents and young adults are somewhat similar to those  
22  
23 obtained with general population. Based on this some authors have suggested that the first sexual  
24  
25 experiences can set the basis for adult sexual functioning (Koch, 1988; O'Sullivan &  
26  
27 Majerovich, 2008), justifying the study of sexual functioning in adolescence and young  
28  
29 adulthood as a way to prevent and manage sexual problems in adulthood.  
30  
31  
32  
33  
34  
35

36 To date, the study of sexual function in adolescents and young adults is still sparse and  
37  
38 most research has been conducted in Canada and the United States. In Spain and Portugal there  
39  
40 is very little research examining sexual functioning. Most of the research that exists has focused  
41  
42 on only one area of sexual functioning, such as sexual satisfaction or sexual desire, and has  
43  
44 examined some aspects other than its prevalence (Carrobes, Gámez-Guadix, & Almendros,  
45  
46 2011; Santos-Iglesias, Calvillo, & Sierra, 2013). Other studies have focused on the study of  
47  
48 sexual functioning but have used samples from the general population without examining  
49  
50 adolescents and young adults in particular (Casas Aranda et al., 2007; Gomes & Nobre, 2014;  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## SEXUAL FUNCTIONING IN YOUNG ADULTS

Martín-Morales et al., 2001; Peixoto & Nobre, 2015; Sierra, Vallejo-Medina, Santos-Iglesias, & Lameiras Fernández, 2012; Teles et al., 2008).

**Arizona Sexual Experience Scale**

Sexuality research conducted in Spain and Portugal has faced another difficulty, that is the scarcity of available instruments to assess sexual functioning. Of the large number of instruments that have been developed for the assessment of sexual function in English-speaking contexts, only a few of them are available in Spanish and Portuguese language. Some of the existing ones are long and intrusive (Pechorro, Diniz, Almeida, & Vieira, 2009; Bobes et al., 2000); were designed to specifically assess sexual function in either men or women (IIEF; Quinta Gomes & Nobre, 2012); or have been validated or used only in clinical samples (Ávila Escribano, Pérez Madruga, Olazábal Ulacia, & López Fidalgo, 2004; García-Portilla et al., 2011; Vallejo-Medina & Sierra, 2012).

McGahuey et al. (2000) developed the Arizona Sexual Experience Scale (ASEX) as a valid, reliable, short, easy-to-use, nonintrusive and with clinical utility. The ASEX is composed of five items that assess five basic areas of sexual functioning in men and women regardless of their sexual orientation: desire, arousal, erection/vaginal lubrication, ability to reach orgasm, and satisfaction with orgasm. The male and female versions differ on the third question. Items are rated on a scale with six response options, ranging from 1 (*hyperfunction*) to 6 (*hypofunction*). Both item and global scores can be interpreted, with higher scores indicating greater sexual dysfunction. McGahuey et al. (2000) proposed that a global score greater than or equal to 19, a score greater than or equal to 5 on any item, or any three items with a score greater than or equal to 4, were indicative of sexual dysfunction.

## SEXUAL FUNCTIONING IN YOUNG ADULTS

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26

The ASEX has some appropriate characteristics for a scale developed to assess sexual dysfunction (DeRogatis, 2008; Meston & DeRogatis, 2002). Previous studies have shown adequate values of internal consistency reliability ( $\alpha = .90$ ) and test-retest reliability ( $r = .80 - .89$ ). It has shown sensitivity and specificity to discriminate sexually functional and dysfunctional individuals (McGahuey et al., 2000). The scale has been subsequently validated in different countries (e.g., France, China, Turkey, Brazil, Tunisia) and with different clinical samples (e.g., anxiety, depression, schizophrenia, Parkinson, kidney patients; Briki et al., 2014; Byerly, Nakonezny, Fisher, Magourik, & Rush, 2006; Jitkrittadukul, Jagota, & Bhidayasiri, 2014; Lin, Juang, Wen, Liu, & Hung, 2012; Nakhli et al., 2014; Nunes et al., 2014; Soykan, 2004), showing its utility regardless of cultural differences or medical conditions.

**The Present Study**

27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

The present study had two objectives. The first goal was to conduct an initial validation of the Spanish and Portuguese translations of the Arizona Sexual Experience Scale. For this purpose, we translated the ASEX into Spanish and Portuguese (see Method for description of translation) and we examined the psychometric properties of each version separately. First, we examined construct validity through an exploratory factor analysis, and we also examined whether the male and female versions were equivalent across country by testing measurement invariance. Then, we analyzed the scale's internal consistency (Cronbach's alpha) and item properties (i.e., mean, standard deviation, and item-total corrected correlation). In order to determine concurrent validity, the relationship between the ASEX scores and a series of associated constructs was examined. To determine divergent validity, we examined the correlation of the scale with constructs that are not associated with sexual functioning. Finally, using ROC curves, we examined discriminant validity by analyzing the ability of the scale to

## SEXUAL FUNCTIONING IN YOUNG ADULTS

differentiate people who had suffered some sexual difficulties from those who had not. We made the following predictions:

H1: Better sexual functioning would be related to higher *sexual excitation* (Bancroft, Graham, Janssen, & Sanders, 2009), higher *sexual sensation seeking* (Johnson, Phelps, & Cottler, 2004), and to lower *sexual inhibition due to the threat of performance failure* (Bancroft et al., 2009).

H2: Sexual functioning and *sexual inhibition due to the threat of performance consequences* would be weakly or not significantly correlated (Bancroft et al., 2009)

The second goal of this study was to examine sexual functioning in male and female young adults in Spain and Portugal. We predicted that:

H3: Women in Spain and Portugal would report poorer sexual function than men (Laumann et al., 1999).

## Methods

### Participants and Procedure

Participants were recruited from a mid-size university in Spain and a large-size university in Portugal through a non-random sampling procedure using each university e-mail distribution list for students. Each student registered on these distribution lists received an e-mail with the purpose of the study, the inclusion criteria (i.e., being a university student at the time of the study, ages ranging from 18 to 26 years old), contact information of the principal investigator, and the link to an online survey. Only those participants who gave their informed consent could access the survey. After completing the survey, they indicated whether they would like to receive summary results of the study, and whether they would like to enter a raffle to win an iPad mini™. Those who responded affirmatively to either question gave their name and an e-mail address,

## SEXUAL FUNCTIONING IN YOUNG ADULTS

1  
2  
3 which were stored on a separate, secure database. These studies were approved by the regional  
4  
5 Ethics Review Board for Clinical Research in Spain and the university Ethics Board of the  
6  
7 Psychology School in Portugal.  
8  
9

10 The final Spanish sample was composed of 520 heterosexual participants (26.20% men).  
11  
12 All participants ranged in age from 18 to 26 years and most of them were in a current romantic  
13  
14 relationship. Most participants had had vaginal sexual intercourse and around 20% of  
15  
16 participants reported anal sexual intercourse. Sample descriptive statistics for men and women  
17  
18 can be found in Table 1. The Portuguese sample consisted of 587 heterosexual young adults  
19  
20 (27.40% men). All participants' ages ranged from 18 to 26 years old and the majority was in a  
21  
22 current romantic relationship. Most participants had engaged in vaginal intercourse and around  
23  
24 30% had engaged in anal intercourse. Three participants from Spain and eight from Portugal  
25  
26 were deleted because they did not have any sexual experience by the time of the study.  
27  
28 Descriptive statistics can be found in Table 1. T-test comparisons revealed that men were older,  
29  
30 had shorter romantic relationships, were older than women when they had their first vaginal  
31  
32 sexual contact, and had a greater number of vaginal sexual partners than women. Comparison of  
33  
34 the Spanish and Portuguese samples showed that more Portuguese students had had vaginal and  
35  
36 anal sexual relationships than Spanish. Also, Portuguese students were younger, had a greater  
37  
38 number of anal sexual partners, and were younger than Spanish when they had their first anal  
39  
40 sexual contact.  
41  
42  
43  
44  
45  
46  
47

**Measures**

48  
49 **Background questionnaire.** Used to gather information about gender, age, sexual  
50  
51 orientation, relationship status, and relationship length.  
52  
53  
54  
55  
56  
57  
58  
59  
60



## SEXUAL FUNCTIONING IN YOUNG ADULTS

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
*Sexual history.* We asked participants whether they had engaged in vaginal sexual contacts with penetration, age of first vaginal sexual contact with penetration, and number of vaginal sexual partners. Parallel questions were used to examine anal sexual experience. Participants also reported whether they had experienced any sexual difficulty in their lifetime.

13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
*Arizona Sexual Experience Scale* (McGahuey et al., 2000). The scale has been described in the Introduction. Participants responded on a scale from 1 (*hyperfunction*) to 6 (*hypofunction*). Because the ASEX do not assess sexual pain, we added another item (i.e., “How strong is your pain in your genitals (your sexual parts) during your sexual activity/intercourse?”). This item was responded using the same response scale from 1 (*No pain at all*) to 6 (*Extremely strong*). However, this item was only used for descriptive purposes and was not included as part of the scale validation.

29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
*Sexual Excitation/Sexual Inhibition Scales – Short Form* (Carpenter, Janssen, Graham, Vorst, & Wicherts, 2011). The Spanish (Moyano & Sierra, 2014) validation and the Portuguese (Quinta Gomes, Janssen, Pinto-Gouveia, & Nobre, 2016) translation were used in this study. Composed of 14 items to assess sexual excitation (SES), sexual inhibition due to the threat of performance failure (SIS1), and sexual inhibition due to the threat of performance consequences (SIS2). Participants responded on a Likert-type scale ranging from 1 (*Strongly agree*) to 4 (*Strongly disagree*). After reversing items’ scores, higher scores indicated greater sexual excitation and inhibition. The scale has good reliability and adequate validity (Carpenter et al., 2011; Moyano & Sierra, 2014; Quinta Gomes et al., 2016). In this study, Cronbach’s alpha values in Spain were .79 (SES), .70 (SIS1), and .58 (SIS2) for men, and .80, .72, and .67 for women, respectively. In Portugal, values were .67, .68, and .54 for men, and .72, .58, and .61 for women.

## SEXUAL FUNCTIONING IN YOUNG ADULTS

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

*Sexual Sensation Seeking Scale* (Kalichman, 2011). The Spanish (Teva & Bermúdez, 2008) and Portuguese (Santos Pechorro et al., 2015) validations were used. It is composed of 10 items to assess sexual sensation seeking on a Likert scale ranging from 1 (*Not at all like me*) to 4 (*Very much like me*). Higher scores indicated greater sexual sensation seeking. Authors reported appropriate validity and reliability (Kalichman, 2011; Santos Pechorro et al., 2015; Teva & Bermúdez, 2008). In the present study, Cronbach's alpha values were .64 for men and .73 for women in Spain, and .57 for men and .55 for women in Portugal.

**Development of the Spanish and Portuguese Version of the ASEX**

The English version of the ASEX (McGahuey et al., 2000) was translated into Spanish by an expert in sexuality research using a forward translation procedure. Both the translated and the original version were given to a bilingual expert in translating psychological and sexological manuscripts to assure the correspondence between the two versions. Then the Spanish translation was sent to two experts in psychological assessment and sexuality research to identify and suggest changes to items that were not clear and understandable. Changes were made when both experts suggested the same modification. No changes were made in this phase of the study. Finally, the resulting version was given to 10 individuals with characteristics similar to the final sample. They were given the same task as experts in psychological assessment and sexuality research. An 80% of agreement was required to make further changes to the items. No changes were made in this phase either (see Appendix). The same procedure was used for the Portuguese translation.

**Results****Arizona Sexual Experiences Scale***Exploratory Factor Analysis*

## SEXUAL FUNCTIONING IN YOUNG ADULTS

Parallel analysis suggested the extraction of one factor. In Spain, results for the men showed a single factor solution that explained 48.37% of the variance (eigenvalue = 2.42). All factor loadings were greater than .60. For the women, a single factor solution explained 48.26% of the variance (eigenvalue = 2.41) with factor loadings greater than .63. In Portugal, results for the men showed a single factor solution that explained 46.85% of the variance (eigenvalue = 2.34). All factor loadings were greater than .53. For the Portuguese women, a single factor solution explained 50.69% of the variance (eigenvalue = 2.53) and all factor loadings were greater than .67 (see Table 2).

***Factorial Invariance***

Factorial invariance across country was tested separately for the male and female version of the ASEX with Mplus 7.0 and using a robust maximum likelihood estimator. Four models were assessed: (a) configural invariance constrained the number of factors and the pattern of free and fixed loadings across both groups; (b) weak invariance tested equality of factor loadings across groups; (c) strong invariance tested equality of intercepts for both groups; and (d) strict invariance assumed that residual variances for all items were equal across groups (Dimitrov, 2010; Wang & Wang, 2012). Comparative Fit Index (CFI) and Tucker-Lewis Index (TLI) values above .90 and Root Mean Square Error of Approximation (RMSEA) under .08 were indicators of good fit (Hu & Bentler, 1999; Wang & Wang, 2012). Additionally, to assess the fit of nested models, we examined the scaled chi-square difference test (Satorra & Bentler, 2010) and changes in CFI values. According to Cheung and Rensvold (2002) a decrease in  $CFI \leq .01$  is a good indicator of measurement invariance.

The results for men showed configural, weak invariance, and partial strong invariance (item 1 was set free to vary across countries). Strict invariance did not hold across country and

## SEXUAL FUNCTIONING IN YOUNG ADULTS

the modification indices did not suggest to free any of the residual variances. The results for women showed configural invariance, partial weak invariance (item 2), and partial strong invariance (item 1) (see Table 3). Similar to the male version, strict invariance was not reached.

***Item Analysis and Reliability***

In Spain, item analysis for men showed mean responses slightly below the theoretical midpoint of the scale and standard deviations below 1.0, which indicates fairly good sexual functioning in all areas, and the items were below desirable levels of discrimination. Corrected item-total correlations were over .25 (Nunnally & Bernstein, 1994), and Cronbach's alpha for the overall scale was .73 (see Table 4). For women, all items showed a mean slightly under the theoretical midpoint of the scale, except for the orgasm item. Standard deviations were below 1.0, except for the orgasm and satisfaction items. All corrected item-total correlations were greater than .25 (Nunnally & Bernstein, 1994), and Cronbach's alpha was .73. In Portugal, item analysis for men showed mean responses below the midpoint of the scale and standard deviations under 1.0, indicating good sexual functioning. Corrected item-total correlations were greater than .25 (Nunnally & Bernstein, 1994), and Cronbach's alpha for the overall scale was .71. In Portuguese women, item means were closer to the midpoint of the scale and standard deviations close or over 1.0, which indicates an average sexual function and items with good discrimination. All corrected item-total correlations were over .25, and Cronbach's alpha was .75 (see Table 4).

***Validity***

Zero-order correlations were used to examine our concurrent and divergent validity hypotheses. As predicted we found better sexual functioning to be related to higher sexual excitement, lower sexual inhibition due to the threat of performance failure, and greater sexual

## SEXUAL FUNCTIONING IN YOUNG ADULTS

sensation seeking both in men and women (H1). On the other hand, as expected, we found that sexual functioning was not correlated or weakly associated with sexual inhibition due to the threat of performance consequences in Spain (H2). Similar results were found in Portugal, except for the relationship between sexual function and sexual inhibition due to the threat of performance consequences, that was statistically significant in both men and women in Portugal (see Table 5).

With respect to discriminant validity of the ASEX, we performed ROC analysis. The criterion standard was whether or not participants reported that they had suffered any sexual difficulty in the past. In Spain, the area under the curve was statistically significant ( $AUC = .76 \pm .05, p < .001$ ). A cut-off score of 15 discriminated those who had experienced some sexual difficulty in the past with a sensitivity of .64 and specificity of .80. In Portugal the area under the curve was also significant ( $AUC = .73 \pm .03, p < .001$ ). A cut-off score of 15 discriminated those who had experienced some sexual difficulty in the past with a sensitivity of .67 and specificity of .72. A binary logistic regression was used to predict the dependent variable (i.e., having experienced or not a sexual difficulty in the past) from the ASEX score. In Spain, a significant model ( $\chi^2 = 25.67, p < .001$ ) correctly classified 72.3% of the sample. The results showed a significant model in Portugal ( $\chi^2 = 25.67, p < .001$ ) that correctly classified a 69.7% of the sample. These results indicate that in both samples the ASEX scores could well discriminate participants who had reported some sexual complaint in the past from those who had not.

### **Sexual Functioning in Male and Female Young Adults**

Table 6 shows the score distributions for the different areas of sexual function in Spanish men and women, as well as the percentage of participants that could be classified as having sexual dysfunctions according to McGahuey et al. (2000) criteria. Men showed a good sexual

## SEXUAL FUNCTIONING IN YOUNG ADULTS

function overall, however, 7.30% of men could be classified as sexually dysfunctional. The most prevalent sexual problem among Spanish men was their ability to reach an orgasm and the least prevalent one was their level of sexual satisfaction. Overall, women showed poorer sexual function than men. A total of 18.40% of women were classified as having sexual dysfunction. The most problematic area for women was their ability to reach an orgasm and the least frequent was their lubrication. A mixed MANOVA was used to compare each of the areas of sexual function (except erection and lubrication) between men and women. The areas of sexual function were treated as within-subjects factor and gender as a between-subjects factor. Results showed a significant effect for the areas of sexual function,  $F(4, 477) = 110.35, p < .001, \eta_p^2 = .48$ ; gender,  $F(1, 480) = 62.97, p < .001, \eta_p^2 = .12$ ; and the interaction of area and gender,  $F(4, 477) = 6.17, p < .001, \eta_p^2 = .05$ . All areas of sexual function differed significantly among them ( $p < .001$ ) except for desire and arousal. Follow-up of the interaction effect showed that women scored significantly higher than men in all areas of sexual function ( $p < .001$ ). For men, all areas were significantly different among them ( $p < .001$ ) except for desire and arousal and desire and orgasm. For the women, all areas were significantly different ( $p < .001$ ) except for desire and arousal.

Men in Portugal showed an overall good function, however 4.90% could be classified as having sexual dysfunction. The most problematic area of sexual function was their sexual interest and the least frequent was their ability to reach an orgasm and their sexual satisfaction (see Table 7). Of the 384 women in Portugal, 28.70% were classified as sexually dysfunctional. The most frequent sexual problem was their ability to reach an orgasm and the least frequent was their sexual interest. Same analysis was conducted to compare the areas of sexual function and gender effects. Results showed a significant effect for the areas of sexual function,  $F(4, 566) =$

## SEXUAL FUNCTIONING IN YOUNG ADULTS

1  
2  
3 102.98,  $p < .001$ ,  $\eta_p^2 = .42$ ; gender,  $F(1, 569) = 80.83$ ,  $p < .001$ ,  $\eta_p^2 = .12$ ; and the interaction of  
4  
5  
6 area and gender,  $F(4, 566) = 9.12$ ,  $p < .001$ ,  $\eta_p^2 = .06$ . All areas of sexual function differed  
7  
8 significantly among them ( $p < .001$ ) except for desire and satisfaction. Follow-up of the  
9  
10 interaction effect showed that women scored significantly higher than men in all areas of sexual  
11  
12 function ( $p < .001$ ). For the men, all areas of sexual function were significantly different ( $p <$   
13  
14  $.001$ ) except for desire and satisfaction, and arousal and orgasm. For the women, all areas were  
15  
16 significantly different ( $p < .001$ ) except for desire and satisfaction only.  
17  
18

**Discussion**

19  
20  
21  
22 This study presents the psychometric properties of the first Spanish and Portuguese  
23  
24 adaptation of the Arizona Sexual Experience Scale (McGahuey et al., 2000). The scale shows  
25  
26 good reliability and validity in general. Additionally, the sexual functioning of Spanish and  
27  
28 Portuguese young adults was examined. Results show that participants have overall adequate  
29  
30 levels of sexual functioning, although sexual difficulties are reported frequently.  
31  
32

33  
34 The exploratory factor analysis of the ASEX showed a univariate structure with good  
35  
36 reliability in both men and women in Spain and Portugal. The Spanish and Portuguese male and  
37  
38 female versions of the ASEX were not completely equivalent. Differences were found in the  
39  
40 factor loadings and the intercepts of items 2 and 1, respectively. These could indicate that these  
41  
42 items might have different meanings across the two groups and suggest that comparisons based  
43  
44 on these items and the factor cannot be made (Gregorich, 2006). Reliability values were lower  
45  
46 than those found in previous studies (Briki et al., 2014; Byerly et al., 2006; Jitkrittadukul et al.,  
47  
48 2014; Lin et al., 2012; McGahuey et al., 2000; Nakhli et al., 2014; Nunes et al., 2014; Soykan,  
49  
50 2004). These differences could be explained by the use of clinical samples in previous studies.  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
Clinical populations generally score higher in the ASEX and have more than one area of sexual

## SEXUAL FUNCTIONING IN YOUNG ADULTS

1  
2  
3 functioning affected (McGahuey et al., 2000), making the scores of the different items to be  
4  
5 more consistent among them. We found that the ASEX could discriminate between participants  
6  
7 who had experienced at least one sexual difficulty in the past. However, this does not mean that  
8  
9 the Spanish and Portuguese validations can be used as clinical tools for detecting sexual  
10  
11 dysfunction already. Future studies should examine their discriminant validity with clinical  
12  
13 populations in order to establish clinically significant cut-off scores.  
14  
15

16  
17 The validity results support the proposed hypotheses (Bancroft et al., 2009; Johnson et al.,  
18  
19 2004) for the majority of cases. We expected that sexual function would be weakly or not  
20  
21 correlated to sexual inhibition due to threat of performance consequences (SIS2), however we  
22  
23 found this association to be significant and moderate in Portugal. Past research indicates that  
24  
25 SIS2 could measure a tendency to respond with inhibited sexual function when a threat is present  
26  
27 (Janssen & Bancroft, 2007). It might be that the context in which sex occurs for young adults  
28  
29 (e.g., lack of sexual experience and sexual education, social pressure to engage in sex; García,  
30  
31 Cañadas-De la Fuente, González-Jiménez, Fernández-Castillo, & García-García, 2011;  
32  
33 O'Sullivan et al., 2014; O'Sullivan & Majerovich, 2008; van de Bongardt, Reitz, Sandfort, &  
34  
35 Dekovic, 2015; Veiga, Teixeira, Martins, & Meliço-Silvestre, 2006) might be perceived as  
36  
37 threatening and therefore affects their sexual function. Additionally, a previous study conducted  
38  
39 in Portugal has shown that men with erectile dysfunction scored significantly higher on both  
40  
41 SIS1 and SIS2 compared to sexually healthy controls (Quinta Gomes, Janssen, Gamesan, &  
42  
43 Nobre, 2016). More research is needed to examine the contextual and cultural factors and  
44  
45 predictors that affect Spanish and Portuguese young adults' sexual function. Based on these  
46  
47 results, we could say that the ASEX provides a fast, simple, and fairly nonintrusive appraisal of  
48  
49 the sexual functioning of individuals (McGahuey et al., 2000), meeting most of the US Food and  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



## SEXUAL FUNCTIONING IN YOUNG ADULTS

1  
2  
3 Drug Administration standards for scales assessing sexual functioning (DeRogatis, 2008). This  
4  
5 adaptation, in particular, provides a valid and reliable instrument that can be used in Spain and  
6  
7 Portugal and in Spanish and Portuguese-speaking contexts.  
8  
9

10 The results obtained from the analysis of sexual functioning allow us to draw several  
11  
12 conclusions. First, a high percentage of participants show adequate sexual functioning in all the  
13  
14 different areas. This allows us to conclude that this sample of university students have levels of  
15  
16 sexual function that are appropriate for a young and healthy population. Actually, our results  
17  
18 reveal lower rates of sexual difficulties than other studies using young adults from universities  
19  
20 (O'Sullivan & Majerovich, 2008) or the general population (Laumann et al., 1999). Second,  
21  
22 while sexual functioning is generally positive, there is still a percentage of young adults –  
23  
24 between 0.60 and 3.10% of the men and between 1.10 and 22.50% of the women- who report  
25  
26 having severe problems in at least one area of sexual functioning (O'Sullivan & Majerovich,  
27  
28 2008). Furthermore, the percentage of participants that are classified as having a sexual  
29  
30 dysfunction using the criteria from McGahuey et al. (2000) ranges from 4.90% in Portuguese  
31  
32 men to 28.70% in Portuguese women, indicating that despite of the overall good sexual function,  
33  
34 sexual dysfunction can have a significant presence in younger adults. However, the percentage of  
35  
36 participants reporting sexual difficulties in this study is lower than the percentage found  
37  
38 previously in the general population and in other studies with adolescents and young adults  
39  
40 (Farmer & Meston, 2007; Fisher & Boroditsky, 2000; Laumann et al., 1999; O'Sullivan et al.,  
41  
42 2014; O'Sullivan & Majerovich, 2008; Peixoto & Nobre, 2015). This could be explained by the  
43  
44 age difference between the general population and our sample. Older adults have higher rates of  
45  
46 biological and psychological problems and unhealthy lifestyles that adversely impact their sexual  
47  
48 functioning (Breslin, Karmakar, Smith, Etches, & Mustard, 2007). Differences could also be  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## SEXUAL FUNCTIONING IN YOUNG ADULTS

1  
2  
3 explained by the short time frame used to assess sexual functioning in this study (i.e., one week),  
4  
5 decreasing the likelihood of participants suffering from sexual problems. Finally, the results also  
6  
7 show that women have poorer sexual functioning than men across all different areas (Farmer &  
8  
9 Meston, 2007; Fisher & Boroditsky, 2000; Laumann et al., 1999; O'Sullivan & Majerovich,  
10  
11 2008). These differences could be due to biological, psychological, and/or social causes  
12  
13 (O'Sullivan et al., 2014; Westheimer & Lopater, 2004). Future studies should examine the  
14  
15 reasons for these differences. As found in previous studies (Gomes & Nobre, 2014; Laumann et  
16  
17 al., 1999; O'Sullivan et al., 2014; O'Sullivan & Majerovich, 2008; Peixoto & Nobre, 2015;  
18  
19 Simons & Carey, 2001), our work shows that the most common problems for women are  
20  
21 difficulty achieving orgasm and a lack of satisfaction with orgasms. For men, the most frequent  
22  
23 problems were difficulty achieving orgasm and a lack of sexual desire, which is inconsistent with  
24  
25 previous studies which indicate that the most frequent problems for men were premature  
26  
27 ejaculation and a lack of sexual desire (Laumann et al., 1999; O'Sullivan et al., 2014; O'Sullivan  
28  
29 & Majerovich, 2008; Simons & Carey, 2001). However, the ASEX does not distinguish between  
30  
31 premature ejaculation and delayed ejaculation, making it difficult to compare our results with  
32  
33 previous studies.  
34  
35  
36  
37  
38  
39

40  
41 Although this study introduces relevant and novel results about the sexual functioning of  
42  
43 young adults in Spain and Portugal, these results must be interpreted with caution. First, although  
44  
45 this first validation of the ASEX has appropriate psychometric properties, it does not examine  
46  
47 areas of sexual functioning, such as premature and delayed ejaculation (American Psychiatric  
48  
49 Association, 2013; Peixoto & Nobre, 2014). Future studies need to document other sources of  
50  
51 reliability (i.e., test-retest) and validity, such as discriminant validity using clinical samples and  
52  
53 construct validity using other scales to assess sexual function. The sample is composed only of  
54  
55  
56  
57  
58  
59  
60

## SEXUAL FUNCTIONING IN YOUNG ADULTS

heterosexual young adults and mostly women, which limits the generalizability of the results.

Finally, some of the measures used for validation purposes showed low reliability, which could have impacted the results of this study.

Overall, this study fills some gaps in the literature on sexual functioning in Spain and Portugal by providing some relevant data on the sexual problems and dysfunction of young adults. Additionally, it validates a scale for the assessment of sexual functioning in Spanish and Portuguese-speaking contexts. The results show that in spite of the adequate sexual functioning in young adults overall, they also experience some problems in sexual functioning and, therefore, it is a topic of research that needs more attention. Knowledge about the adolescents' and young adults' sexual functioning as well as early detection of sexual problems and dysfunctions is essential to promote a healthy sexual lifestyle. These results can be used to inform the development of sexual health education programs, in which sexual function is frequently neglected.

### Acknowledgements

Part of this research was funded by the "Antonio Gargallo-Ibercaja" Foundation (Spain) and granted to Dr. Ángel Castro, and by the Fundação para a Ciência e Tecnologia (Portugal). Authors would like to thank Ciaran Rowe for his helpful comments on the translation of the scale.

### References

American Psychiatric Association (2013). *Diagnostic and Statistical Manual of Mental Disorders*. Fifth ed. Arlington, VA: American Psychiatric Publishing.

Ávila Escribano, J. J., Pérez Madruga, A., Olazábal Ulacia, J. C., & López Fidalgo, J. (2004).

## SEXUAL FUNCTIONING IN YOUNG ADULTS

Disfunciones sexuales en alcoholismo. *Adicciones*, 16, 1-6.

Bancroft, J., Graham, C. A., Janssen, E., & Sanders, S. A. (2009). The Dual Control Model: Current status and future directions. *Journal of Sex Research*, 46, 121-142. doi: 10.1080/00224490902747222

Bobes, J., González, M. P., Rico-Villandemoros, F., Bascaran, M. T., Sarasa, P., & Clayton, A. (2000). Validation of the Spanish version of the Changes in Sexual Functioning Questionnaire (CSFQ). *Journal of Sex & Marital Therapy*, 26, 119-131. doi: 10.1080/009262300278524

Breslin, F. C., Karmakar, S. D., Smith, P., Etches, J., & Mustard, C. (2007). Time allocation between work and recreation and the associated injury risks among young people. *Journal of Safety Research*, 38, 373-379. doi: 10.1016/j.jsr.2007.01.009

Breyer, B. N., Smith, J. F., Eisenberg, M. L., Ando, K. A., Rowen, T. S., & Shindel, A. W. (2010). The Impact of Sexual Orientation on Sexuality and Sexual Practices in North American Medical Students. *Journal of Sexual Medicine*, 7, 2391-2400. doi: 10.1111/j.1743-6109.2010.01794.x

Briki, M., Haffen, E., Monnin, J., Tio, G., Nicolier, M., Sechter, D., & Vandiel, P. (2014). Sexual dysfunction and depression: Validity of a French version of the ASEX scale. *Encephale*, 40, 114-122. doi: 10.1016/j.encep.2012.10.008

Byerly, M. J., Nakonezny, P. A., Fisher, R., Magouirk, B., & Rush, J. (2006). An empirical evaluation of the Arizona Sexual Experiences Scale and a simple one-item screening test for assessing antipsychotic-related sexual dysfunction in outpatients with schizophrenia and schizoaffective disorder. *Schizophrenia Research*, 81, 311-316. doi: 10.1016/j.schres.2005.08.013

## SEXUAL FUNCTIONING IN YOUNG ADULTS

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
- Carpenter, D. L., Janssen, E., Graham, C. A., Vorst, H., & Wicherts, J. (2011). The Sexual Inhibition/Sexual Excitation Scales-Short Form (SIS/SES-SF). In T. D. Fisher, C. M. Davis, W. L. Yarber, & S. L. Davis (Eds.), *Handbook of Sexuality-Related Measures* (pp. 236-239). London, UK: Routledge.
- Carrobbles, J. A., Gámez-Guadix, M., & Almendros, C. (2011). Sexual functioning, sexual satisfaction, and subjective and psychological well-being in Spanish women. *Anales de Psicología, 27*, 27-34.
- Casas Aranda, I., Guirao Sánchez, L., García-Giralda, L., Alfaro González, J. V., Sandoval Martínez, C., & García-Giralda, F. J. (2007). SyDSF-AP: un cuestionario de 21 ítems para aproximarnos a la salud sexual y detectar las disfunciones sexuales femeninas en Atención Primaria. *Semergen, Medicina de Familia, 33*, 9-13.
- Cheung, G. W., & Rensvold, R. B. (2002). Evaluating goodness-of-fit indexes for testing measurement invariance. *Structural Equation Modeling, 9*, 235-255.  
doi:10.1207/S15328007SEM0902\_5
- De Rogatis, L. R. (2008). Assessment of sexual function/dysfunction in patient reported outcomes. *International Journal of Impotence Research, 20*, 35-44.
- Dimitrov, D. M. (2010). Testing for factorial invariance in the context of construct validation. *Measurement and Evaluation in Counseling and Development, 43*, 121-149.  
doi:10.1177/0748175610373459
- Farmer, M. A., & Meston, C. M. (2007). Predictors of genital pain in young women. *Archives of Sexual Behavior, 36*, 831-843. doi: 10.1007/s10508-007-9199-7
- Fisher, W. A., & Borodistky, R. (2000). Sexual activity, contraceptive choice, and sexual and reproductive health indicators among single Canadian women aged 15-29: Additional

## SEXUAL FUNCTIONING IN YOUNG ADULTS

- 1  
2  
3 findings from the Canadian Contraceptive Study. *Canadian Journal of Human Sexuality*,  
4 9, 79-93.  
5  
6  
7  
8 García, M. A., Cañadas-De la Fuente, G. A., González-Jiménez, E., Fernández Castillo, R., &  
9  
10 García-García, I. (2011). Educar en conductas sexuales saludables: una innovación  
11 docente en promoción de la salud. *Revista Médica de Chile*, 139, 1269-1275. doi:  
12 10.4067/S0034-98872011001000004  
13  
14  
15  
16  
17 García-Portilla, M. P., Sáiz, P. A., Fonseca, E., Al-Halabi, S., Bobes-Bascarán, M. T., Arrojo,  
18  
19 M., ... Bobes, J. (2011). Psychometric Properties of the Spanish Version of the Changes  
20 in Sexual Functioning Questionnaire Short-Form (CSFQ-14) in Patients with Severe  
21 Mental Disorders. *Journal of Sexual Medicine*, 8, 1371-1382. doi: 10.1111/j.1743-  
22 6109.2010.02043.x  
23  
24  
25  
26  
27  
28  
29 Gomes, A. L. Q., & Nobre, P. Q. (2014). Prevalence of sexual problems in Portugal: Results of a  
30 population-based study using a stratified sample of men aged 18 to 70 years. *Journal of*  
31 *Sex Research*, 51, 13-21. doi: 10.1080/00224499.2012.744953  
32  
33  
34  
35  
36 Gregorich, S. E. (2006). Do self-report instruments allow meaningful comparisons across diverse  
37 population groups? Testing measurement invariance using the confirmatory factor  
38 analysis framework. *Medical Care*, 44, S78-S94.  
39  
40  
41  
42  
43 Hu, L., & Bentler, P. M. (1999). Cutoff criteria for fit indexes in covariance structure analysis:  
44 Conventional criteria versus new alternatives. *Structural Equation Modeling*, 6, 1-55.  
45  
46  
47  
48  
49  
50  
51 Janssen, E., & Bancroft, J. (2007). The Dual Control Model: The role of sexual inhibition and  
52 excitation in sexual arousal and behavior. In E. Janssen (Ed.), *The psychophysiology of*  
53 *sex* (pp. 197-222). Bloomington, IN: Indiana University Press.  
54  
55  
56  
57  
58  
59  
60

## SEXUAL FUNCTIONING IN YOUNG ADULTS

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
- Jitkritisadakul, O., Jagota, P., & Bhidayasiri, R. (2014). The Arizona Sexual Experiences Scale: A Validity and Reliability Assessment of the Thai Translation (ASEX-Thai) in Parkinson's Disease. *Journal of Parkinson's Disease*, 4, 205-210. doi: 10.3233/JPD\_130271
- Johnson, S. D., Phelps, D. L., & Cotler, L. B. (2004). The association of sexual dysfunction and substance use among a community epidemiological sample. *Archives of Sexual Behavior*, 33, 55-63. doi: 10.1023/B:ASEB.0000007462.97961
- Kalichman, S. J. (2011). Sexual Sensation Seeking Scale. In T. D. Fisher, C. M. Davis, W. L. Yarber, & S. L. Davis (Eds.), *Handbook of Sexuality-Related Measures* (pp. 564-565). London, UK: Routledge.
- Koch, P. B. (1988). The relationship of first intercourse to later sexual functioning concerns in adolescents. *Journal of Adolescent Research*, 3, 345-362. doi: 10.1777/074355488833009
- Laumann, E. O., Paik, A., & Rosen, R. C. (1999). Sexual dysfunction in the United States: Prevalence and predictors. *Journal of American Medical Association*, 281, 537-544. doi: 10.1001/jama.281.6.537
- Lin, C. F., Juang, Y. Y., Wen, J. K., Liu, C. Y., & Hung, C. I. (2012). Correlations between sexual dysfunction, depression, anxiety, and somatic symptoms among patients with major depressive disorder. *Chang Gung Medical Journal*, 35, 323-331.
- Martín-Morales, A., Sánchez-Cruz, J. J., de Tejada, I. S., Rodríguez-Vela, L., Jiménez-Cruz, J. F., & Burgos-Rodríguez, R. (2001). Prevalence and independent risk factors for erectile dysfunction in Spain: Results of the Epidemiologia de la Disfuncion Erectil Masculina study. *Journal of Urology*, 166, 569-574. doi: 10.1016/S0022-5347(05)65986-1

## SEXUAL FUNCTIONING IN YOUNG ADULTS

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
- McGahuey, C. A., Gelenberg, A. J., Laukes, C. A., Moreno, F. A., Delgado, P. L., McKnight, K. M., & Manber, R. (2000). The Arizona Sexual Experience Scale (ASEX): Reliability and Validity. *Journal of Sex & Marital Therapy*, 26, 25-40. doi: 10.1080/009262300278623
- Mercer, C. H., Fenton, K. A., Johnson, A. M., Wellings, K., Macdowall, W., McManus, S., ... Erens, B. (2003). Sexual function problems and help seeking behaviour in Britain: national probability sample survey. *British Medical Journal*, 327, 426-427. doi: 10.1136/bmj.327.7412.426
- Meston, C., & DeRogatis, L. R. (2002). Validated instruments for assessing female sexual function. *Journal of Sex & Marital Therapy*, 28, 155-164. doi: 10.1080/00926230252851276
- Moyano, N., & Sierra, J. C. (2014). Validación de las Escalas de Inhibición/Excitación Sexual-Forma Breve (EIS/EES-FB). *Terapia Psicológica*, 32, 87-100.
- Nakhli, J., El Kissi, Y., Bouhlel, S., Amamou, B., Nabli, T. A., Nasr, S. B., & Ali, B. B. (2014). Reliability and validity of the Arizona Sexual Experiences Scale-Arabic version in Tunisian patients with schizophrenia. *Comprehensive Psychiatry*, 55, 1473-1477. doi: 10.1016/j.comppsy.2014.04.006
- Nunes, L. V. A., Dieckmann, L. H. J., Lacaz, F. S., Bressan, R., Matsuo, T., & Mari, J. D. (2009). The accuracy of the Arizona Sexual Experience Scale (ASEX) to identify sexual dysfunction in patients of the schizophrenia spectrum. *Revista de Psiquiatria Clínica*, 36, 197-204. doi: 10.1590/s0101-60832009000500002
- Nunnally, J. C., & Bernstein, I. H. (1994). *Psychometric theory*. New York, NY: McGraw-Hill.



## SEXUAL FUNCTIONING IN YOUNG ADULTS

O'Sullivan, L. F., Brotto, L. A., Byers, E. S., Majerovich, J. A., & Wuest, J. A. (2014).

Prevalence and characteristics of sexual functioning among sexually experienced middle to late adolescents. *Journal of Sexual Medicine*, *11*, 630-641. doi: 10.1111/jsm.12419

O'Sullivan, L. F., & Majerovich, J. A. (2008). Difficulties with sexual functioning in a sample of male and female late adolescents and young adult university students. *Canadian Journal of Human Sexuality*, *17*, 109-121.

Pechorro, P., Diniz, A., Almeida, S., & Vieira, R. (2009). Portuguese validation of Female Sexual Functioning Index. *Laboratório de Psicologia*, *7*, 33-44.

Peixoto, M. M., & Nobre, P. (2014). Prevalence of sexual problems and associated distress among gay and heterosexual men. *Sexual and Relationship Therapy*, *30*, 211-225. doi: 10.1080/14681994.2014.986084

Peixoto, M. M., & Nobre, P. (2015). Prevalence and sociodemographic predictors of sexual problems in Portugal: a population-based study with women aged 18 to 79 years. *Journal of Sex & Marital Therapy*, *41*, 169-180. doi: 10.1080/0092623X.2013.842195

Quinta Gomes, A. L., Janssen, E., Ganesan, A., & Nobre, P. J. (2016). Sexual inhibition and sexual excitation as psychobiological risk factors for erectile disorder. *Manuscript in preparation*.

Quinta Gomes, A.L., Janssen, E., Pinto-Gouveia, J., & Nobre, P. (2016). Validation of the Sexual Inhibition and Sexual Excitation Scales (SIS/SES) in Portugal: Assessing gender differences and predictors of sexual functioning. *Manuscript submitted for publication*.

Quinta Gomes, A. L. & Nobre, P. (2012). The International Index of Erectile Function (IIEF-15): Psychometric properties of the Portuguese version. *Journal of Sexual Medicine*, *9*, 180-187. doi: 10.1111/j.1743-6109.2011.02467.x

## SEXUAL FUNCTIONING IN YOUNG ADULTS

- 1  
2  
3 Santos-Iglesias, P., Calvillo, G., & Sierra, J. C. (2013). A further examination of Levine's model  
4 of sexual desire. *Psychology & Sexuality, 4*, 34-45. doi: 10.1080/19419899.2011.576697  
5  
6  
7  
8 Santos Pechorro, P., Monteiro Pascoal, P., Soares Figueredo, C., Almeida, A. I., Vieira, R. X., &  
9  
10 Jesus, S. N. (2015). Validação portuguesa da Escala de Busca de Sensações Sexuais  
11 (SSSS). *Revista Internacional de Andrología, 13*, 125-130. doi:  
12  
13 10.1016/j.androl.2014.11.003  
14  
15  
16  
17 Satorra, A., & Bentler, P. M. (2010). Ensuring positiveness of the scaled difference chi-square  
18 test statistic. *Psychometrika, 75*, 243-248. doi:10.1007/s11336-009-9135-y  
19  
20  
21  
22 Sierra, J. C., Vallejo-Medina, P., Santos-Iglesias, P., & Lameiras Fernández, M. (2012).  
23 Validación del Massachusetts General Hospital-Sexual Functioning Questionnaire  
24 (MGH-SFQ) en población española. *Atención Primaria, 44*, 516-526. doi:  
25  
26 10.1016/j.aprim.2012.02.004  
27  
28  
29  
30  
31 Simons, J. S., & Carey, M. P. (2001). Prevalence of sexual dysfunctions: results from a decade of  
32 research. *Archives of Sexual Behavior, 30*, 177-219. doi: 10.1023/A:1002729318254  
33  
34  
35  
36 Soykan, A. (2004). The reliability and validity of Arizona sexual experiences scale in Turkish  
37 ESRD patients undergoing hemodialysis. *International Journal of Impotence Research,*  
38  
39  
40  
41  
42 16, 531-534. doi: 10.1038/sj.ijir.3901249  
43  
44 Teles, A. G., Carreira, M., Alarcão V., Sociol, D., Aragüés, J. M., Lopes, L., ... Costa, J. G.  
45 (2008). Prevalence, severity, and risk factors for erectile dysfunction in a representative  
46 sample of 3,548 portuguese men aged 40 to 69 years attending primary healthcare  
47 centers: Results of the portuguese erectile dysfunction study. *Journal of Sexual Medicine,*  
48  
49  
50  
51  
52 5, 1317-1324. doi: 10.1111/j.1743-6109.2007.00745.x  
53  
54  
55  
56  
57  
58  
59  
60

## SEXUAL FUNCTIONING IN YOUNG ADULTS

- 1  
2  
3 Teva, I., & Bermúdez, M. P. (2008). Adaptación castellana y propiedades psicométricas de la  
4  
5 Escala de Búsqueda de Sensaciones Sexuales en adolescentes españoles. *Revista*  
6  
7 *Mexicana de Psicología*, 25, 129-137.  
8  
9
- 10 Vallejo-Medina, P., & Sierra, J. C. (2012). Adaptation, equivalence, and validation of the  
11  
12 Changes in Sexual Functioning Questionnaire-Drugs in a sample of drug-dependent men.  
13  
14 *Journal of Sex & Marital Therapy*, 39, 368-384. doi: 10.1080/0092623x.642493  
15  
16
- 17 van de Bongardt, R., Reitz, E., Sandfort, T., & Dekovic, M. (2015). A meta-analysis of the  
18  
19 relations between three types of peer norms and adolescent sexual behavior. *Personality*  
20  
21 *and Social Psychology Review*, 19, 203-234. doi: 10.1177/1088868314544223  
22  
23
- 24 Veiga, L., Teixeira, F., Martins, I., & Melo-Silvestre, A. (2006). Sexuality and human  
25  
26 reproduction: a study of scientific knowledge, behaviours and beliefs of Portuguese future  
27  
28 elementary schools teachers. *Sex Education*, 6, 17-29. doi: 10.1080/14681810500508915  
29  
30
- 31 Wang, J., & Wang, X. (2012). *Structural equation modeling. Applications using Mplus*.  
32  
33 Chichester, United Kingdom: Higher Education Press.  
34  
35
- 36 Westheimer, R., & Lopater, S. (2004). *Human sexuality: A psychosocial perspective*. Baltimore,  
37  
38 MD: Lippincot Williams & Wilkins.  
39  
40
- 41 World Health Organization (2006). *Defining sexual health: Report of a technical consultation on*  
42  
43 *sexual health, 28-31 January 2002*. Geneva: World Health Organization.  
44  
45
- 46 World Health Organization (2010). *Developing sexual health programmes. A framework for*  
47  
48 *action*. Geneva: World Health Organization.  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## SEXUAL FUNCTIONING IN YOUNG ADULTS

## Appendix

## ASEX Portugal

Para cada um dos seguintes itens indica o teu nível **GLOBAL** durante a **ÚLTIMA SEMANA**, incluindo **HOJE**.

1.- Quão forte (intenso) é o teu desejo sexual?

1	2	3	4	5	6
Extremamente forte (intenso)	Muito forte (intenso)	Algo forte (intenso)	Algo débil	Muito débil	Nenhum desejo sexual

2.- Com que facilidade consegues excitar-te sexualmente?

1	2	3	4	5	6
Extremamente fácil	Muito fácil	Algo fácil	Algo difícil	Muito difícil	Nunca me excito

3.- (**SÓ PARA HOMENS**) Com que facilidade consegues atingir e manter uma erecção?

1	2	3	4	5	6
Extremamente fácil	Muito fácil	Algo fácil	Algo difícil	Muito difícil	Nunca

4.- (**SÓ PARA MULHERES**) Com que facilidade a tua vagina fica molhada (lubrificada) durante as relações sexuais?

1	2	3	4	5	6
Extremamente fácil	Muito fácil	Algo fácil	Algo difícil	Muito difícil	Nunca

5.- Com que facilidade consegues atingir um orgasmo?

1	2	3	4	5	6
Extremamente fácil	Muito fácil	Algo fácil	Algo difícil	Muito difícil	Nunca consigo atingir orgasmo

6.- Quão satisfatórios são os teus orgasmos?

1	2	3	4	5	6
Extremamente satisfatórios	Muito satisfatórios	Algo satisfatórios	Algo insatisfatórios	Muito insatisfatórios	Não consigo atingir orgasmos

## SEXUAL FUNCTIONING IN YOUNG ADULTS

28

## ASEX Spain

Para cada uno de los siguientes ítems indica tu nivel **GLOBAL** durante la **ÚLTIMA SEMANA**, incluyendo **HOY**.

1.- ¿Cómo de fuerte (intenso) es tu deseo sexual?

1	2	3	4	5	6
Extremadamente fuerte (intenso)	Muy fuerte (intenso)	Algo fuerte (intenso)	Algo débil	Muy débil	Ningún deseo sexual

2.- ¿Con qué facilidad logras excitarte sexualmente?

1	2	3	4	5	6
Extremadamente fácil	Muy fácil	Algo fácil	Algo difícil	Muy difícil	Nunca me excito

3.- **(SOLO PARA HOMBRES)** ¿Con qué facilidad puedes conseguir y mantener una erección?

1	2	3	4	5	6
Extremadamente fácil	Muy fácil	Algo fácil	Algo difícil	Muy difícil	Nunca

4.- **(SOLO PARA MUJERES)** ¿Con qué facilidad se humedece (lubrica) tu vagina durante las relaciones sexuales?

1	2	3	4	5	6
Extremadamente fácil	Muy fácil	Algo fácil	Algo difícil	Muy difícil	Nunca

5.- ¿Con qué facilidad puedes lograr un orgasmo?

1	2	3	4	5	6
Extremadamente fácil	Muy fácil	Algo fácil	Algo difícil	Muy difícil	Nunca consigo un orgasmo

6.- ¿Cómo de satisfactorios son tus orgasmos?

1	2	3	4	5	6
Extremadamente satisfactorios	Muy satisfactorios	Algo satisfactorios	Algo insatisfactorios	Muy insatisfactorios	No puedo conseguir orgasmos

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

Table 1

*Sociodemographic characteristics of the Spanish and Portuguese samples*

Variable	Spain				Portugal				Gender	Country
	Men		Women		Men		Women			
	<i>M</i> / %	<i>SD</i>	<i>M</i> / %	<i>SD</i>	<i>M</i> / %	<i>SD</i>	<i>M</i> / %	<i>SD</i>		
Age	21.97	2.13	21.48	2.11	21.27	2.23	20.71	2.06	12.99***	26.12***
Romantic relationship (Yes)	64.90%		66.90%		64.00%		67.40%		0.73	.01
Length (years)	2.41	1.82	3.00	2.00	2.49	1.71	2.82	1.86	6.97**	0.07
Vaginal sexual contact (Yes)	93.30%		93.90%		92.50%		89.40%		0.52	4.47*
Age of first contact	17.54	1.73	17.07	1.66	17.38	2.01	17.15	1.77	7.60**	0.12
Number of partners	4.43	5.06	3.91	4.17	3.59	4.53	2.36	2.23	10.53**	19.84***
Anal sexual contact (Yes)	19.40%		22.20%		34.20%		26.50%		0.99	7.39**
Age of first contact	19.73	2.44	19.47	2.15	19.04	2.20	18.88	2.07	0.49	4.56*
Number of partners	1.27	0.53	1.48	2.08	1.80	1.51	1.21	0.57	0.98	0.48

\*\* *p* < .05, \*\* *p* < .01, \*\*\* *p* < .001

Table 2

*Exploratory factor analysis of the ASEX in men and women in Spain and Portugal*

Items	Spain		Portugal	
	Men	Women	Men	Women
	Loading	Loading	Loading	Loading
1. How strong is your sex drive?	.74	.68	.78	.72
2. How easily are you sexually aroused (turned on)?	.75	.76	.81	.73
3. Can you easily get and keep an erection? (Men only)	.72		.71	
3. How easily does your vagina become moist during sex? (Women only)		.63		.67
4. How easily can you reach an orgasm?	.64	.74	.53	.73
5. Are your orgasms satisfying?	.60	.66	.54	.72

Table 3

*Testing for factorial invariance of the ASEX across countries for men and women*

Model	$\chi^2$	df	$\Delta\chi^2$	$\Delta$ df	CFI	$\Delta$ CFI	TLI	RMSEA
<b>Men</b>								
Configural	19.75*	9			.955		.901	.09
Weak	21.52	13	1.41	4	.965	+ .01	.946	.06
Strong	46.12***	18	23.76***	5	.883	- .082	.870	.07
Partial strong <sup>b</sup>	25.33	17	4.04	4	.965	.000	.959	.06
Strict	38.91*	22	13.50*	5	.930	- .035	.936	.12
<b>Women</b>								
Configural	17.71*	7			.988		.967	.06
Weak	36.39***	11	18.64***	4	.973	- .015	.950	.07
Partial weak <sup>a</sup>	22.36*	10	4.51	3	.987	- .001	.973	.05
Strong	95.84***	15	70.19***	5	.913	- .074	.884	.11
Partial strong <sup>b</sup>	33.85**	14	11.34*	4	.979	- .008	.969	.06
Strict	82.62***	19	53.10***	5	.931	- .048	.928	.09

*Note:* *N* Spain = 131 men, 384 women. *N* Portugal = 158 men, 426 women. <sup>a</sup>The factor loading of item 2 was set free to vary across groups. <sup>b</sup>The intercept of item 1 was set free to vary across groups.

\*  $p < .05$ , \*\*  $p < .01$ ; \*\*\*  $p < .001$ .



Table 4

*Item analysis for men and women in Spain and Portugal*

Items	Spain						Portugal					
	Men			Women			Men			Women		
	M	SD	Corrected	M	SD	Corrected	M	SD	Corrected	M	SD	Corrected
	item-total correlation			item-total correlation			item-total correlation			item-total correlation		
Interest	2.46	0.84	.53	2.84	0.80	.46	2.07	1.01	.56	2.52	0.92	.50
Arousal	2.42	0.76	.55	2.87	0.78	.54	2.44	0.83	.62	2.95	0.85	.52
Erection	2.08	0.84	.52				2.26	1.06	.48			
Lubrication				2.43	0.86	.41				2.65	1.13	.47
Orgasm	2.61	0.94	.44	3.34	1.03	.57	2.60	0.76	.34	3.53	1.22	.57
Satisfaction	2.04	0.66	.41	2.32	1.01	.47	2.04	0.70	.34	2.42	1.31	.55
Total	11.62	2.82		13.79	3.11		11.40	2.99		14.06	3.87	

Table 5

*Zero-order correlations between sexual functioning and sexual arousal, inhibition, and sensation seeking in men and women*

	Sexual functioning			
	Spain		Portugal	
	Men	Women	Men	Women
Sexual excitation	-.37***	-.19**	-.28***	-.21***
Sexual inhibition due to the threat of performance failure	.32***	.30***	.25**	.37***
Sexual inhibition due to the threat of performance consequences	.08	.12*	.24**	.25***
Sexual sensation seeking	-.21*	-.30***	-.29***	-.28***

\*  $p < .05$ , \*  $p < .01$ , \*\*\*  $p < .001$

Table 6

*Percent of male and female sexual problems in Spain*

	Men						Dysfunction <sup>a</sup>
	1	2	3	4	5	6	
Interest	8.80	46.00	37.20	5.10	2.90	0	2.90
Arousal	8.10	47.10	39.00	4.40	1.50	0	1.50
Erection	21.90	53.30	19.00	4.40	1.50	0	1.50
Orgasm	9.70	37.30	35.80	14.20	3.00	0	3.00
Satisfaction	16.70	64.40	17.40	0.80	0.80	0	0.80
Pain	66.20	18.80	11.30	2.30	2.30	0	2.30
	Women						
Interest	2.10	29.40	50.90	14.50	2.90	0.30	3.20
Arousal	3.70	25.50	50.40	18.90	1.60	0	1.60
Lubrication	12.70	43.60	33.10	9.50	0.80	0.30	1.10
Orgasm	1.10	19.30	38.30	28.50	9.20	3.50	12.70
Satisfaction	11.50	61.20	19.10	4.10	0.30	3.80	4.10
Pain	37.00	32.40	18.90	10.30	0.50	0.80	1.30

*Note.* Lower scores indicate hyperfunction and higher scores hypofunction (1: Extremely strong/easily/satisfying/painful; 2: Very strong/easily/satisfying/painful; 3: Somewhat strong/easily/satisfying/painful; 4: Somewhat weak/difficult/unsatisfying/painful; 5: Very weak/difficult/unsatisfying/painful; 6: No sex drive/never aroused/never/never reach orgasm/Can't reach orgasm/Extremely painful). <sup>a</sup> Based on McGahuey et al. (2000).

Table 7

*Percent of male and female sexual problems in Portugal*

	Men						Dysfunction <sup>a</sup>
	1	2	3	4	5	6	
Interest	26.10	51.60	18.00	1.20	3.10	0	3.10
Arousal	11.80	41.00	39.10	6.80	1.20	0	1.20
Erection	22.40	52.20	3.70	21.10	0.60	0	0.60
Orgasm	6.80	36.60	46.60	9.90	0	0	0
Satisfaction	19.30	59.60	18.00	3.10	0	0	0
Pain	56.00	24.50	8.20	9.40	1.90	0	1.90
	Women						
Interest	10.40	41.20	35.00	10.90	2.30	0.20	2.50
Arousal	3.50	24.50	45.40	23.80	2.80	0	2.80
Lubrication	14.80	34.20	28.60	17.80	3.00	1.60	4.60
Orgasm	3.10	17.30	31.00	26.20	14.90	7.60	22.50
Satisfaction	19.40	49.10	17.80	3.60	2.80	7.30	10.10
Pain	32.20	25.20	20.20	17.50	4.30	0.50	4.80

*Note.* 1: Extremely strong/easily/satisfying/painful; 2: Very strong/easily/satisfying/painful; 3: Somewhat strong/easily/satisfying/painful; 4: Somewhat weak/difficult/unsatisfying/painful; 5: Very weak/difficult/unsatisfying/painful; 6: No sex drive/never aroused/never/never reach orgasm/Can't reach orgasm/Extremely painful.<sup>a</sup>

Based on McGahuey et al. (2000).