

## Accepted Manuscript

Title: Effects of sex and menstrual cycle phase on cardiac response and alpha- amylase levels in psychosocial stress

Authors: Laura Espin, Carolina Villada, Vanesa Hidalgo, Alicia Salvador



PII: S0301-0511(18)30224-2  
DOI: <https://doi.org/10.1016/j.biopsycho.2018.12.002>  
Reference: BIOPSY 7627

To appear in:

Received date: 7 April 2018  
Revised date: 30 October 2018  
Accepted date: 4 December 2018

Please cite this article as: Espin L, Villada C, Hidalgo V, Salvador A, Effects of sex and menstrual cycle phase on cardiac response and alpha- amylase levels in psychosocial stress, *Biological Psychology* (2018), <https://doi.org/10.1016/j.biopsycho.2018.12.002>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Effects of sex and menstrual cycle phase on cardiac response and alpha- amylase levels  
in psychosocial stress.**

Laura Espin<sup>1,2\*</sup>, Carolina Villada<sup>2,3</sup>, Vanesa Hidalgo<sup>2,4</sup>, Alicia Salvador<sup>2</sup>

<sup>1</sup> Department of Human Anatomy and Psychobiology, University of Murcia, 30100 Murcia, Spain

<sup>2</sup> Laboratory of Social Cognitive Neuroscience, IDOCAL, Department of Psychobiology, University of Valencia, 46010 Valencia, Spain

<sup>3</sup> Department of Cognitive and Behavioral Neurobiology. Institute of Neurobiology. National Autonomous University of Mexico, Juriquilla, Querétaro, Mexico.

<sup>4</sup> IIS Aragón, Department of Psychology and Sociology, Area of Psychobiology, University of Zaragoza, Teruel, Spain

\* Corresponding Author: Laura Espin, Department of Human Anatomy and Psychobiology, University of Murcia, 30100 Murcia, Spain. Tel.: +34 868 888 665; Fax: +34 868 884 111.

E-mail address: lespin@um.es (L. Espin)

**Highlights**

- Follicular women exhibited blunted alpha-amylase reactivity to the TSST
- Luteal women and men showed higher alpha-amylase reactivity to the TSST
- Men showed higher LF/HF response to the TSST compared to the control condition

- No condition differences were found in the LF/HF responses of the two groups of women
- Sex and menstrual cycle factors should be considered in the stress response

### ABSTRACT

The impact of sex and the menstrual cycle phase on the autonomic response to psychosocial stress remains controversial. This study explored autonomic nervous system activity through salivary alpha-amylase, heart rate, and heart rate variability responses to the Trier Social Stress Test (TSST) in healthy young people. The sample was composed of 25 men, 26 women in the luteal phase, and 25 women in the follicular phase, from 18 to 25 years of age. Participants were exposed to the TSST or a control condition. The results indicate that women in their follicular phase showed a blunted alpha-amylase response to stress compared to men and women in the luteal phase. In addition, men showed higher sympatho-vagal activity in the stress condition compared to the two groups of women. These results confirm that sex and the menstrual cycle phase are potential modulators of autonomic nervous system reactivity to psychosocial stress.

#### **Keywords:**

Salivary alpha-amylase; psychosocial stress; sex differences; menstrual cycle; heart rate variability.

## 1. INTRODUCTION

Many studies have shown that men and women differ in the way they respond to stressors, based on their endocrinological and behavioral responses (Taylor et al., 2000), and this difference seems to be reflected in the sex-specific prevalence rates for several diseases (Gobinath et al., 2017). Thus, women suffer more often from autoimmune illnesses associated with anxiety and depression, whereas men are more susceptible to developing cardiovascular disease (Kajantie & Phillips, 2006). A stressful stimulus results in the activation of several physiological pathways, including the autonomic nervous system (ANS) and the hypothalamic–pituitary–adrenal axis (HPA), which seem to be affected by sex. Therefore, studying sex differences in stress reactivity could help us to find a mechanism that explains why some diseases are more common in men and others in women.

A large amount of research has been published on sex-differences in ANS and HPA-axis responses to stress. Differences have been found between men and women on several baseline or pre-stress cardiovascular measures, such as heart rate (HR) and blood pressure (Suarez et al., 2004), as well as differences in the effects of sex and hormonal status on the physiological response to acute psychosocial stress (Stark et al., 2006; Wolf et al., 2001). In addition, a higher salivary cortisol response has been found in men than in women in reaction to social stress (Hidalgo et al., 2014; Kirschbaum et al., 1999; Stephens et al., 2016; Uhart et al., 2006). The role of sex hormones is an issue that needs to be considered to better explain why some studies show no sex differences in stress responsiveness. For example, the number of cortisol responders to social stress decreases during the follicular phase and in postmenopausal women, compared to women in the luteal phase (Villada et al., 2017), possibly because women in the luteal phase show higher levels of progesterone and estrogens than in other phases of the menstrual cycle or even during the postmenopausal period (Gordon &

Girdler, 2014).

Some authors have also suggested that estrogens have a diminishing effect on sympathetic activity, but the effect of progesterone is still unclear (Kajantie & Phillips, 2006). Moreover, the circulating levels of allopregnenolone (a neuroactive metabolite of progesterone) seem to be involved in the sensitivity to stress. Specifically, the larger allopregnenolone levels after stress induction (aversive movies) have been related to a higher HR response, greater negative affect, and lower phasic activity in the amygdala and medial Prefrontal Cortex (mPFC) in women in the luteal phase than in women in the follicular phase (Ossewarde et al., 2010).

Despite the relevance of sex differences in the autonomic stress response and its interaction with stress hormones, only a few studies have focused on sex differences in salivary alpha-amylase (sAA) levels or responses, with ambiguous results (Kivlighan & Granger, 2006; Nater et al., 2006; Takai et al., 2007). sAA has been proposed as an indirect marker of the sympatho-adrenal medullar (SAM) response to different stressors (Filaire et al., 2010; Nater & Rohleder, 2009). In recent years, sAA activity has received considerable attention (Nater et al., 2005, 2006; Rohleder & Nater, 2009; van Stegeren, Wolf, & Kindt, 2008), and its response profile to stress has been studied in several age groups, including children (Räikkönen et al., 2010; Spinrad et al., 2009), adolescents (Sumter, Bokhorst, Miers, Van Pelt, & Westenberg, 2010; Susman et al., 2010), young people (Nater et al., 2005, 2006; Rohleder, Wolf, Maldonado, & Kirschbaum, 2006; Schoofs, Hartmann, & Wolf, 2008) and older adults (Almela et al., 2011; Strahler, Mueller, Rosenlocher, Kirschbaum, & Rohleder, 2010). Discussion about whether sAA is a good index for sympathetic nervous system (SNS) activity is still ongoing because the available literature is inconclusive. Positive associations between stress-induced sAA responses and markers of SNS activity have been observed, for example, in plasma concentrations of norepinephrine (Thoma, Kirschbaum, Wolf, &

Rohleder, 2012) and in measures of the sympatho-vagal balance, with increases in heart rate (HR) (Nater et al., 2006) and decreases in the cardiac parasympathetic activity (Bosch, Geus, Veerman, Hoogstraten, & Nieuw Amerongen, 2003). However, other studies have reported no associations between stress-induced sAA levels and plasma noradrenaline (Petrakova et al., 2015) or cardiovascular measures (Strahler et al., 2010).

Another currently debated aspect is whether there are sex differences in the sAA response to acute stress. Although no sex differences in basal sAA levels were found in some studies (Nater, Rohleder, Schlotz, Ehlert, & Kirschbaum, 2007), others reported greater noradrenergic arousal (higher sAA at baseline) in men compared to women (Carr, Scully, Webb & Felmingham, 2016). Several studies have reported that men and women have a comparable sAA responsivity to standardized stressors such as the cold pressor stress (CPS) test (van Stegeren et al., 2008) or the Trier Social Stress Test (TSST) (Almela et al., 2011; Hidalgo et al., 2012; Hlavacova et al., 2017; Maruyama et al., 2012; Schoofs & Wolf, 2011; Thoma et al., 2012). However, Smeets (2010) reported sex differences in the acute sAA response to the TSST, with a higher sAA response in men than in women (women were tested only in the luteal phase). More recently, Carr et al. (2016) found higher sAA reactivity to stress in young women (tested in several phases of the menstrual cycle) than in men, in response to the CPS test. Therefore, in light of these mixed findings, it seems necessary to take into account the women's menstrual cycle phase when studying the noradrenergic response to acute stress.

The impact of sex on other biomarkers of the stress response, such as HR or heart rate variability (HRV), also remains controversial. Several studies have reported no sex differences in HR responses to the TSST (Kirschbaum et al., 1999; Kelly, Tyrka, Anderson, Price, & Carpenter et al., 2008; Villada, Hidalgo, Almela & Salvador, 2018), whereas other studies have found an enhanced HR response in women (Kudielka, Buske-Kirschbaum,

Hellhammer, & Kirschbaum, 2004; Smeets, 2010). Results from studies on the effects of sex hormones on HR responses are unclear. Some studies on cardiac variability across phases of the menstrual cycle suggest that estrogen may attenuate HR reactivity (Kajantie & Phillips, 2006). However, years later, one study showed that women tested in the luteal phase exhibited a greater HR response to the TSST than men, but not greater than women in the follicular phase (Childs et al., 2010). Moreover, no differences have been found in the HR and HRV parameters in response to social stress in young women in different phases of the menstrual cycle (Gordon & Girdler, 2014; Pico-Alfonso et al., 2007; Villada et al., 2014).

HRV, a measurement of the beat-to-beat difference in HR, is mediated by the parasympathetic and sympathetic cardiac nerves and reflects the capacity for the parasympathetic inhibition of autonomic arousal. There is evidence indicating that this balance may be affected by various psychiatric conditions, as well as by the presence of stressful stimuli (Berntson and Cacioppo, 2004), showing an increased response of the sympathetic component and a decrease in the parasympathetic response during the stress condition (Kim et al., 2018; Nater et al., 2006). Regarding sex differences in HRV, a recent meta-analysis showed that, in general, HRV in women is characterized by a relative dominance of vagal, parasympathetic activity, despite a higher mean HR (Koenig and Thayer et al., 2016). In addition, some studies report that HRV displays physiological changes throughout the menstrual cycle (Brar et al., 2015; Uckuyu et al., 2013; Vishrutha et al., 2012). Vishrutha et al. (2012) concluded that the HF component of HRV was higher in the follicular phase, whereas the LF component was higher in the luteal phase. These results suggest a parasympathetic predominance during the follicular phase and greater sympathetic activity in the luteal phase (Brar et al., 2015; Vishrutha et al., 2012), although other studies did not find differences in HRV related to the menstrual cycle phase (Das et al., 2015; Leich et al., 2003; Nakagawa et al., 2006). Thus, this relationship between the menstrual cycle and

autonomic activity is still an unresolved question (Das et al., 2015).

Due to a relative lack of knowledge and consistency regarding the role of sex and the menstrual cycle phase in ANS markers of the stress response, the current study aims to clarify the sex and menstrual cycle phase differences in the ANS response to stress. To address this response, we integrated the sAA response with cardiac markers (HR and HRV) in a comprehensive range of complementary indicators of ANS activity. Because personality traits, specifically trait anxiety, can play an important role in the psychophysiological stress response (Chida & Hamer, 2008), in order to control individual differences, a measure of trait anxiety was included. Based on previous literature, we expect a sympathetic activation in response to the TSST, compared to the control condition, measured through sAA, HR, and the ratio of low and high frequency HRV bands (LF/HF), and a decrease in the root mean square of successive differences (RMSSD) as a parasympathetic modulation. In relation to sex differences and the effects of the menstrual cycle phase in the response to stress, due to mixed results in the literature, we aim to explore the full profile of stress responses in women in different phases of the menstrual cycle.

## **2. METHODS**

### **2.1. Participants**

The final sample was composed of seventy-six healthy university students, 25 men (mean age = 19.56; SD = 1.80) and 51 women (mean age = 18.90; SD = 1.48). From the women's subgroup, 26 were tested in the luteal phase of the menstrual cycle (mean age = 19; SD = 1.62), and 25 were tested in the follicular phase of the menstrual cycle (mean age = 18.80; SD = 1.35).

To recruit participants, informative talks were held in various campus classrooms. Volunteers were interviewed by trained psychologists, and they completed an extensive questionnaire to check whether they met the study prerequisites. From the initial sample of 119



volunteers, 20 women were excluded because they had spontaneous anovulatory cycles, 13 subjects were not selected because of the exclusion criteria, and 10 subjects were eliminated due to a variety of problems during the experimental procedure. The exclusion criteria were: alcohol or other drug abuse, visual or hearing problems, presence of cardiovascular, endocrine, neurological, or psychiatric disease, and the presence of a stressful life event during the past year. To obtain this information participants were asked about the presence or not of any important (positive or negative) event in the past year that they considered stressful and that changed their daily life significantly: e.g. traffic accident, change of habits, winning the lottery, death of someone close). Each event was evaluated based on the impact (positive vs. negative) and based on the magnitude: from 1 (not relevant) to 4 (very relevant). Subjects were also excluded if they were using any medication directly related to emotional or cognitive function, or one that was able to influence hormonal and sAA levels, such as glucocorticoids, beta-blockers, antidepressants, benzodiazepines, asthma medication, thyroid therapies, and psychotropic substances. Vitamins, sporadic use of painkillers, and natural therapies were allowed. Because there is no conclusive data available on the impact of the use of oral contraceptives on sAA (Rohleder & Nater, 2009), women using oral contraceptives were excluded from participation. None of the participants were habitual smokers, although in each condition, five participants reported sporadic smoking (less than 10 cigarettes a week).

Subjects in the final sample were randomly assigned to one of two conditions: the TSST procedure versus a control condition. The TSST group was made up of 13 men, 13 women in the luteal phase (4th to 8th day before the onset of the new menstrual cycle), and 11 women in the follicular phase (5th to 8th day after the onset of the new menstrual cycle). The subjects in the control condition were 12 men, 13 women in the luteal phase, and 14 women in the follicular phase.

The menstrual cycle phase was calculated using two estimation procedures. First, in order to establish the date of each subject's appointment, all the cycles were converted to a standard 28-day cycle, taking as reference points the day of onset of the last menstruation and the real length of the studied cycle (Rossi & Rossi, 1980). Second, to confirm the previous estimation and estimate the ovulation point, Basal Body Temperature (BBT) was recorded daily during two complete menstrual cycles by means of sublingual temperature, taken for 5 min before getting up. To analyze the BBT, the “smoothed curve” (SMC) method was used, as described by McCarthy and Rockette (1983, 1986).

Participants who met the criteria were contacted by telephone and asked to attend a session that took place in a laboratory at the Faculty of Psychology. Before each session, participants were asked to maintain their general habits, sleep as long as usual, refrain from heavy physical activity the day before the session, and not consume alcohol since the night before the session.

Additionally, they were instructed not to brush their teeth at least 1 h prior to the session, and to drink only water, without eating, smoking, or taking any stimulants, such as coffee, cola, caffeine, tea or chocolate, 2 h prior to the session, because these factors have been shown to modify sAA activity (Rohleder & Nater, 2009). The study was conducted in accordance with the Declaration of Helsinki, and the protocol was approved by the Ethics Research Committee of the University of Murcia. All the participants received verbal and written information about the study and signed an informed consent.

## **2.2.Procedure**

Experimental sessions were held in the laboratory at the university between 2pm and 5pm (the sequence is presented schematically in Figure 1). This study employed a between-subjects design where participants were tested individually and in a single session. After arriving at the laboratory, participants' weight and height were measured, and the HR

measurement device was connected and activated. Experimental sessions consisted of several phases of equal durations in both conditions. Each session took approximately 1 h to complete. In the last part of the session, all the participants completed the Inventory of Situations and Responses of Anxiety (ISRA).

[INSERT FIGURE 1 HERE]

### **2.2.1. Stress condition**

The protocol started with a habituation phase of 15 min to allow the participants to adapt to the laboratory setting. During this phase, the participants remained seated, and baseline measures were obtained for HR and HRV, salivary alpha-amylase, anxiety and mood. After the habituation phase, the introduction phase started. In this phase, the participants were informed about the procedure for the stress task. They received the instructions in front of an audience in the same room where the task took place. Next, the participants returned to the first room and started the preparation for the TSST (10 min duration).

The TSST was performed according to the description provided by Kirschbaum et al. (1993). This test consists of a 10-min preparation phase that includes instructions for the speech, 5 min of free speech (a simulated job interview), and a 5 min mental arithmetic task (serial subtractions as quickly and accurately as possible) in front of a committee composed of a man and a woman.

Once it was over, the participants returned to the first room, and they had 20 min to recover while they answered the trait anxiety questionnaire. Then, the HR and HRV recorders were turned off.

### **2.2.2. Control condition**

The control condition was designed to be as similar as possible to the stress condition in mental workload and overall physical activity, but without the main components capable of provoking stress, such as evaluative threat and uncontrollability (Dickerson & Kemeny, 2004). The control task was composed of a preparation phase of 10 min, where the participants read a chapter from a book with neutral content, followed by 5 minutes of reading aloud and an arithmetic task that involved counting by one for 5 minutes. The control condition was performed in the same room as the stress condition, but all the stress-producing elements were removed prior to starting it (video camera, tape recorder, committee, and microphone). The timing of the saliva samples, the questionnaires used, and the phase durations were the same for both conditions.

## **2.3. Measures**

### **2.3.1. Salivary Alpha-amylase (sAA)**

sAA concentrations were estimated from saliva samples obtained using Salivettes (Sarstedt, Nümbrecht, Germany) during a 55 min period with six assessment points:  $t-5$ ,  $t_0$ ,  $t+10$ ,  $t+15$ ,  $t+20$  and  $t+40$  min, from the start of the stressor (see Figure 1). Participants were instructed to introduce the cotton swab into their mouths for 1 min, not chew the cotton because it may affect salivary protein composition as well as the flow rate (Bosch et al., 2011), and move the swab around in a circular pattern to collect saliva from all the salivary glands (Rohleder and Nater, 2009). Saliva samples were frozen at  $-20$  °C from the end of the session until the analyses took place. The concentration of alpha-amylase in saliva was measured by an enzyme kinetic method following the protocol specified in Rohleder, Wolf, Maldonado and Kirschbaum (2006). Inter- and intra-assay variation was below 10%.

### **2.3.2. Heart Rate (HR) and Heart Rate Variability (HRV)**

HR and HRV were measured using an HR monitor (Suunto, model T6, Suunto Oy, Vantaa, Finland), which consists of a chest belt for heart beat detection and transmission and a “watch” for data collection and storage. Heartbeat detection is performed with an accuracy of 1 ms, and these types of monitors have shown good validity (Roy, Boucher, & Comtois, 2009). After eliminating the artifacts, the HR and HRV means for each 5-min phase (habituation, preparation, speech, arithmetic, and recovery) were computed. HR was recorded in real-time and expressed as beats per minute (bpm). To evaluate HRV, we used two parameters: the root mean square of successive differences (RMSSD in ms) from time domain analysis, reflecting the short-time HRV, predominantly as a response to changes in parasympathetic tone; and the ratio between low and high frequency components (LF/HF ratio) of HRV spectra as an index of vagal balance (Task Force, 1996).

### **2.3.3. Trait Anxiety**

The Inventory of Situations and Responses of Anxiety (ISRA, Cano-Vindel & Miguel-Tobal., 1999) is a self-report questionnaire that measures, based on Lang’s model (Lang, 1968), different components (cognitive, physiological, and motor) of trait anxiety on three subscales containing 24 items each (Cronbach’s alpha ranges from  $\alpha = 0.95$  to 0.98). A total score is obtained by simply adding together the scores on the three subscales.

## **2.4. Statistical Analyses**

Data were tested for normal distribution and homogeneity of variance using the Kolmogorov–Smirnov and Shapiro–Wilk tests before the statistical procedures were applied. These analyses revealed significant deviations from normality on all the absolute sAA levels. Therefore, they were log-transformed prior to analyses, and all subsequent analyses were conducted with log-transformed data.

One-way ANOVAs were conducted to evaluate possible differences between groups on demographic and anthropometric variables and trait anxiety scores (ISRA). ANOVAs for repeated measures were used to analyze sAA levels and HR and HRV parameters (RMSSD, LF/HF), with group (luteal vs. follicular vs. men) and condition (TSST vs. control) as between-subjects factors. We employed time as within-subjects factor (For sAA: t-5, t0, t+10, t+15, t+20 and t+40 min; for HR- HRV parameters: habituation, preparation, speech, arithmetic task and recovery). Furthermore, the area under the curve with respect to the increase (AUC<sub>i</sub>) was calculated using the trapezoid formula specified in Pruessner, Kirschbaum, Meinlschmid, and Hellhammer (2003) for sAA, HR, RMSSD, and LF/HF. Univariate ANOVAs were computed to compare the two conditions and groups on all the physiological indices.

All the results were corrected using the Greenhouse-Geisser procedure, where appropriate. Post-hoc comparisons were performed using Bonferroni adjustments for multiple comparisons. As a measure of the effect size, we report Partial Eta Squared ( $\eta^2_p$ ) values.

### **3. RESULTS**

#### **3.1. Demographic and anthropometric variables**

The main characteristics of the groups are shown in Table 1. Significant differences were found only on BMI, with men showing higher BMI than both groups of women. All the groups showed similar scores on the ISRA total and each of its subscales, with no significant differences.

[INSERT TABLE 1 HERE]

#### **3.2. Biochemical and cardiovascular measures**

Significant effects of the time x condition interaction for sAA ( $F(5,350) = 2.79; p = 0.01; \eta^2_p = 0.03$ ), HR ( $F(4,280) = 14.37; p < 0.001; \eta^2_p = 0.17$ ) and RMSSD of HRV ( $F(4, 280) = 2.976; p = 0.02; \eta^2_p = 0.04$ ) were found. The participants exposed to the TSST condition

showed higher sAA levels than those exposed to the control condition at all the sampling times ( $p \leq 0.025$ ), except at baseline (t-5) and t+0 (for both  $p > 0.05$ ). For HR, significant differences between the two conditions were found in the preparation ( $p < 0.001$ ), speech ( $p = 0.001$ ), and arithmetic task ( $p = 0.004$ ), with higher HR in the TSST condition than in the control condition. For RMSSD, there were no significant differences between conditions in the baseline and recovery periods ( $p > 0.10$  in both cases), but a significant difference between conditions was found during the recovery period ( $p = 0.03$ ) with lower RMSSD levels found in the stress condition than in the control condition.

The time x group interaction was significant for LF/HF ( $F(8, 280) = 4.583$ ;  $p \leq 0.001$ ;  $\eta^2_p = 0.11$ ), as men showed higher LF/HF than luteal women on the arithmetic task ( $p \leq 0.001$ ). No differences were found between men and follicular women ( $p = 0.12$ ) or between the two groups of women ( $p = 0.10$ ). In addition, results revealed significant increases from preparation to the speech in follicular women ( $p = 0.04$ ), whereas no differences were found between these phases in luteal women (all  $p > 0.10$ ). Men showed significant increases from habituation to speech ( $p = 0.004$ ), habituation to arithmetic ( $p \leq 0.001$ ), speech to recovery ( $p = 0.002$ ), and arithmetic to recovery ( $p \leq 0.001$ ).

The condition x group interaction was significant only for LF/HF ( $F(2, 70) = 4.270$ ;  $p = 0.018$ ;  $\eta^2_p = 0.10$ ). In the TSST condition, men showed a higher LF/HF ratio than the two groups of women (both  $p < 0.02$ ). There were no significant differences between the two groups of women ( $p > 0.99$ ) (see Figure 2).

**[INSERT FIGURE 2 HERE]**

In addition, only men showed higher LF/HF levels in the stress condition than in the control condition ( $p \leq 0.001$ ). In the control condition, no significant differences between groups were found (all  $p > 0.10$ ) (see Figure 3).

[INSERT FIGURE 3 HERE]

With regard to the AUC<sub>i</sub> for the sAA, the condition x group interaction was significant ( $F(2, 70) = 4.82; p = 0.01; \eta^2_p = 0.12$ ). In the TSST condition, men and luteal women showed higher sAA output than their counterparts in the control condition ( $p=0.03$  and  $p=0.009$ , respectively). No condition differences were found in follicular women ( $p=0.16$ ). In the control condition, no significant differences between groups were found (all  $p > 0.99$ ), whereas in the TSST condition, follicular women showed a significantly less pronounced stress-induced sAA output, compared to men ( $p=0.03$ ) and luteal women ( $p=0.002$ ). There were no significant differences between luteal women and men ( $p=0.87$ ) (see Figure 4).

[INSERT FIGURE 4 HERE]

For the AUC<sub>i</sub> for the HR, a significant effect of the condition was found, showing higher cardiac reactivity in the TSST condition than in the control condition ( $F(1, 70) = 12.68; p = 0.001; \eta^2_p = 0.15$ ), but there were no other significant differences (all  $p > 0.10$ ).

#### 4. DISCUSSION

Our aim was to study the effects of sex and the menstrual cycle phase on sAA and cardiac responses to psychosocial stress. As expected, the participants exposed to the TSST showed significant increases in sAA and HR (sympathetic response) and a decrease in the RMSSD (parasympathetic response). Thus, our results confirm previous findings showing an



increase in sAA (Nater et al., 2005, 2006) and HR (Nater et al., 2005) and lower parasympathetic activity in response to the TSST (Villada et al., 2018).

A ~~negative~~ less pronounced sAA reactivity to the TSST in follicular women was found, compared to men and luteal women, but no differences were found between men and luteal women. In addition, the LF/HF response to the TSST increased only in men, compared to both groups of women in the TSST condition and men in the control condition.

The sAA response was different between groups in the experimental condition. Women in the follicular phase showed a less pronounced sAA reactivity (AUC<sub>i</sub>) to the TSST compared to men and women in the luteal phase. Physiological differences in estrogen levels probably contribute to the differences observed in the stress responsiveness of women in the luteal phase because estrogen concentrations are high during this phase (Kajantie & Phillips, 2006). However, estrogen levels are also known to be high during the mid-to-late follicular phase compared to the early follicular phase (Gordon & Girdler, 2014). The luteal and follicular phases differ most in progesterone levels, with women in the luteal phase presenting the highest levels of this hormone (Gordon & Girdler, 2014; Kajantie & Phillips, 2006). The effects of progesterone on the physiological response to stress are unclear. Some studies show a sedative effect (de Wit, Schmitt, Purdy, & Hauger, 2001; Soderpalm, Lindsey, Purdy, Hauger, & de Wit, 2004), whereas others find inconsistent effects (Childs et al., 2010). In our study, luteal women (high levels of progesterone) showed an increased sAA response compared to follicular women, and so it is possible that progesterone levels have an enhancing effect on the physiological response to stress. The result found in luteal women is consistent with other study showing that progesterone increases in response to acute stress (Childs et al., 2010), and that its effects are bimodal and paradoxical depending on the released dose of this hormone and the amount of associated allopregnanolone. This metabolite of progesterone has biphasic effects, with low doses increasing an adverse anxiogenic effect,

and high doses decreasing this effect and having more calming properties. The exact mechanism of this phenomenon is not known, but it is often referred to as a biphasic or bimodal effect (Andreen et al., 2009). Along these lines, one study found an association between greater stress-induced increases in plasma allopregnanolone in luteal women compared to follicular women, directly linking the activation of the sympathoadrenal system to greater availability of progesterone, a precursor of allopregnanolone in luteal women (Girdler et al., 2006). These higher levels of allopregnanolone during the luteal phase has also been related to higher sensitivity to stress, measured through HR and noradrenaline, and lower phasic activity in the amygdala and in the mPFC (Ossewarde et al., 2010). Findings from another study showed that women in the follicular phase exhibited blunted noradrenaline responses to the TSST (Childs et al., 2010). However, a recent study found a higher sAA response to social stress in women in the follicular phase compared to women in the luteal phase (Hlavacova et al., 2017).

In our study, considering the sAA response as an indirect indicator of catecholaminergic activity, we suggest that the blunted sAA reactivity to the TSST found in follicular women could be explained by the blunted catecholaminergic activity in this group. These results provide new information about the effects of the menstrual cycle on sAA reactivity to stress because most of the studies have only included men and women in the luteal phase (Smeets, 2010; Strahler et al., 2010; Thoma et al., 2012), only women who use oral contraceptives (Almela et al., 2011), or only women in the early follicular phase (Hidalgo et al., 2014).

The results highlight the blunted sAA reactivity to stress found in women in the follicular phase, compared to men and women in the luteal phase. These cycle-related differences may provide clues about sex differences in the prevalence of stress-related disorders. For example, blunted sAA reactivity was recently associated with atopic diseases

(i.e. allergic rhinitis, asthma) in adult men and women (Hlavacova et al., 2017), and in children with asthma between 8-18 years old (Wolf et al., 2008). Thus, a repeated blunted adrenocortical response to acute stress in women could contribute to heightened susceptibility to some autoimmune diseases. In this regard, the low reactivity associated with the menstrual cycle phase could influence the manifestation and treatment of some disorders associated with women. In women, an increase in anxiety symptoms has been found to coincide with the peak in gonadal hormones (Kessler et al., 2005), and sex differences have been observed in depression (Ge et al., 2001). Therefore, this aspect should be taken into account in clinical practice because these hormonal changes during the menstrual cycle could correspond to changes in the presentation of psychiatric illnesses, thus affecting treatment success. Moreover, women in the follicular phase seem to show more psychological complaints than women in other phases of the menstrual cycle (Guillermo et al., 2010). Personality traits and stress coping styles are factors that could be measured in future studies. Understanding how the mechanisms involved in the stress response interact with each other will have important benefits for future interventions and help patients to improve their strategies for coping with challenging life events better than classical pharmacological approaches. From these results, some questions arise to investigate in future studies. The underlying mechanisms through which progesterone may moderate sex differences require further study. In addition, the impact of other sex-related hormones or factors that we did not measure should be considered, such as estrogen and progesterone levels to confirm the menstrual cycle phase evaluated, as well as allopregnone.

Furthermore, men exhibited a higher LF/HF response to the TSST than the two groups of women, whereas the HR and RMSSD response to the TSST did not differ according to sex or menstrual cycle. These data suggest that sympathetic nervous system activity in response to the TSST appears to be more dominant in men than in women (both luteal and follicular).

These results partly match other studies using the TSST that did not find sex differences in the HR response (Kelly et al., 2008; Kirschbaum et al., 1999; Smeets, 2010), and they contrast with results obtained by other authors (Lustyk, Olson, Gerrish, Holder, & Widman, 2010) who found that women in the luteal phase were significantly more HR reactive than women in the follicular phase. However, these latest results did not consider men and, therefore, are not completely comparable to ours. Previous studies have shown that, in young adults, there is a predominance of sympathetic regulation in men and a dominant parasympathetic influence in women (Evans et al., 2001). Furthermore, stress tasks in women have generally reported a higher HF component, whereas men exhibited a higher LF/HF and LF component (Sato & Miyake, 2004). The reason for these differences is not clear. One could speculate that sex differences could be due to the type of stressor used, but Ordaz and Luna (2012) stated that any sex differences reflect the sex-specific sensitivity of the physiological domain rather than the type of stressor that elicited the response. Along these lines, one possible explanation for our findings would be that female reproductive hormones such as estrogen or progesterone might inhibit sympathetic activation. In some settings, estrogen has had an attenuating effect on sympathetic activity (Ryan et al., 1994), and during the luteal phase, higher estradiol concentrations were associated with lower cardiac output responses to a speech task (Sita & Miller, 1996). In addition, some differences have been found between women in the late follicular and early follicular phases, with higher HR in the late follicular group characterized by higher levels of estradiol (Pollard et al., 2007).

A limitation of this study could be that the design used was between-subjects instead of intra-subject. Despite criticism of the between-subjects design, we think this design could be one of the strong points of the present study. The inclusion of a control condition and strictly random assignment of the subjects to the experimental and control groups guaranteed that both groups were equivalent in the important dimensions. Therefore, we can assume that

the differences found in the two groups were solely due to the experimental manipulation, thus guaranteeing the internal validity of the study. Another limitation is the fact that the sex hormones were not measured. Although we used a rigorous method to estimate the menstrual cycle phase and check the ovulatory phase, further studies should include the concentrations of the women's sex hormones in each menstrual cycle phase.

Importantly, despite these possible limitations, the current study addresses several gaps in the literature on the menstrual cycle and stress reactivity. The simultaneous measurement of both autonomic and neuroendocrine reactivity in the same individuals advanced our current understanding of the autonomic mechanisms contributing to stress reactivity in women across the reproductive cycle. In addition, we confirmed that the sAA response to acute stress is sensitive to hormonal variations associated with the menstrual cycle phases evaluated. It is important to note that the stressful task used, the TSST, was able to induce a significant stress response. Therefore, we can confirm that it is a valid method to make possible sex differences emerge in the autonomic reactivity to acute stress in healthy young men and women. In addition, the TSST also increased our understanding of how women's hormonal status can modulate the autonomic reactivity to acute stress, which might be useful in future research.

Overall, these findings make an important contribution to the literature by clarifying the impact of sex and the menstrual cycle phase on ANS indicators of psychosocial stress reactivity. Moreover, they support the importance of understanding the underlying mechanisms that influence this association. Because men and women tend to react differently to stress, psychologically and biologically, these reactivity differences need to be considered in order to better understand some gender-related disorders and how they can be prevented and/or treated.

### **Role of the funding source**

This research study was supported by the Spanish Education and Science Ministry with grants no. PSI2013-46889, PROMETEOII/2015/020 and Government of Aragón, Department of Innovation, Research and University and FEDER “Construyendo desde Aragón”.

### **Conflict of interest**

The authors report no conflicts of interest. The authors alone are responsible for the content and writing of the paper.

### **Acknowledgements**

The authors wish to thank Prof. Dr. Clemens Kirschbaum for his collaboration on the biochemical analyses of sAA and Ms. Cindy DePoy for the revision of the English text.

### **References**

- Almela, M., Hidalgo, V., Villada, C., van der Meij, L., Espín, L., Gómez-Amor, J., & Salvador, A. (2011). Salivary alpha-amylase response to acute psychosocial stress: the impact of age. *Biological psychology*, 87(3), 421-429.
- Andreen, L., Nyberg, S., Turkmen, S., van Wingen, G., Fernandez, G., & Backstrom, T. (2009). Sex steroid induced negative mood may be explained by the paradoxical effect mediated by GABAA modulators. *Psychoneuroendocrinology*, 34, 1121–1132.
- befriend, not fight-or-flight. *Psychol Rev* 107(3):411–429.

- Berntson, G. G., & Cacioppo, J. T. (2004). Heart rate variability: Stress and psychiatric conditions. *Dynamic electrocardiography*, 57-64.
- Bosch, J.A., de Geus, E.J., Veerman, E.C., Hoogstraten, J., & Nieuw Amerongen, A.V. (2003). Innate secretory immunity in response to laboratory stressors that evoke distinct patterns of cardiac autonomic activity. *Psychosomatic Medicine*, 65, 245–258.
- Bosch, J.A., Veerman, E.C., de Geus, E.J., Proctor, G.B. (2011). Alpha-amylase as a reliable and convenient measure of sympathetic activity: don't start salivating just yet! *Psychoneuroendocrinology* 36, 449–453.
- Brar, T. K., Singh, K. D., & Kumar, A. (2015). Effect of different phases of menstrual cycle on heart rate variability (HRV). *Journal of clinical and diagnostic research: JCDR*, 9(10), CC01.
- Cano-Vindel, A., & Miguel-Tobal, J. J. (1999). Evaluación de la ansiedad desde un enfoque interactivo y multidimensional: El inventario de Situaciones y Respuestas de Ansiedad (ISRA). *Psicología Contemporánea*, 6(1), 14-21.
- Carr, A. R., Scully, A., Webb, M., & Felmingham, K. L. (2016). Gender differences in salivary alpha-amylase and attentional bias towards negative facial expressions following acute stress induction. *Cognition and Emotion*, 30(2), 315-324.
- Chida, Y., & Hamer, M. (2008). Chronic psychosocial factors and acute physiological responses to laboratory-induced stress in healthy populations: a quantitative review of 30 years of investigations. *Psychological bulletin*, 134(6), 829.

- Childs, E., Dlugos, A., & De Wit, H. (2010). Cardiovascular, hormonal, and emotional responses to the TSST in relation to sex and menstrual cycle phase. *Psychophysiology*, 47, 550-559.
- Das, A., Chakraborti, D., Saha-Roy, S., Bal, C., & Chatterjee, S. (2015). Effect Of Endogenous Female Sex Hormone Fluctuations During Menstrual Cycle On Heart Rate Variability. *IOSR Journal of Dental and Medical Sciences*, 14(7), pp 01-05.
- de Wit, H., Schmitt, L., Purdy, R., & Hauger, R. (2001). Effects of acute progesterone administration in healthy postmenopausal women and normally-cycling women. *Psychoneuroendocrinology*, 26, 697-710.
- Dickerson, S. S., & Kemeny, M. E. (2004). Acute stressors and cortisol responses: a theoretical integration and synthesis of laboratory research. *Psychological bulletin*, 130(3), 355-391.
- Evans, J.M., Ziegler, M.G., Patwardhan, A.R., Ott, J.B., Kim, C.S., Leonelli, F.M., & Knapp, C.F. (2001). Gender differences in autonomic cardiovascular regulation: spectral, hormonal, and hemodynamic indexes. *Journal of Applied Physiology*, 91, 2611-2618.
- Filaire, E., Portier, H., Massart, A., Ramat, L., & Teixeira, A. (2010). Effect of lecturing to 200 students on heart rate variability and alpha-amylase activity. *European journal of applied physiology*, 108(5), 1035-1043.
- Ge, X., Conger, R. D., & Elder Jr, G. H. (2001). Pubertal transition, stressful life events, and the emergence of gender differences in adolescent depressive symptoms. *Developmental psychology*, 37(3), 404-417.



- Girdler, S. S., Beth Mechlin, M., Light, K. C., & Morrow, L. A. (2006). Ethnic differences in allopregnanolone concentrations in women during rest and following mental stress. *Psychophysiology*, 43, 331–336.
- Gobinath, A. R., Choleris, E., & Galea, L. A. (2017). Sex, hormones, and genotype interact to influence psychiatric disease, treatment, and behavioral research. *Journal of neuroscience research*, 95(1-2), 50-64.
- Gordon, J. L., & Girdler, S. S. (2014). Mechanisms underlying hemodynamic and neuroendocrine stress reactivity at different phases of the menstrual cycle. *Psychophysiology*, 51(4), 309-318.
- Guillermo, C. J., Manlove, H. A., Gray, P. B., Zava, D. T., & Marrs, C. R. (2010). Female social and sexual interest across the menstrual cycle: the roles of pain, sleep and hormones. *BMC women's health*, 10(1), 1-10.
- Hlavacova, N., Solarikova, P., Marko, M., Brezina, I., & Jezova, D. (2017). Blunted cortisol response to psychosocial stress in atopic patients is associated with decrease in salivary alpha-amylase and aldosterone: Focus on sex and menstrual cycle phase. *Psychoneuroendocrinology*, 78, 31-38.
- Hidalgo, V., Villada, C., Almela, M., Espín, L., Gómez-Amor, J., & Salvador, A. (2012). Enhancing effects of acute psychosocial stress on priming of non-declarative memory in healthy young adults. *Stress*, 15(3), 329-338.

- Hidalgo, V., Almela, M., Villada, C., & Salvador, A. (2014). Acute stress impairs recall after interference in older people, but not in young people. *Hormones and behavior*, 65(3), 264-272.
- Kajantie, E., & Phillips, D. I. (2006). The effects of sex and hormonal status on the physiological response to acute psychosocial stress. *Psychoneuroendocrinology*, 31, 151-178.
- Kelly, M.M., Tyrka, A.R., Anderson, G.M., Price, L.H., & Carpenter, L.L. (2008). Sex differences in emotional and physiological responses to the Trier Social Stress Test. *Journal of Behavior Therapy and Experimental Psychiatry*, 39, 87-98.
- Kessler, R. C., Berglund, P., Demler, O., Jin, R., Merikangas, K. R., & Walters, E. E. (2005). Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the National Comorbidity Survey Replication. *Archives of general psychiatry*, 62(6), 593-602.
- Kim, H. G., Cheon, E. J., Bai, D. S., Lee, Y. H., & Koo, B. H. (2018). Stress and Heart Rate Variability: A Meta-Analysis and Review of the Literature. *Psychiatry investigation*, 15(3), 235.
- Kirschbaum, C., Kudielka, B.M., Gaab, J., Schommer, N.C., & Hellhammer, D.H. (1999). Impact of gender, menstrual cycle phase, and oral contraceptives on the activity of the hypothalamus-pituitary-adrenal axis. *Psychosomatic Medicine*, 61, 154-162.
- Kirschbaum, C., Pirke, K.M., & Hellhammer, D.H. (1993). The "Trier Social Stress Test" a tool for investigating psychobiological stress responses in a laboratory setting. *Neuropsychobiology*, 28, 76-81.

- Kivlighan, K. T., & Granger, D. A. (2006). Salivary  $\alpha$ -amylase response to competition: Relation to gender, previous experience, and attitudes. *Psychoneuroendocrinology*, 31(6), 703-714.
- Koenig, J., & Thayer, J. F. (2016). Sex differences in healthy human heart rate variability: a meta-analysis. *Neuroscience & Biobehavioral Reviews*, 64, 288-310.
- Kudielka, B.M., Buske-Kirschbaum, A., Hellhammer, D.H., & Kirschbaum, C. (2004). Differential heart rate reactivity and recovery after psychosocial stress (TSST) in health children, younger adults, and elderly adults: the impact of age and gender. *International Journal of Behavioral Medicine*, 11, 116–121.
- Lang, P.J. (1968). Fear reduction and fear behavior: problems in treating a construct. In J.M. Shleien (Ed.). *Research in Psychotherapy, III*. Washington: American Psychological Association.
- Leicht, A. S., Hirning, D. A., & Allen, G. D. (2003). Heart rate variability and endogenous sex hormones during the menstrual cycle in young women. *Experimental physiology*, 88(3), 441-446.
- Lustyk, M.K.B., Olson, K.C., Gerrish, W.G., Holder, A., & Widman, L. (2010). Psychophysiological and neuroendocrine response to laboratory stressors in women: implications of menstrual cycle phase and stressor type. *Biological Psychology*, 83, 84-92.
- Maruyama, Y., Kawano, A., Okamoto, S., Ando, T., Ishitobi, Y., Tanaka, Y., ... & Ninomiya, T. (2012). Differences in salivary alpha-amylase and cortisol responsiveness following exposure to electrical stimulation versus the Trier Social Stress Tests. *PLoS One*, 7(7),

e39375.

McCarthy, J.J., Rockette, H.E., 1983. A comparison of methods to interpret the basal body temperature graph. *Fertil. Steril.* 39, 640–646.

McCarthy, J.J., Rockette, H.E., 1986. Prediction of ovulation with basal body temperature. *J. Reprod. Med.* 31, 742–747.

Nakagawa, M., Ooie, T., Takahashi, N., Taniguchi, Y., Anan, F., Yonemochi, H., & Saikawa, T. (2006). Influence of menstrual cycle on QT interval dynamics. *Pacing and Clinical Electrophysiology*, 29(6), 607-613.

Nater, U. M., & Rohleder, N. (2009). Salivary alpha-amylase as a non-invasive biomarker for the sympathetic nervous system: current state of research. *Psychoneuroendocrinology*, 34(4), 486-496.

Nater, U. M., La Marca, R., Florin, L., Moses, A., Langhans, W., Koller, M. M., & Ehlert, U. (2006). Stress-induced changes in human salivary alpha-amylase activity-associations with adrenergic activity. *Psychoneuroendocrinology*, 31, 49–58.

Nater, U. M., Rohleder, N., Gaab, J., Berger, S., Jud, A., Kirschbaum, C., & Ehlert, U. (2005). Human salivary alpha-amylase reactivity in a psychosocial stress paradigm. *International Journal of Psychophysiology*, 55, 333–342.

Nater, U.M., Rohleder, N., Schlotz, W., Ehlert, U., & Kirschbaum, C. (2007). Determinants of the diurnal course of salivary alpha-amylase. *Psychoneuroendocrinology*, 32, 392–401.

Olson, K.C. (2006). Relationship among psychological and physiological stress reactivity:

- assessment of menstrual cycle phase and autonomic perception effects. Doctoral Dissertation. Seattle Pacific University.
- Ordaz, S., & Luna, B. (2012). Sex differences in physiological reactivity to acute psychosocial stress in adolescence. *Psychoneuroendocrinology*, 37, 1135-1157.
- Ossewaarde, L., Hermans, E. J., van Wingen, G. A., Kooijman, S. C., Johansson, I. M., Bäckström, T., & Fernández, G. (2010). Neural mechanisms underlying changes in stress-sensitivity across the menstrual cycle. *Psychoneuroendocrinology*, 35(1), 47-55.
- Petrakova, L., Doering, B. K., Vits, S., Engler, H., Rief, W., Schedlowski, M., & Grigoleit, J. S. (2015). Psychosocial stress increases salivary alpha-amylase activity independently from plasma noradrenaline levels. *PloS one*, 10(8), e0134561.
- Pico-Alfonso, M. A., Mastorci, F., Ceresini, G., Ceda, G. P., Manghi, M., Pino, O., ... & Sgoifo, A. (2007). Acute psychosocial challenge and cardiac autonomic response in women: The role of estrogens, corticosteroids, and behavioral coping styles. *Psychoneuroendocrinology*, 32(5), 451-463.
- Pollard, T. M., Pearce, K. L., Rousham, E. K., & Schwartz, J. E. (2007). Do blood pressure and heart rate responses to perceived stress vary according to endogenous estrogen level in women?. *American Journal of Physical Anthropology: The Official Publication of the American Association of Physical Anthropologists*, 132 (1), 151–157.
- Pruessner, J. C., Kirschbaum, C., Meinlschmid, G., & Hellhammer, D. H. (2003). Two formulas for computation of the area under the curve represent measures of total

- hormone concentration versus time-dependent change. *Psychoneuroendocrinology*, 28(7), 916-931.
- Räikkönen, K., Matthews, K.A., Pesonen, A.K., Pyhälä, R., Paavonen, E.J., Feldt, K., et al. (2010). Poor sleep and altered hypothalamic–pituitary–adrenocortical and sympatho-adrenal-medullary system activity in children. *Journal of Clinical Endocrinology and Metabolism*, 95, 2254–2261.
- Rohleder, N., & Nater, U. M. (2009). Salivary alpha-amylase in humans. Determinants and methodological considerations. *Psychoneuroendocrinology*, 34, 469–485.
- Rohleder, N., Wolf, J.M., Maldonado, E.F., & Kirschbaum, C. (2006). The psychosocial stress-induced increase in salivary alpha-amylase is independent of saliva flow rate. *Psychophysiology*, 43, 645–652.
- Rossi, A.S., Rossi, P.E., 1980. Body time and social time: mood patterns by menstrual cycle phase and day of week. In: Parsons, J. (Ed.), *The psychology of sex differences and sex roles*. Hemisphere, New York, pp. 269–303.
- Roy, R.A., Boucher, J.P., & Comtois, A.S. (2009). Heart rate variability modulation after manipulation in pain-free patients vs patients in pain. *Journal of Manipulative and Physiological Therapeutics*, 32, 277–286.
- Ryan SM, Goldberger AL, Pincus SM, Mietus J, & Lipsitz LA. (1994). Gender and age-related differences in heart rate dynamics: Are women more complex than man? *Journal of the American College of Cardiology*, 24, 1700–1707
- Sato, N., & Miyake, S. (2004). Cardiovascular reactivity to mental stress: Relationship with menstrual cycle and gender. *Journal of Physiological Anthropology and Applied*

- Human Science, 23, 215–223.
- Schoofs, D., & Wolf, O.T. (2011). Are salivary gonadal steroid concentrations influenced by acute psychosocial stress? A study using the Trier Social Stress Test (TSST). *International Journal of Psychophysiology*, 80, 36–43.
- Schoofs, D., Hartmann, R., & Wolf, O.T. (2008). Neuroendocrine stress responses to an oral academic examination: no strong influence of sex, repeated participation and personality traits. *Stress*, 11, 52–61.
- Sita, A., & Miller, S. B. (1996). Estradiol, progesterone and cardiovascular response to stress. *Psychoneuroendocrinology*, 21, 339–346.
- Smeets, T. (2010). Autonomic and hypothalamic-pituitary-adrenal stress resilience: Impact of cardiac vagal tone. *Biological Psychology*, 84 (2), 290-295.
- Soderpalm, A. H., Lindsey, S., Purdy, R. H., Hauger, R., & de Wit, H. (2004). Administration of progesterone produces mild sedative-like effects in men and women. *Psychoneuroendocrinology*, 29, 339–354.
- Spinrad, T. L., Eisenberg, N., Granger, D. A., Eggum, N. D., Sallquist, J., Haugen, R. G., ... & Hofer, C. (2009). Individual differences in preschoolers' salivary cortisol and alpha-amylase reactivity: Relations to temperament and maladjustment. *Hormones and Behavior*, 56(1), 133-139.
- Stark, R., Wolf, O. T., Tabbert, K., Kagerer, S., Zimmermann, M., Kirsch, P., ... & Vaitl, D. (2006). Influence of the stress hormone cortisol on fear conditioning in humans: evidence for sex differences in the response of the prefrontal cortex. *Neuroimage*, 32(3), 1290-1298.
- Stephens, M. A. C., Mahon, P. B., McCaul, M. E., & Wand, G. S. (2016). Hypothalamic–pituitary–adrenal axis response to acute psychosocial stress: effects of biological sex

- and circulating sex hormones. *Psychoneuroendocrinology*, 66, 47-55.
- Strahler, J., Mueller, A., Rosenloecher, F., Kirschbaum, C., & Rohleder, N. (2010). Salivary alpha-amylase stress reactivity across different age groups. *Psychophysiology*, 47, 587–595.
- Suarez, E.C., Saab, P.G, Llabre, M.M., Kuhn, CM., Zimmerman, E. (2004). Ethnicity, gender, and age effects on adrenoceptors and physiological responses to emotional stress. *Psychophysiology*, 41, 450–460.
- Sumter, S.R., Bokhorst, C.L., Miers, A.C., Van Pelt, J., & Westenberg, P.M. (2010). Age and puberty differences in stress responses during a public speaking task: do adolescents grow more sensitive to social evaluation? *Psychoneuroendocrinology*, 35, 1510–1516.
- Susman, E.J., Dockray, S., Granger, D.A., Blades, K.T., Randazzo, W., Heaton, J.A., et al. (2010). Cortisol and alpha amylase reactivity and timing of puberty: vulnerabilities for antisocial behaviour in young adolescents. *Psychoneuroendocrinology*, 35, 557–569.
- Takai, N., Yamaguchi, M., Aragaki, T., Eto, K., Uchihashi, K., & Nishikawa, Y. (2007). Gender- Specific Differences in Salivary Biomarker Responses to Acute Psychological Stress. *Annals of the New York Academy of Sciences*, 1098(1), 510-515.
- Task Force of the European Society of Cardiology and the North American Society of Pacing and Electrophysiology. (1996). Heart rate variability: standards of measurement, physiological interpretation and clinical use. *Circulation*, 93, 1043-1065.
- Taylor, S. E., Klein, L. C., Lewis, B. P., Gruenewald, T. L., Gurung, R. A., & Updegraff, J. A. (2000). Biobehavioral responses to stress in females: tend-and-befriend, not fight-or-flight. *Psychological review*, 107(3), 411.
- Thoma, M.V., Kirschbaum, C., Wolf, J.M., & Rohleder, N. (2012). Acute stress responses in



- salivary alpha-amylase predict increases of plasma norepinephrine. *Biological Psychology*, 91, 342-348.
- Uckuyu, A., Toprak, E., Ciftci, O., & Ciftci, F. C. (2013). The fluctuation in the heart rate variability throughout ovulation induction cycle: is the case different in polycystic ovary syndrome. *Gynecol Obstet*, 3(5), 172-6.
- Uhart, M., Chong, R. Y., Oswald, L., Lin, P. I., & Wand, G. S. (2006). Gender differences in hypothalamic–pituitary–adrenal (HPA) axis reactivity. *Psychoneuroendocrinology*, 31(5), 642-652.
- Van Stegeren, A., Wolf, O. T., & Kindt, M. (2008). Salivary alpha amylase and cortisol responses to different stress tasks: Impact of sex. *International Journal of Psychophysiology*, 69, 33–40.
- Villada, C., Espin, L., Hidalgo, V., Rubagotti, S., Sgoifo, A., & Salvador, A. (2017). The influence of coping strategies and behavior on the physiological response to social stress in women: The role of age and menstrual cycle phase. *Physiology and Behavior*, 170, 37–46.
- Villada, C., Hidalgo, V., Almela, M., & Salvador, A. (2018). Assessing Performance on an Evaluated Speaking Task. *Journal of Psychophysiology*, 32 (2), 64-74.
- Villada, C., Hidalgo, V., Almela, M., Mastorci, F., Sgoifo, A., & Salvador, A. (2014). Coping with an acute psychosocial challenge: behavioral and physiological responses in young women. *PloS one*, 9(12), e114640.
- Vishrutha, K. V., Harini, N., Ganaraja, B., Pavanchand, A., & Veliath, S. (2012). A study of cardiac autonomic control and pulmonary functions in different phases of menstrual

cycle. *International Journal of Applied Biology and Pharmaceutical Technology*, 3(3), 306-311.

Wolf, O. T., Schommer, N. C., Hellhammer, D. H., McEwen, B. S., & Kirschbaum, C.

(2001). The relationship between stress induced cortisol levels and memory differs between men and women. *Psychoneuroendocrinology*, 26(7), 711-720.

Wolf, J. M., Nicholls, E., & Chen, E. (2008). Chronic stress, salivary cortisol, and  $\alpha$ -amylase in children with asthma and healthy children. *Biological psychology*, 78(1), 20-28.

ACCEPTED MANUSCRIPT

## Figure captions

**Fig. 1.** Timeline of the TSST and control (C) conditions. Sequential sAA sampling (from T1 to T6). The Inventory of Situations and Responses of Anxiety (ISRA) was applied during the recovery period

**Fig. 2.** Estimated means for each moment of the TSST condition in the three groups for sAA (a), HR (b), RMSSD (c) and LF/HF (d) measures. Error bars represent standard errors of the mean (SEM).

**Fig. 3.** Means of LF/HF ratio in the TSST and Control conditions in the three groups analyzed. Error bars represent standard error of the mean (SEM)

**Fig. 4.** Means of sAA AUCi in the TSST and control conditions in the three groups analyzed. Error bars represent standard error of the mean (SEM)




Fig. 1. Timeline of the TSST and control (C) conditions. Sequential sAA sampling (from T1 to T6). The Inventory of Situations and Responses of Anxiety (ISRA) was applied during the recovery period.




Fig. 2. Estimated means for each moment of the TSST condition in the three groups for sAA (a), HR (b), RMSSD (c) and LF/HF (d) measures. Error bars represent standard errors of the mean (SEM).




Fig. 3. Means of LF/HF ratio in the TSST and Control conditions in the three groups analyzed. Error bars represent standard error of the mean (SEM).




Fig. 4. Means of sAA AUCi in the TSST and control conditions in the three groups analyzed. Error bars represent standard error of the mean (SEM).

### Table captions

**Table 1:** One-way ANOVA for descriptive characteristics of age, body mass index (BMI), and scores on the subscales and total ISRA score. The values represent means and standard error of the mean (S.E.M.)

N=76	Groups	Mean	S.E.M	Ranges	ANOVA
Age (years)	Luteal	19.00	0.31	18-23	F= 1.50, p=0.22
	Follicular	18.80	0.27	18-24	
	Men	19.56	0.36	18-25	
BMI (kg/m <sup>2</sup> )	Luteal	21.94	0.63	18.65-29.85	F= 5.56, p=0.006
	Follicular	20.81	0.57	16.14-28.96	
	Men	23.69	0.62	18.05-30.39	
ISRA Cognitive (direct scores)	Luteal	90.15	5.82	35-141	F= 2.55, p=0.08
	Follicular	84.48	6.64	23-169	
	Men	70.20	6.83	27-154	

ISRA Physiology (direct scores)	Luteal	56.55	6.52	12.50-128	F= 1.79, p=0.17
	Follicular	50.20	6.93	10-160.50	
	Men	39.22	6.19	6-111	
ISRA Motor (direct scores)	Luteal	56.84	7.33	7-142	F= 0.58, p=0.56
	Follicular	57.44	6.71	5-141	
	Men	48.12	6.24	6-108	
ISRA Total (direct scores)	Luteal	204.86	17.58	77.00-375.50	F= 2.03, p=0.13
	Follicular	192.52	18.45	48.00-434.50	
	Men	155.54	17.90	45.50-371.50	

**Table 1:** One-way ANOVA for descriptive characteristics of age, body mass index (BMI), and scores on the subscales and total ISRA score. The values represent means and standard error of the mean (S.E.M.)