

[Click here to view linked References](#)

## The muscle-bone unit in adolescent swimmers

1  
2 *Alejandro Gomez-Bruton*<sup>1,2,3,4</sup>; *Alejandro Gonzalez-Aguero*<sup>1,2,3,5,8</sup>; *Angel Matute-Llorente*<sup>1,2,3,5</sup>; *Gabriel*  
3 *Lozano-Berges*<sup>1,3,5</sup>; *Alba Gomez-Cabello*<sup>1,2,3,6,8</sup>; *Luis A. Moreno*<sup>1,2,7,8</sup>; *Jose A. Casajus*<sup>1,2,3,5,8</sup>; *Germán*  
4 *Vicente-Rodríguez*<sup>1,2,3,5,8</sup>

5  
6 <sup>1</sup> GENUD (Growth, Exercise, NUtrition and Development) Research Group, Zaragoza, Spain

7 <sup>2</sup> Centro de Investigación Biomédica en Red de Fisiopatología de la Obesidad y Nutrición  
8 (CIBEROBN), Madrid, Spain

9 <sup>3</sup> EXERNET red de investigación en ejercicio físico y salud para poblaciones especiales

10 <sup>4</sup> Universidad Isabel I, Burgos, Spain

11 <sup>5</sup> Faculty of Health and Sport Science (FCSD), Department of Physiatry and Nursing. Universidad de  
12 Zaragoza, Ronda Misericordia 5, 22001-Huesca, Spain.

13 <sup>6</sup> Centro Universitario de la Defensa, Zaragoza, Spain

14 <sup>7</sup> Faculty of Health Science (FCS). Universidad de Zaragoza. Zaragoza, Spain

15 <sup>8</sup> Instituto Agroalimentario de Aragón (IA2), Zaragoza, Spain

### Corresponding author:

16 German Vicente-Rodriguez, PhD.

17 **GENUD (Growth, Exercise, NUtrition and Development) Research Group**

18 Faculty of Health and Sport Sciences

19 University of Zaragoza

20 Grupo GENUD Edificio SAI 2ª planta, Zaragoza

21 *Phone:* +34 876553755 or 976761000 (Ext. 843754)

22 *Fax:* +34 974238422 / +34 974239384

23 *Email:* [gervicen@unizar.es](mailto:gervicen@unizar.es)

1  
2 **Abstract**  
3

4 Purpose: To examine the functional and structural muscle-bone unit in adolescent swimmers.  
5  
6

7 Methods: Sixty-five swimmers (34 girls/31 boys) and 119 controls (51 girls/68 boys) participated in the  
8 study. Muscle cross-sectional area (MCSA), bone mineral content (BMC) and polar strength-strain index  
9 (SSIPOL) were measured in the non-dominant radius by peripheral quantitative computed tomography  
10 (pQCT). Subtotal BMC and lean mass were evaluated with dual-energy X-ray absorptiometry (DXA).  
11  
12

13 Handgrip and isometric knee extension (IKE) tests were performed to determine muscle force. The effect  
14 of MCSA, lean and force on SSIPOL and BMC were tested, and the functional and structural muscle-  
15 bone ratios of swimmers and controls were compared.  
16  
17

18 Results: Both muscle size (MCSA and lean) and muscle force (handgrip and IKE) influenced BMC and  
19 SSIPOL in swimmers and controls similarly. Swimmers presented normal MCSA and lean values for  
20 their height, but when compared to controls, swimmers presented a higher amount of lean and MCSA for  
21 the same BMC or SSIPOL (structural muscle-bone unit). For the functional muscle-bone unit, different  
22 results were found for the lower and upper limbs, as no differences were found for the upper limbs, while  
23 for the lower limbs, swimmers presented higher muscle force for the same amount of BMC.  
24  
25

26 Conclusions: The contradictory results regarding BMC in swimmers found in previous studies could  
27 partly be explained with the findings of the present study that reinforce the idea that swimming is not an  
28 effective sport to practice regarding bone mass, and that the muscle bone-unit is different in swimmers  
29 than in controls.  
30  
31  
32  
33  
34  
35

36 **Keywords: Musculoskeletal; Body composition; Bone Health; Children; Exercise; Swimming**  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## Introduction

The positive effects of physical activity and exercise on bone mineral density (BMD) and bone strength have been recently graded with an A and B level of evidence respectively by the National Osteoporosis Foundation[1]. These positive effects are thought to be due to the mechanical forces that are generated either through impact with the ground (i.e. gravitational or ground-reaction forces) or through skeletal muscle contractions (i.e. muscle forces) performed while engaging in different physical activities. It is nowadays clear that different sports have distinct effects on bone mass, with those without ground-reaction forces like swimming[2] or cycling[3] showing the weakest effect on BMD or bone mineral content (BMC).

In order to try to explain the positive effect of exercise on bone, Frost designed the Mechanostat theory which predicts that an increase in muscle mass or force during growth would create a moderate overload, which would lead to an increase in bone mass or strength[4]. Consequently, many studies adjust BMD or BMC by lean or muscle mass when comparing athletes to controls. These adjustments are done with the premise that lean affects bone equally in both groups (athletes and controls), and are very important as they can change results substantially. For example, in a previous study in which we compared BMD and BMC in swimmers and controls[5], we found few differences between groups (some of them favouring the swimmers), whereas when data were adjusted, swimmers presented lower BMD values for most of the measured variables. In this line, a recent meta-analysis evaluating the effects of swimming on BMD suggested that swimming might increase lean mass without increasing BMD[6], which could entail that the muscle-bone association is different in swimmers than in other athletes and controls.

In order to study the muscle-bone association, Schoenau et al.[7] proposed an algorithm that took height, BMC and muscle into account with the aim of describing the muscle-bone unit in different populations. Two different muscle-bone units can be described. The first one is the structural muscle bone unit in which a ratio of bone and muscle mass is estimated. This structural muscle-bone unit has been criticized as muscle force is not necessarily proportional to muscle size[8]. Consequently, some researchers have advocated for the functional muscle-bone unit which is a ratio of bone and muscle force[8]. Although in other athletic disciplines[9, 10] and in non-athletes[8, 11] this muscle-bone association has been deeply studied, there is a lack of literature exploring this association in adolescent swimmers. It is extremely

1 important to evaluate this muscle-bone association, as if muscle is affecting bone differently in swimmers  
2 than in other athletes or sedentary controls, adjusting by muscle or lean mass when comparing these  
3 groups could be incorrect.  
4

5  
6 Therefore, the aims of the present study were to evaluate if swimmers present a similar functional and  
7 structural muscle-bone unit when compared to controls.  
8  
9

## 10 11 **Material and methods**

12  
13  
14 Study design, protocol and consent forms were performed in accordance with the Helsinki Declaration of  
15 1964 (revised in Fortaleza, 2013) and were reviewed and approved by the Research Ethics Committee of  
16 the Government of Aragon (ref. CP08/2012, CEICA, Spain). Sample size calculations for the present  
17 project have been explained elsewhere[12], and the sample size is similar to that presented in previous  
18 articles evaluating the muscle-bone unit in children and adolescents[13, 14].  
19  
20  
21  
22  
23

### 24 25 *Participants*

26  
27  
28 All participants had to be between the ages of 11 and 18 years, Caucasian, non-smokers, with no chronic  
29 disease or musculoskeletal disorders (fibromyalgia, gout, osteoarthritis, rheumatoid arthritis, tendinitis),  
30 bone fractures or medication. Swimmers had to have a history of swimming and competing in regional  
31 tournaments for more than 3 years and current training for a minimum of 6 hours per week. The normo-  
32 active controls (CG) could not be performing any aquatic activity on a regular basis or more than 3 hours  
33 of weight-bearing physical activity per week. Swimmers were recruited from 4 swimming clubs while  
34 controls were recruited from 4 schools and high-schools of Zaragoza (Spain).  
35  
36  
37  
38  
39  
40  
41  
42

### 43 44 *Pubertal stage and anthropometric measurements*

45  
46 Pubertal maturation was determined by self-assessment of secondary sexual characteristics according to  
47 the criteria devised by Tanner[15]. This method has been reported to be both valid and reliable in  
48 assessing sexual maturity among adolescent athletes[16].  
49  
50  
51  
52  
53

54 Height was measured with a stadiometer to the nearest 0.1 cm (SECA 225, SECA, Hamburg, Germany),  
55 and weight with a scale to the nearest 0.1 kg (SECA 861, SECA, Hamburg, Germany). Both  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 anthropometric measures were taken without shoes and minimal clothing following the procedures by  
2 International Society for the Advancement in Kinanthropometry (ISAK)[17].  
3  
4

### 5 *Muscle force*

6

7  
8  
9 Maximum isometric forearm force was determined with a digital handgrip dynamometer (Takei TKK  
10 5401, Takei scientific instruments, Tokyo, Japan). Participants were instructed to exert their maximal grip  
11 force with the upper limb in extension in two attempts with each hand, with one minute pauses between  
12 them. The dynamometer was adapted for each participant according to their hand span as proposed by  
13 Ruiz et al.[18] The best attempt was selected for further analysis.  
14  
15  
16  
17

18  
19  
20 Maximum isometric peak torque through knee extension (IKE) was measured using a strain gauge  
21 (MuscleLab, Force Sensor, Norway). The participant was sitting with an anchorage placed on the distal  
22 third of the tibia. This anchorage was connected to the strain gauge, registering force data during the 6  
23 seconds that the participant performed the maximum knee extension. Two attempts were allowed for each  
24 leg, recording the best performance.  
25  
26  
27  
28  
29

### 30 *Bone assessments*

31

32  
33  
34 Whole body dual-energy X-ray absorptiometry (DXA) scans were performed with the pediatric version of  
35 the QDR-Explorer software (Hologic Corp. Software version 12.4. Bedford. Masssachusettes, USA).  
36

37  
38 Outcomes of interest were BMC and lean mass of the subtotal body (whole body – head) and upper- and  
39 lower-limbs (obtained from the whole-body scan). All the DXA analyses were performed by the same  
40 operator (AML). The coefficients of variation of the DXA in our laboratory for a whole body scan were  
41 2.3% for BMC and 1.9% for lean mass. For DXA, the final sample consisted of 65 swimmers (34 girls /  
42 31 boys) and 119 controls (51 girls / 68 boys).  
43  
44  
45  
46  
47

48  
49 The radius was also measured with a stratec XCT-2000 L peripheral quantitative computed tomography  
50 (pQCT) scanner (Stratec Medizintechnik, Pforzheim, Germany). Scanning procedure and coefficients of  
51 variation in our laboratory have been described in detail elsewhere[19]. For the present study, data of the  
52 66% of the total radius length is presented. The same researcher (AML) performed all the scans. Firstly, a  
53 scout view was performed to manually locate a reference line on the distal end of the radius. The  
54 reference line was placed at the endplate of the radius as suggested by the Stratec XCT-2000 manual.  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 Radius total BMC, polar bone strength strain index (SSIPOL; an estimate of the resistance to bending and  
2 torsion of the bone)[20], and muscle cross-sectional area (MCSA), were evaluated. As some of the  
3 participants presented moved scans, the sample with good quality pQCT scans consisted of 55 swimmers  
4 (29 girls / 26 boys) and 88 controls (39 girls /49 boys).  
5  
6

### 7 8 *Statistical analyses* 9

10 Independent t-tests were performed to compare anthropometric data between swimmers and controls.  
11

12  
13 Linear regressions were performed to evaluate the association between height and muscle mass and to  
14 determine if this association was similar in controls and swimmers. Further linear regressions were  
15 performed to test the influence of muscle size (MCSA or lean) or muscle force (handgrip or IKE) on bone  
16 (BMC or SSIPOL), and determine if this influence was similar in controls and swimmers.  
17  
18  
19  
20  
21

22  
23 Lean mass, BMC, and MCSA mean values from the control group (dividing the controls into height  
24 groups spanning 10 cm each) were used as a reference, and five different age- and pubertal-adjusted ratios  
25 (a. BMC to MCSA; b. BMC to Handgrip; c. BMC to Lean; d. SSIPOL to MCSA; e. SSIPOL to  
26 Handgrip) were calculated for this group. Z-scores were after calculated for the swimmer group with the  
27 previously described control values. To evaluate whether a parameter was significantly different from the  
28 result in controls, the difference of the mean Z score to zero was assessed with one-sample t-tests. A  
29 significant difference was assumed when the 95% CI of the mean Z-score did not include zero.  
30  
31  
32  
33  
34  
35  
36  
37

38 For all analyses, statistical significance was set as  $p < .05$ . All analyses were performed with SPSS version  
39 21.0.  
40  
41

## 42 43 **Results** 44

### 45 46 *Descriptive characteristics* 47

48 Anthropometric characteristics are summarized in Table 1. No age differences were found between  
49 groups ( $p > .05$ ; Table 1). Male swimmers were taller and heavier than controls ( $p < .05$ ; Table 1).  
50  
51

52 Female swimmers presented lower leg BMC values when compared to controls ( $p < .05$ ; Table 1). No  
53 other difference was found between groups for any of the measured bone variables independently of the  
54 used device (DXA or pQCT). Male swimmers presented higher values of subtotal and arms lean than  
55 controls (both  $p < .05$ ; Table 1).  
56  
57  
58  
59  
60  
61  
62

*Influence of height, lean, force and group (swimmer or control) on bone mineral content and bone strength*

Height was positively associated with MCSA and lean mass in both girls and boys ( $p < .05$ ; data not presented). No significant group by height interactions were found, suggesting that height influenced lean and MCSA similarly in both groups (swimmers and controls).

All muscle (MCSA and lean) and force variables (handgrip and IKE) were positively associated with BMC (measured with DXA and pQCT) and SSIPOL (pQCT) in both girls and boys (all  $p < .05$ ; Table 2). Regarding the effect of group, both male and female swimmers presented lower values than controls for subtotal, arms and legs BMC measured with DXA when adjusted by lean (Table 2). Nonetheless, when adjusting arms BMC by muscle force these differences disappeared ( $p > .05$ ; Table 2) for both genders. When adjusting legs BMC by IKE the differences between groups also disappeared for males. No differences were found between groups for BMC or SSIPOL measured with pQCT (all  $p > .05$ ; Table 2).

*Schoenau algorithm*

Data from Tables 3 and 4 were used to calculate z-scores and apply the Schoenau algorithm[7].

For step 1 of the Schoenau muscle-bone unit[7] (is muscle adequate for height?), sex- and height-specific z-scores were computed with data from Table 3. The mean height-dependent Z-score of subtotal lean and MCSA were not significantly different from zero in the swimmers (both  $p > .05$ ; Table 5), suggesting that muscle mass (MCSA or Lean mass) was adequate for height in swimmers.

Regarding step 2 of the Schoenau muscle-bone unit[7] (is BMC adequate for muscle mass or force?), all the structural muscle-bone unit ratios were significantly lower in swimmers than in controls (both age and Tanner dependent  $p < .05$ ; Table 5), suggesting that BMC was not adequate for muscle mass in swimmers.

Regarding the functional muscle-bone unit, the legs BMC/force ratio was lower in swimmers than in controls (both age and Tanner dependent  $p < .05$ ; Table 5), which entails that legs BMC was not adequate for legs muscle force.

**Discussion**

The main finding of the present study is that swimmers present lower structural muscle-bone unit ratios, suggesting that they have a lower amount of bone for the same amount of lean mass or higher amount of

1 lean mass for the same amount of bone. For the functional muscle-bone unit, different results were found  
2 for the lower and upper limbs, as no differences were found for the upper limbs between swimmers and  
3 controls, while for the lower limbs, swimmers presented lower BMC for the same amount of muscle force  
4 or higher muscle force for the same BMC.  
5  
6

7  
8  
9 The results found with the linear regressions suggest that both muscle (lean or MCSA) and force  
10 (handgrip or IKE) are determinant to BMC and SSIPOL in swimmers. These findings are in line with  
11 previous studies that evaluated this association in healthy children and adults[21] and children with  
12 different conditions[7]; all finding a strong association between MCSA and BMC. No group by muscle  
13 interactions were found, suggesting that the mechanisms regulating the muscle-bone interaction in  
14 swimmers and controls are similar. These results were expected, as swimmers are not a population with a  
15 specific disease or condition that could regulate their bone-muscle mechanisms in any way as found in  
16 other populations[7].  
17  
18  
19  
20  
21  
22  
23  
24

25  
26 The negative Z-scores found for the structural muscle bone-unit ratios in swimmers, both when taking  
27 into account age or Tanner stage could be interpreted in two different ways: 1) swimmers present a higher  
28 amount of muscle for each unit of BMC/SSIPOL, or 2) swimmers present a lower amount of BMC and a  
29 weaker bone per each unit of lean/MCSA. When focusing on boys, it seems clear that although there are  
30 no significant differences between BMC, SSIPOL or MCSA between groups (Table 1), swimmers present  
31 higher mean values for MCSA (not significant) and lean (significant). Therefore, swimmers present more  
32 MCSA and lean for the same amount of BMC or SSIPOL. This would imply that swimming is effective  
33 for improving lean mass but not for stimulating bone mass, at least in the regions in which measurements  
34 were performed. This is in line with previous studies that compared swimmers with sedentary controls  
35 finding higher lean mass values in the swimmers without higher values in BMC[22] or BMD[23]. This  
36 lack of bone improvements in swimmers could be due to two different and not necessarily independent  
37 factors; the lack of impact, and the low mechanical strains produced while swimming.  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

50  
51 Focusing on impact it has been widely demonstrated that sports that entail high impacts improve bone  
52 mass[24–26], and those reviews that included a variety of sports always found that athletes involved in  
53 high impact sports presented higher bone values when compared to non-impact sports[27, 28].  
54  
55 Nonetheless, not all impacts have the same effect, as low impacts repeated many times might not be as  
56 useful, as demonstrated in a review performed by Tenforde and Fredericson, who showed that athletes  
57  
58  
59  
60  
61  
62  
63  
64  
65



1 engaged in high-impact and multidirectional loading sports consistently had greater BMD and geometric  
2 properties when compared to distance runners[27]. It consequently seems like impacts are determinant to  
3 bone mass, while the only impacts received by swimmers are those performed while kicking against the  
4 wall which will probably entail very low loads.  
5  
6

7  
8 Regarding the mechanical strains produced while swimming, Meaking et al.[29] described in a review  
9 article the mechanisms of adaptation to mechanical loading in bone. Furthermore they described some  
10 main points that when specifically analysed and extrapolated to the characteristics of swimming are just  
11 the opposite of what the sport demands: 1) "Loading-related bone formation correlates with peak strain  
12 magnitude": swimming, like most endurance sports, is characterized by low magnitudes and therefore  
13 peak strains will be relatively low; 2) "Increasing strain rate during loading stimulates bone formation":  
14 the rate of change of strain magnitude or in other words the acceleration or deceleration which is very low  
15 in swimming is also a key factor to bone stimulation; 3) "Number of loading cycles required to maximally  
16 stimulate bone formation is small"; relatively few cycles of loading are required to maximize the amount  
17 of new bone formed, while in swimming there are thousands of cycles in each training session; 4)  
18 "Inserting rest between loading cycles increases bone formation". An animal study by Robling et al.[30]  
19 showed that dividing 360 loading cycles into 4 bouts of 90 cycles or 6 bouts of 60 cycles per day  
20 enhanced the osteogenic response to loading. Nevertheless, when competing in swimming unlike other  
21 sports (i.e. soccer, basketball) there are no rest periods between loading cycles. Therefore, these 4  
22 described points in addition to the lack of eccentric movements in swimming, which have shown to be  
23 more effective than concentric movements in improving bone mass[31], could mostly explain why the  
24 structural muscle-bone ratios were lower in swimmers than in CG.  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43

44 When focusing on the functional muscle-bone unit, results were different, as no differences for the z-  
45 scores were found for the BMC/handgrip and SSIPOL/handgrip ratios, suggesting that swimmers and  
46 controls have the same muscle force for the same amount of bone. This does not necessarily imply that  
47 swimming is not improving muscle force, as we found that when measuring force with a specific test that  
48 simulated swimming stroke, swimmers performed higher amounts of force than controls (unpublished  
49 personal findings). Nevertheless, the handgrip test measures the maximal isometric force performed by  
50 the forearms and while adolescent swimmers perform many repetitive movements that involve the  
51 forearm muscles, they never reach maximum force that, therefore, will probably not be increased as  
52 shown in the present study. Surprisingly, the leg functional bone-unit was different between swimmers  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62

1 and controls, probably because swimmers presented higher (although not significant) leg muscle force  
2 values and similar bone values. The different results found for the upper and lower limb functional  
3 muscle-bone unit, could be due to the used non-specific tests to assess muscle force, as although both are  
4 unspecific tests, swimmers while performing the turn phase under-water complete a quick and strong  
5 push against the wall, which might strengthen the quadriceps (the main muscles involved in the isometric  
6 knee extension test). It is possible that if two specific swimming tests had been performed to measure the  
7 upper and lower body force, swimmers would have indeed presented higher force values when compared  
8 to the controls, and therefore, results for the functional muscle-bone unit would have been similar to the  
9 structural muscle bone unit, with swimmers presenting higher muscle force values for the same amount of  
10 bone.

11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21 Although this study presents several strengths, such as the use of two different devices to measure bone  
22 mass, and the estimation of both functional and structural muscle-bone units, it is not extent of  
23 limitations, with the main one being the type of muscle force assessment; isometric tests. A type of  
24 muscle contraction that is never performed by swimmers while swimming. Additionally, and as explained  
25 in previous studies[8] maximum forces that stimulate bone will be reached with eccentric contractions,  
26 and not with isometric ones. Therefore, further studies, using other types of strength assessment in  
27 swimmers are needed in order to confirm our results.

28  
29  
30  
31  
32  
33  
34  
35  
36 The present findings could partly explain the inconsistent results regarding bone mass in adolescent  
37 swimmers, as although literature is scarce for pQCT studies, there is a vast amount of literature regarding  
38 DXA studies with diverse results[2]. This variety of results could partly be explained by the effect of lean  
39 mass, as some researchers adjust BMD or BMC by lean mass. The assumption behind this common  
40 adjustment is that lean mass influences bone mass in a similar manner in swimmers and CG. However,  
41 our data suggest that the ratio BMC to lean is not similar so lean adjustment should be avoided as when  
42 comparing swimmers to CG, swimmers will have more lean mass (although the ratio of BMC to lean  
43 mass will be lower in the swimmers). Therefore, when adjusting by lean mass, BMC will become lower  
44 in the swimmers when compared to the CG due to the effect of the covariate. Thus, it is possible that  
45 differences in bone mass might emerge between both groups when those differences in raw data are not  
46 present. This happened in the current study, as when comparing swimmers to controls without adjustment  
47 no differences were found for bone variables between groups (Table 1), but when groups were compared  
48 adjusting by lean, swimmers presented lower BMC values (Table 2).

1 According to the muscle-bone algorithm proposed by Schoenau et al.[7], swimmers would be classified  
2 as a population with a primary bone defect, as they do present a normal MCSA for their height, but they  
3 do not present an adequate BMC for their MCSA/lean. Nevertheless, the mentioned algorithm was  
4 proposed for pediatric bone disease, and populations that are studied with it will generally present normal  
5 muscle and low bone as found in kidney transplant patients[7], unlike the swimmers of the present study  
6 that presented high muscle mass and normal BMC. Consequently, swimmers should not be considered as  
7 a population with primary bone defects. Nonetheless, the idea that swimming is not an effective sport to  
8 practice regarding bone mass[6] was reinforced by the present study. Swimming coaches should  
9 implement weight-bearing trainings in addition to water training in order to try to improve adolescent  
10 swimmers bone mass.  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22

#### 23 **Competing interests:**

24 All authors declare no disclosures  
25  
26  
27  
28  
29  
30  
31

#### 32 **Authors' contributions**

33 AGB, AML, AGA, AGC and GLB collected all the data. AGB and AGA prepared the database and the  
34 statistical analyses and drafted the first manuscript. LM, JAC and GVR conceived the study, participated  
35 in its design and obtained funding. GVR and AGB coordinated the study. All authors have read and  
36 approved the final version of the manuscript.  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## References

1. Weaver CM, Gordon CM, Janz KF, et al (2016) The National Osteoporosis Foundation's position statement on peak bone mass development and lifestyle factors: a systematic review and implementation recommendations. *Osteoporos Int* 27:1281–1386
2. Gomez-Bruton A, Gonzalez-Aguero A, Gomez-Cabello A, et al (2013) Is bone tissue really affected by swimming? A systematic review. *PLoS One* 8:e70119
3. Olmedillas H, Gonzalez-Aguero A, Moreno LA, et al (2012) Cycling and bone health: a systematic review. *BMC Med* 10:168
4. Burr DB (1997) Muscle strength, bone mass, and age-related bone loss. *J Bone Min Res* 12:1547–1551
5. Gomez-Bruton A, González-Agüero A, Gómez-Cabello A, et al (2015) The effects of swimming training on bone tissue in adolescence. *Scand J Med Sci Sport* 25:
6. Gomez-Bruton A, Montero-Marín J, González-Agüero A, et al (2016) The Effect of Swimming During Childhood and Adolescence on Bone Mineral Density: A Systematic Review and Meta-Analysis. *Sport Med* 46: . doi: 10.1007/s40279-015-0427-3
7. Schoenau E, Neu CM, Beck B, et al (2002) Bone mineral content per muscle cross-sectional area as an index of the functional muscle-bone unit. *J Bone Min Res* 17:1095–1101
8. Anliker E, Toigo M (2012) Functional assessment of the muscle-bone unit in the lower leg. *J Musculoskelet Neuronal Interact* 12:46–55
9. Ireland A, Maden-Wilkinson T, McPhee J, et al (2012) Upper limb muscle-bone asymmetries and bone adaptation in elite youth tennis players. *Med Sci Sport Exerc* 45:1749–1758
10. Calbet JA, Moysi JS, Dorado C, Rodriguez LP (1998) Bone mineral content and density in professional tennis players. *Calcif Tissue Int* 62:491–496
11. Schoenau E (2005) The “functional muscle-bone unit”: a two-step diagnostic algorithm in pediatric bone disease. *Pediatr Nephrol* 20:356–359
12. Gomez-Bruton A, Gonzalez-Aguero A, Gomez-Cabello A, et al (2015) Bone structure of adolescent swimmers; a peripheral quantitative computed tomography (pQCT) study. *J Sci Med Sport* 19:707–712
13. Veilleux L-N, Pouliot-Laforte A, Lemay M, et al (2015) The functional muscle–bone unit in patients with osteogenesis imperfecta type I. *Bone* 79:52–57 . doi: 10.1016/j.bone.2015.05.019
14. Soucek O, Matyskova J, Anliker E, et al (2015) The muscle–bone interaction in Turner syndrome. *Bone* 74:160–165 . doi: 10.1016/j.bone.2015.01.017
15. Tanner JM, Whitehouse RH, Takaishi M (1966) Standards from birth to maturity for height, weight, height velocity, and weight velocity: British children, 1965. II. *Arch Dis Child* 41:613–635
16. Leone M, Comtois AS (2007) Validity and reliability of self-assessment of sexual maturity in elite adolescent athletes. *J Sport Med Phys Fit* 47:361–365
17. Marfell-Jones M, Olds T, Stewart A, Carter L (2006) International Standards for Anthropometric assessment. International Society for the Advancement of Kinanthropometry (ISAK). Potchefstroom, South Africa
18. Ruiz JR, Espana-Romero V, Ortega FB, et al (2006) Hand span influences optimal grip span in male and female teenagers. *J Hand Surg Am* 31:1367–1372
19. Gomez-Bruton A, González-Agüero A, Casajus JA, Vicente-Rodríguez G (2014) Swimming training repercussion on metabolic and structural bone development. Benefits of the incorporation

of whole body vibration or pliometric training. The RENACIMIENTO project. *Nutr Hosp* 30:399–409

20. Schiessl H, Ferretti JL, Tysarczyk-Niemeyer G, Willnecker J (1996) Noninvasive bone strength index as analysed by peripheral quantitative computed tomography (pQCT). In: Schoenau E (ed) *Paediatric osteology: new developments in diagnostics and therapy*. Elsevier, Amsterdam, The Netherlands, pp 141–146
21. Schoenau E, Neu CM, Mokov E, et al (2000) Influence of puberty on muscle area and cortical bone area of the forearm in boys and girls. *J Clin Endocrinol Metab* 85:1095–1098
22. Jurimae J, Cicchella A, Tillmann V, et al (2009) Effect of pubertal development and physical activity on plasma ghrelin concentration in boys. *J Endocrinol Invest* 32:18–22
23. Ferry B, Lespessailles E, Rochcongar P, et al (2013) Bone health during late adolescence: Effects of an 8-month training program on bone geometry in female athletes. *Jt Bone Spine* 80:57–63
24. Hagman M, Helge EW, Hornstrup T, et al (2018) Bone mineral density in lifelong trained male football players compared with young and elderly untrained men. *J Sport Heal Sci* 7:159–168 . doi: 10.1016/J.JSHS.2017.09.009
25. Lozano-Berges G, Matute-Llorente Á, González-Agüero A, et al (2017) Soccer helps build strong bones during growth: a systematic review and meta-analysis. *Eur J Pediatr*. doi: 10.1007/s00431-017-3060-3
26. Burt LA, Greene DA, Ducher G, Naughton GA (2013) Skeletal adaptations associated with pre-pubertal gymnastics participation as determined by DXA and pQCT: a systematic review and meta-analysis. *J Sci Med Sport* 16:231–239
27. Tenforde AS, Fredericson M (2011) Influence of sports participation on bone health in the young athlete: a review of the literature. *PM&R* 3:861–867
28. Guadalupe-Grau A, Fuentes T, Guerra B, Calbet JA (2009) Exercise and bone mass in adults. *Sport Med* 39:439–468
29. Meakin LB, Price JS, Lanyon LE (2014) The Contribution of Experimental in vivo Models to Understanding the Mechanisms of Adaptation to Mechanical Loading in Bone. *Front Endocrinol* 5:154
30. Robling AG, Burr DB, Turner CH (2000) Partitioning a daily mechanical stimulus into discrete loading bouts improves the osteogenic response to loading. *J Bone Min Res* 15:1596–1602
31. Hawkins SA, Schroeder ET, Wiswell RA, et al (1999) Eccentric muscle action increases site-specific osteogenic response. *Med Sci Sport Exerc* 31:1287–1292

Table 1. Anthropometric characteristics, bone and muscle strength variables

	Girls (n=85)		Boys (n=99)	
	CG (n=51)	SWI (n=34)	CG (n=68)	SWI (n=31)
<i>Age</i> (y)	14.2±2.3	13.9±1.9	14.9±2.3	15.1±1.5
<i>Weight</i> (kg)	50.9±10.9	47.8±8.4	55.9±12.6*	61.7±10.1
<i>Height</i> (cm)	156.2±7.8	156.8±9.1	164.8±12.4*	171.4±8.9
<i>Tanner stage</i> (I/II/III/IV/V)	2/9/16/10/14	1/8/8/16/1	1/10/10/15/32	0/2/8/13/8
<b>DXA</b>				
<i>Subtot_BMC</i> (g)	1271.134±316.906	1188.993±310.165	1540.004±482.605	1598.500±374.931
<i>Arms_BMC</i> (g)	101.711±26.937	99.255±25.836	121.572±41.426	135.299±33.480
<i>Legs_BMC</i> (g)	318.505±75.187*	283.895±66.770	402.011±119.235	404.741±93.503
<i>Subtot_Lean</i> (g)	30591.035±5779.726	29968.373±5767.138	38415.18±9860.463*	43028.005±8214.649
<i>Legs_Lean</i> (g)	5680.941±1116.848	5333.074±996.291	7274.547±1863.74	7949.562±1545.221
<i>Arms_Lean</i> (g)	1502.969±273.985	1569.289±308.990	2136.03±634.143*	2517.67±547.38
<b>pQCT<sup>#</sup></b>				
<i>Radius length</i> (mm)	232.82±14.17	234.59±12.43	252.41±20.92	259.58±21.04
<i>Radius BMC</i> (mg/mm)	0.81±0.14	0.77±0.14	0.90±0.22	0.90±0.15
<i>Radius SSIPOL</i> (N)	207.8±52.53	200.33±48.7	256.14±82.57	268.71±54.09
<i>Radius MCSA</i> (mm <sup>2</sup> )	2041.19±295.54	2101.31±358.14	2748.30±697.22	2907.58±630.95
<b>Muscle strength</b>				
<i>Handgrip</i> (kg)	22.1±4.9	22.3±5.2	29.7±9.7	32.0±7.3
<i>Leg force</i> (kg)	38.3±11.5	37.9±10.6	47.3±14.5	51.4±10.4

Values are mean ± standard deviation

\*P<0.05 Difference between groups within gender.

# for pQCT variables there were 68 girls (39 controls vs. 29 swimmers) and 75 boys (49 controls vs. 26 boys)

Arms=Mean value of the left and right arm; BMC=Bone mineral content; DXA=Dual energy x-ray; Handgrip=Mean values of the right and left arm;

Leg force= Mean leg force of the left and right legs; Legs=Mean value of the left and right arm; MCSA=Muscle cross-sectional area;

pQCT=Peripheral quantitative computed tomography; SSIPOL=Polar strain strength index; Subtot=Whole body less the head.

Table 2. Results of the different centered linear regressions including both swimmers and controls

<i>Independent variables</i>	Structural muscle bone unit					Functional muscle bone unit			
		DXA		PQCT		DXA		PQCT	
<i>Girls</i>	Subt. BMC	Arms BMC	Legs BMC	BMC <sub>1</sub>	SSIPOL <sub>1</sub>	Arms BMC	Legs BMC	BMC <sub>2</sub>	SSIPOL <sub>2</sub>
<i>β Group</i>	-0.080 <sup>‡</sup>	-0.144*	-0.093 <sup>‡</sup>	-0.272	-0.168	-0.034	-0.208*	-0.109	-0.065
<i>β Lean or MCSA</i>	0.921*	0.921*	0.876*	0.762*	0.697*	-	-	-	-
<i>β Muscle strength</i>	-	-	-	-	-	0.855*	0.715*	0.788*	0.785*
<i>β Interaction</i>	-0.011	-0.066	-0.001	-0.100	-0.050	-0.021	0.048	0.019	-0.021
<i>Model r square</i>	0.849	0.763	0.802	0.515	0.448	0.707	0.607	0.655	0.601
<i>Boys</i>	Subt. BMC	Arms BMC	Legs BMC	BMC <sub>1</sub>	SSIPOL <sub>1</sub>	Arms BMC	Legs BMC	BMC <sub>2</sub>	SSIPOL <sub>2</sub>
<i>β Group</i>	-0.145*	-0.100*	-0.141*	-0.053	0.064	0.084	-0.083	0.060	0.155
<i>β Lean or MCSA</i>	0.986*	0.977*	0.971*	0.917*	0.912*	-	-	-	-
<i>β Muscle strength</i>	-	-	-	-	-	0.851*	0.799*	0.842*	0.808*
<i>β Interaction</i>	-0.011	-0.037	-0.060-	-0.096	-0.167	0.007	0.022	-0.037	-0.091
<i>Model r square</i>	0.884	0.874	0.866	0.754	0.705	0.754	0.642	0.679	0.598

\*p<0.05; <sup>‡</sup>p<0.07

*β*=Beta standardized coefficient

Group=A negative symbol indicates that swimmers have lower values than controls

Muscle force=For the arms handgrip and for the legs maximum isometric force for leg extension

Interaction=Interaction between group (SWI or CG) and lean mass or muscle force

Table 3. Height-dependent results for BMC and lean for the subtotal body (whole body less the head), arms and legs of the controls

Height range cm	DXA								pQCT							
	Girls	Boys	Girls	Boys	Girls	Boys	Girls	Boys	Girls	Boys	Girls	Boys				
	Subtotal				Arms				Legs				Radius			
	BMC (kg)				BMC (kg)				BMC (kg)				BMC (mg/mm <sup>2</sup> )			
<150	9	10	0.842±0.134	0.857±0.115	0.065±0.010	0.065±0.010	0.221±0.032	0.228±0.032	8	7	0.644±0.087	0.610±0.095				
150-160	24	15	1.210±0.200	1.212±0.128	0.098±0.019	0.089±0.012	0.299±0.047*	0.328±0.035	21	11	0.817±0.119	0.771±0.099				
160-170	17	18	1.563±0.200	1.551±0.243	0.124±0.017	0.127±0.022	0.390±0.045	0.401±0.054	9	12	0.913±0.092	0.939±0.092*				
170-180		16		1.941±0.339		0.151±0.028		0.501±0.083		12		1.014±0.184				
>180		9		2.106±0.313		0.172±0.025		0.542±0.080		7		1.124±0.136				
	Lean (kg)				Lean (kg)				Lean (kg)				MCSA (mm <sup>2</sup> )			
<150	9	10	22.063±3.139	22.929±2.483	1.137±0.142	1.202±0.142	4.144±0.629	4.366±0.593	8	7	1696.2±270.9	1750.2±175.9				
150-160	24	15	30.224±3.793	32.045±3.117	1.482±0.185*	1.698±0.268	5.576±0.762*	6.219±0.653	21	11	2101.5±229.2*	2331.3±334.3				
160-170	17	18	35.252±3.478*	38.838±4.303	1.714±0.222*	2.176±0.310	6.534±0.733*	7.200±0.751	9	12	2200.1±248.7*	2751.8±519.2				
170-180		16		47.380±4.892		2.709±0.415		8.970±0.794		12		3376.3±511.2				
>180		9		49.453±5.660		2.802±0.285		9.396±1.441		7		3319.1±220.2				

Values are mean ± standard deviation

\*Significant differences between results in girls and boys of the same height group (p<0.05 in each case)

Arms=Mean value of left and right arm; DXA=Dual energy X-ray; Legs=Mean value of left and right leg; MCSA= muscle cross-sectional area (mm<sup>2</sup>); pQCT=Peripheral quantitative computed tomography; Subtotal=Whole body excluding the head.



Table 4. Variation with age and Tanner stage of following ratios (only for controls): BMC/lean mass, BMC/force, BMC/MCSA, SSIPOL/MCSA, SSIPOL/Handgrip

DXA	n		Structural muscle-bone unit						Functional muscle-bone unit			
			Subtotal body		Arms		Legs		Arms		Legs	
			BMC <sub>(g)</sub> /Lean <sub>(kg)</sub> ratio		BMC <sub>(g)</sub> /lean <sub>(kg)</sub> ratio		BMC <sub>(g)</sub> /Lean <sub>(kg)</sub> ratio		BMC <sub>(g)</sub> /Handgrip <sub>(kg)</sub> ratio		BMC <sub>(g)</sub> /Leg force <sub>(kg)</sub> ratio	
Girls	Boys	Girls	Boys	Girls	Boys	Girls	Boys	Girls	Boys	Girls	Boys	
Age (y)												
<13	22	17	39.28±3.68	37.26±3.26	60.31±8.21*	54.85±6.37	54.46±5.69	52.12±4.67	4.34±0.74	4.15±0.54	8.82±2.16	8.27±1.56
13-15	9	15	40.86±3.85	38.89±2.34	67.36±8.88*	53.15±4.69	53.54±4.32	54.55±3.70	4.84±0.97*	3.88±0.62	9.34±1.48	8.93±1.37
15-17	11	16	42.24±3.08	41.06±4.40	72.38±5.50*	58.72±6.86	56.58±5.06	56.08±5.24	4.86±0.59	4.36±0.66	8.31±1.96	8.22±0.98
>17	9	20	44.96±5.18 <sup>‡</sup>	41.18±3.89	76.47±9.57*	58.57±7.23	60.89±6.24	56.61±5.93	4.81±0.31*	4.02±0.68	7.57±1.25	7.96±1.67
Tanner stage												
II	9	10	38.07±2.34	39.04±3.15	56.64±4.27	55.61±4.69	55.52±3.80	54.45±4.87	3.99±0.62	4.21±0.60	8.40±1.55	8.28±1.53
III	16	10	41.29±4.05*	36.99±3.84	65.57±7.80*	53.05±6.99	53.91±3.56	52.17±5.34	4.49±0.77 <sup>‡</sup>	3.88±0.66	9.40±1.87	8.59±1.51
IV	10	15	42.92±4.70*	38.51±1.99	74.92±7.39*	54.43±6.01	55.45±6.39	53.94±3.15	4.82±0.85*	3.99±0.54	7.92±1.73	8.53±1.62
V	17	37	43.15±5.38	41.34±4.55	72.27±10.50*	58.93±6.73	57.16±7.81	56.50±5.79	5.12±0.40*	4.19±0.70	8.04±1.96	8.18±1.38
PQCT	n		BMC <sub>(mg/mm<sup>2</sup>)</sub> /MCSA <sub>(mm<sup>2</sup>)</sub>		SSIPOL <sub>(mm<sup>3</sup>)</sub> /MCSA <sub>(kg)</sub>		BMC <sub>(mg/mm)</sub> /Hangrip <sub>(kg)</sub>		SSIPOL <sub>(mm<sup>3</sup>)</sub> /Hangrip <sub>(kg)</sub>			
			ratio <sup>Δ</sup>		ratio		ratio		ratio			
Age												
<13	16	11	3.75±0.35 <sup>#</sup>	3.42±0.51	0.090±0.015	0.083±0.021			39.98±9.13	38.75±8.99	9.473±1.982	9.336±2.862
13-15	7	11	3.88±0.32 <sup>#</sup>	3.47±0.54	0.102±0.016	0.097±0.020			37.48±3.22	33.92±5.21	9.792±1.473	9.498±1.774
15-17	9	12	4.04±0.39*	3.27±0.46	0.112±0.013*	0.093±0.014			37.16±4.14*	28.08±4.13	10.299±1.093*	7.959±1.185
>17	7	15	4.38±0.75*	3.16±0.35	0.115±0.029 <sup>#</sup>	0.097±0.016			37.02±6.32*	27.70±5.37	9.688±2.250	8.524±2.168
Tanner stage												
II	7	5	3.71±0.29	3.73±0.56	0.092±0.014	0.094±0.019			39.59±6.07	44.27±6.84	9.813±1.791	11.075±1.394
III	13	8	3.85±0.38	3.55±0.62	0.100±0.015	0.104±0.024			37.33±5.27	36.69±6.80	9.671±1.679	10.715±2.349
IV	8	12	4.35±0.72*	3.29±0.30	0.115±0.029*	0.087±0.023			38.00±5.40*	32.23±4.73	9.939±2.015	8.41±2.194
V	10	24	3.98±0.37*	3.16±0.40	0.103±0.018*	0.092±0.012			36.39±4.39*	27.09±4.21	9.381±1.746*	7.855±1.232

\*Significant differences between girls and boys of the same age or Tanner group (p<0.05).

<sup>‡</sup> Tendency towards differences between girls and boys of the same age or Tanner group (p≤0.07)

Table 5. Z-score values for the different analyses

			Mean dif.	SE	CI		p
Subtotal lean - Height dependent			-0.018	0.125	-0.267	0.231	0.887
MCSA – Height dependent			0.141	0.159	-0.178	0.459	0.380
<i>Age dependent z-scores</i>							
<i>Structural muscle-bone unit</i>	DXA	Subtotal BMC/Lean	<b>-0.720</b>	<b>0.148</b>	<b>-1.015</b>	<b>-0.424</b>	<b>&lt;0.001</b>
		Arms BMC/Lean	<b>-0.451</b>	<b>0.113</b>	<b>-0.676</b>	<b>-0.225</b>	<b>&lt;0.001</b>
		Legs BMC/Lean	<b>-0.744</b>	<b>0.151</b>	<b>-1.045</b>	<b>-0.444</b>	<b>&lt;0.001</b>
	PQCT	BMC/MCSA	<b>-0.596</b>	<b>0.139</b>	<b>-0.875</b>	<b>-0.317</b>	<b>&lt;0.001</b>
		SSIPOL/MCSA	<b>-0.273</b>	<b>0.127</b>	<b>-0.528</b>	<b>-0.018</b>	<b>0.036</b>
<i>Functional muscle-bone unit</i>	DXA	Arms BMC/Handgrip	0.095	0.122	-0.149	0.339	0.440
		Legs BMC/force	<b>-0.654</b>	<b>0.111</b>	<b>-0.878</b>	<b>-0.431</b>	<b>&lt;0.001</b>
	PQCT	BMC/Handgrip	-0.077	0.153	-0.384	0.231	0.620
		SSIPOL/Handgrip	0.076	0.156	-0.2370	0.3883	0.630
	<i>Tanner dependent z-scores</i>						
<i>Structural muscle-bone unit</i>	DXA	Subtotal BMC/Lean	<b>-0.559</b>	<b>0.169</b>	<b>-0.896</b>	<b>-0.222</b>	<b>0.002</b>
		Arms BMC/Lean	<b>-0.525</b>	<b>0.133</b>	<b>-0.791</b>	<b>-0.258</b>	<b>&lt;0.001</b>
		Legs BMC/Lean	<b>-0.646</b>	<b>0.157</b>	<b>-0.960</b>	<b>-0.331</b>	<b>&lt;0.001</b>
	PQCT	BMC/MCSA	<b>-0.622</b>	<b>0.138</b>	<b>-0.898</b>	<b>-0.346</b>	<b>&lt;0.001</b>
		SSIPOL/MCSA	<b>-0.249</b>	<b>0.106</b>	<b>-0.249</b>	<b>-0.462</b>	<b>0.022</b>
<i>Functional muscle-bone unit</i>	DXA	Arms BMC/Handgrip	0.164	0.132	-0.100	0.429	0.219
		Legs BMC/force	<b>-0.473</b>	<b>0.092</b>	<b>-0.656</b>	<b>-0.289</b>	<b>&lt;0.001</b>
	PQCT	BMC/Handgrip	-0.186	0.140	-0.467	0.096	0.191
		SSIPOL/Handgrip	0.047	0.130	-0.214	0.308	0.719

All the z-scores are sex specific.

Mean dif=Mean difference between the groups (Swimmers vs. CG)

SE=Standard error; CI=Confidence interval

p<0.05