

GABA_B receptors: modulation of thalamocortical dynamics and synaptic plasticity

Maria V. Sanchez-Vives,^{a,b*} Almudena Barbero-Castillo,^a Maria Perez-Zabalza^c and Ramon Reig^d

^a Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), Barcelona, Spain

^b Institució Catalana de Recerca i Estudis Avançats (ICREA), Barcelona, Spain

^c Biomedical Signal Interpretation and Computational Simulation Group, University of Zaragoza, Spain

^d Instituto de Neurociencias de Alicante, Alicante, Spain

Abstract—GABA_B-receptors (GABA_B-Rs) are metabotropic, G protein-coupled receptors for the neurotransmitter GABA. Their activation induces slow inhibitory control of the neuronal excitability mediated by pre- and postsynaptic inhibition. Presynaptically GABA_B-Rs reduce GABA and glutamate release inhibiting presynaptic Ca²⁺ channels in both inhibitory and excitatory synapses while postsynaptic GABA_B-Rs induce robust slow hyperpolarization by the activation of K⁺ channels. GABA_B-Rs are activated by non-synaptic or volume transmission, which requires high levels of GABA release, either by the simultaneous discharge of GABAergic interneurons or very intense discharges in the thalamus or by means of the activation of a neurogliaform interneurons in the cortex. The main receptor subunits GABA_{B1a}, GABA_{B1b} and GABA_{B2} are strongly expressed in neurons and glial cells throughout the central nervous system and GABA_B-R activation is related to many neuronal processes such as the modulation of rhythmic activity in several brain regions. In the thalamus, GABA_B-Rs modulate the generation of the main thalamic rhythm, spindle waves. In the cerebral cortex, GABA_B-Rs also modulate the most prominent emergent oscillatory activity—slow oscillations—as well as faster oscillations like gamma frequency. Further, recent studies evaluating the complexity expressed by the cortical network, a parameter associated with consciousness levels, have found that GABA_B-Rs enhance this complexity, while their blockade decreases it. This review summarizes the current results on how the activation of GABA_B-Rs affects the interchange of information between brain areas by controlling rhythmicity as well as synaptic plasticity. © 2020 IBRO. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Keywords: GABA_B-Rs, metabotropic, perigeniculate, reticularis, cortical complexity, Up states, slow oscillations.

SYNAPTIC TRANSMISSION THROUGH METABOTROPIC GABA_B RECEPTORS IN THALAMUS AND CORTEX

GABAergic neurons are the major inhibitory cell type in the mammalian brain and exhibit a wide variety of morphological and physiological properties. The activation of GABAergic neurons can inhibit postsynaptic target cells through increases in Cl⁻ conductance, mediated by the GABA_A receptor, and increases in K⁺ conductance, mediated by the GABA_B receptor (GABA_B-R). GABA_B-Rs are widely distributed in the central nervous system and have been found at both excitatory and inhibitory neurons (Ulrich and Bettler, 2007) as well as in other types of neurons such as dopaminergic (Boyes and Bolam, 2003) or striatal cholinergic interneurons (Yung et al., 1999; Waldvogel

et al., 2004). In addition, GABA_B-Rs are expressed in glial cells (López-Bendito et al., 2004); in fact most of the neurons in the central nervous system contain GABA_B-Rs (Steiger et al., 2004). GABA_B-Rs are G protein-coupled receptors for the neurotransmitter GABA. Interestingly, GABA_B-Rs are similar in structure to (and in the same receptor family as) metabotropic glutamate receptors (Kaupmann et al., 1997). GABA_B-Rs are expressed pre- and postsynaptically, controlling the release of the neurotransmitter and the excitability of the receiving neurons respectively. Initially, GABA_B-Rs were found to be insensitive to the bicuculline component of inhibitory responses, which was activated by L-baclofen (Bowerly et al., 1980). For many years it was thought that the GABA_B-R family would be found to consist of numerous receptor subtypes, due to the variety in responses obtained with the current agonist and antagonist. However, further investigations demonstrated that there are only two receptor subtypes, GABA_{B1a} and GABA_{B1b}, both of them combined with GABA_{B2} subunits to form heteromeric GABA_{B(1a,2)} and GABA_{B(1b,2)} receptors (Jones

*Corresponding author at: IDIBAPS, Carrer del Rosselló, 149, 08036 Barcelona, Spain.

E-mail address: msanche3@clinic.cat (M. V. Sanchez-Vives).

<https://doi.org/10.1016/j.neuroscience.2020.03.011>

0306-4522/© 2020 IBRO. Published by Elsevier Ltd.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

et al., 1998; Kaupmann et al., 1998; White et al., 1998). These receptor units are produced by a different promoter form by the GABA_B1 gene (Bischoff et al., 1999; Steiger et al., 2004), and they are different in their N-terminal ectodomain by a pair of sushi repeats only in GABA_B1a (Blein et al., 2004). Both isoforms are found in different locations; GABA_B1a are mainly found in presynaptic terminals, where they control neurotransmitter release (Vigot et al., 2006), which depending on the specific type of synapse, excitatory or inhibitory, may inhibit or disinhibit respectively. The presynaptic action of GABA_B-Rs inhibiting glutamate release occurs in many areas of the central nervous system (Willcockson et al., 1984; Dutar and Nicoll, 1988; Huston et al., 1995; Emri et al., 1996; Lin et al., 1996) and most probably in all cortical areas (Deisz and Prince, 1989; Kang, 1995; Ramoa and Sur, 1996; Ziakopoulos et al., 2000). GABA_B-R-mediated inhibition requires the activation of the adenylyl cyclase/protein kinase A second messenger pathway, through the triggering of G α i/o-type G proteins and liberation of G β γ subunits, activating G protein-coupled inward-rectifying K⁺ channels and inhibition of voltage-gated Ca²⁺ chan-

nels (Bettler et al., 2004) (Figure 1). Presynaptic GABA_B-Rs, usually the GABA_B1a subunit, inhibit neurotransmitter release through the inhibition of voltage-gated Ca²⁺ channels (Gassmann et al., 2004). Presynaptic GABA_B-Rs are termed autoreceptors when controlling GABA release, and heteroreceptors when inhibiting other types of neurotransmitter (Figure 1). Often, by means of the activation of presynaptic GABA_B-Rs, GABA decreases glutamate release mainly by blocking voltage-sensitive Ca²⁺ channels and also by the modulation of synaptic vesicle priming (Maguire et al., 1989; Mintz and Bean, 1993; Sakaba and Neher, 2003). Postsynaptically, GABA_B1b are located in the dendritic spines, controlling K⁺ channels that are involved in the postsynaptic inhibition (Pérez-Garci et al., 2006), generating slow inhibitory potentials by the activation of Kir3-type K⁺ channels (Luscher et al., 1997) (Figure 1). GABA_B-Rs also decrease dendritic Ca²⁺ by blocking voltage gate calcium channels in the distal dendrites (Chalifoux and Carter, 2011).

GABA_B-Rs are broadly distributed along the CNS and can generate a large combination of outcomes, making GABA_B-Rs a key element for wide dynamic control of synaptic transmission. Thus, in order to understand the role of GABA_B-Rs in synaptic transmission, it is crucial to identify the specific components underlying the studied circuit, as well as the pre- and postsynaptic location of the GABA_B-Rs. Moreover, in addition to the structural components, the dynamic properties of synaptic transmission should be taken into account, including the different regimens of spontaneous brain activity and the previous history of synaptic activity at the related synapses.

One early idea about the function of GABA_B-Rs was that they controlled slow changes in neuronal excitability. In most cases, the activation of GABA_B autoreceptors needs a strong activation to become active, by means of synchronous activity or repetitive stimulation. This was clearly demonstrated in the thalamus (Kim et al., 1997) and also in the hippocampus (Scanziani, 2000). In line with this evidence, it has been shown that GABA_B-Rs are located distal to the release sites that require GABA volume transmission (Bettler and Tiao, 2006). This is true for most brain areas; however, more recent evidence has demonstrated a very important exception:




Figure 1. Distribution and role of GABA_B-Rs. GABA_B-Rs are located presynaptically on extrasynaptic membranes. Presynaptic GABA_B-Rs prevent neurotransmitter release by down-regulating the activity of voltage-sensitive Ca²⁺ channels or inhibiting the release machinery. GABA_B autoreceptors inhibit the release of GABA, whereas GABA_B heteroreceptors inhibit the release of glutamate and several other neurotransmitters. GABA_B-Rs are also expressed postsynaptically. They induce sIPSCs by activating Kir3-type K⁺ -channels, which hyperpolarizes the membrane, favors voltage-sensitive Mg²⁺ block of NMDA receptors and shunts excitatory currents. GABA_B-Rs in spines and dendrites are expressed to regulate the excitability of the network and to counteract excess excitation during population oscillation and epileptiform activity. Dendritic GABA_B-Rs inhibit back propagating action potentials through activation of K⁺ -channels. During high-frequency transmission GABA depresses its own release by an action on GABA_B autoreceptors, which permits sufficient NMDA receptor activation for the induction of LTP. Reprinted from "Chapter 19: GABA_B Receptor and Absence Epilepsy," by Hua A. Han, Miguel A. Cortez, and O. Carter Snead in *Jasper's Basic Mechanisms of the Epilepsies (4th edn)*, Jeffrey Noebels, Massimo Avoli, Michael Rogawski, Richard Olsen, and Antonio Delgado-Escueta (eds), Oxford University Press: Oxford. Reused with permission.

in the cerebral cortex GABA_B-Rs can be strongly activated by the activity of neurogliaform interneurons. In fact, it was shown that a single action potential of the neurogliaform cells releases enough GABA to generate a slow inhibition mediated by GABA_A and GABA_B receptor activation (Tamas et al., 2003). This type of interneuron targets both dendritic spines of pyramidal neurons (Tamas et al., 2003) and different types of interneurons (Oláh et al., 2007).

Neurogliaform neurons were initially described by Ramon y Cajal (Ramon y Cajal, 1922). They are located in all cortical layers, but those in layer 1 are of special functional relevance in the long-lasting inhibition induced by GABA_B that was demonstrated to elucidate interhemispheric inhibition in mice (Palmer et al., 2012). This type of indirect long-lasting inhibition is used to explain how one hemisphere imposes its dominance during a specific sensory or motor process. The activation of the layer 1 neurons by callosal transmission induced a delayed inhibition by hundreds of milliseconds in the layer 5 pyramidal neurons. This type of inhibition occurred in the distal dendrites, induced by the GABA release activating GABA_{B1b} receptors on the postsynaptic pyramidal neurons (Palmer et al., 2012) and suggests the activation of neurogliaform interneurons induces it. Interestingly, beyond the callosal transmission, this evidence shows that the activation of layer 1 inhibits pyramidal neurons for hundreds of milliseconds. In fact, previous studies demonstrated that the stimulation of distal dendritic regions of pyramidal neurons recruited a GABA_B-R-mediated inhibition (Dutar and Nicoll, 1988; Pérez-Garci et al., 2006). In addition to the synaptic transmission and the sensory processing, this long-lasting inhibition mediated by GABA_B-Rs has been proposed as a mechanism to control brain states. In an elegant *in vitro* study in active cortical slices, Mann and colleagues analyzed how electrical stimulation of layer 1 could trigger a transition from Up to Down states (see below). This interesting result demonstrates that persistent cortical activity can be switched off by the inhibition mediated by GABA_B-R activation (Mann et al., 2009). Additional studies showed the specific role for two types of GABA_B-Rs, GABA_{B1a} and GABA_{B1b}. Using mouse lines of knockout phenotypes for both GABA_B-Rs, Mann et al. demonstrated that the transition to the Down state previously described is only mediated by the activation of the GABA_{B1b} subunit, and not by the GABA_{B1a}. This result strongly suggests a postsynaptic mechanism underlying the modulation of transition states.

Role of GABA_B receptors in synaptic plasticity in the thalamocortical loop

In order to understand the role of GABA_B-Rs in synaptic plasticity, it is crucial to identify the specific components underlying the studied circuit, as well as the pre- and postsynaptic location of the GABA_B-Rs. Moreover, in addition to the structural components, the dynamic properties of synaptic transmission should be taken into account, including the different regimes of spontaneous brain activity and the previous history of synaptic activity at related synapses. This section summarizes the

knowledge of the GABA_B-Rs mediating synaptic transmission and plasticity in thalamus and cortex.

As mentioned above, cortical inhibition mediated by the activation of GABA_B-Rs plays a crucial role regulating neuronal excitability and plasticity. GABA_B-Rs contribute to the regulation of inhibition presynaptically, through the autoinhibition of inhibitory interneurons, and postsynaptically, by means of long-lasting inhibitory postsynaptic potentials (Morrisett et al., 1991; Olpe et al., 1993; Deisz, 1999; Stäubli et al., 1999). Many early studies tried to understand the role of the GABA_B-Rs in synaptic plasticity by pharmacological blockade of GABA_B-Rs, which often generated contradictory results showing either an increase (Olpe and Karlsson, 1990; Olpe et al., 1993; Stäubli et al., 1999) or a reduction of long-term potentiation (Davies et al., 1991; Mott and Lewis, 1991). These apparently conflicting results may be explained by a leading effect of GABA_B-R modulation on either GABA_B-R-mediated inhibitory postsynaptic potentials or autoinhibition of neurotransmission through GABA_A-Rs (Stäubli et al., 1999). In the cerebral cortex, as well as in other brain regions, presynaptic GABA_B-Rs inhibit glutamate release (Kang, 1995; Ramoa and Sur, 1996). Experiments in thalamocortical slices described the effect of the GABA_B-Rs mediating either intracortical or thalamo-cortical excitatory synaptic transmission. The application of the GABA_B-R agonist baclofen reduced the amplitude of intracortical EPSPs, also inducing a slight increase in the paired-pulse ratio, but without affecting thalamocortical transmission (Gil et al., 1997). Similar selective suppression of intrinsic but not afferent fiber synaptic transmission by baclofen was also previously described in the piriform cortex (Tang and Hasselmo, 1994). These results suggest that cortical GABA release selectively suppresses specific pathways by activating presynaptic GABA_B-Rs. This type of input selection mediated by GABA_B-Rs was also demonstrated in the perirhinal cortex. The electrical stimulation in either the entorhinal or temporal cortex induced paired pulse depression (PPD) in the perirhinal cortex, during inter-pulse interval from 100 to 1000 milliseconds. PPD was mediated by GABA spilling over onto excitatory synapses, activating presynaptic GABA_B-Rs in glutamatergic terminals. GABA_B-R-mediated PPD was greater when the temporal cortex was stimulated, suggesting that this pathway has a preferred role controlling the firing and synaptic plasticity of the perirhinal cortex (Ziakopoulos et al., 2000).

As mentioned previously, postsynaptically GABA_B-Rs activate K⁺ channels and also directly inhibit Ca²⁺ mediated dendritic spikes, throughout the dendritic arbor, in cortical and hippocampal pyramidal neurons (Pérez-Garci et al., 2006; Larkum et al., 2007; Chalifoux and Carter, 2011). Ca²⁺ influx via voltage-sensitive Ca²⁺-channels has an important role as an intracellular messenger, activating intracellular signaling cascades that are ultimately responsible for altered synaptic efficacy (Grover and Teyler, 1990; Huang and Malenka, 1993). Therefore, GABA_B-Rs could modulate long-term plasticity by direct action on dendritic spines. *In vivo* experiments also studied the functions of GABA_B-R

blockage on short-term synaptic plasticity. Jia and colleagues described the impact of the blockage of GABA_B-Rs in the thalamocortical visual pathway (Jia et al., 2004). They did so by means of electrical stimulation in the dLGN and recordings in V1 before and after iontophoresis of 2-hydroxy-saclofen. The authors described that both paired pulse and short-term depression were reduced by blocking GABA_B-Rs.

GABA_B-Rs are also involved in the induction of long-term potentiation at visual cortical inhibitory synapses (Komatsu, 1996). Moreover, GABA_B-R activation is necessary for the induction of inhibitory LTP at fast-spiking cell to pyramidal cell synapses, which converts LTP to LTD at convergent excitatory pyramidal cell synapses (Wang and Maffei, 2014). Following these results and given that long-term plasticity changes are related to the deprived eye responses, GABA_B-Rs have also been related to the plasticity of the critical period for ocular dominance in the primary visual cortex of cats. During the critical period, monocular deprivation induces an ocular dominance shift. In juvenile animals, the application of the GABA_B-R agonist enhanced the plasticity facilitating the ocular dominance shift, while the GABA_B-R antagonist impaired it. However, in adult cats the infusion of the same neuromodulators did not induce any significant impact on the ocular dominance shift. These results evidence the role of GABA_B-Rs in juvenile ocular dominance plasticity (Cai et al., 2017).

GABA_B RECEPTORS AND EMERGENT RHYTHMIC PATTERNS: THALAMIC ACTIVITY

Thalamocortical rhythms have a relevant role in the triggering of brain rhythms across different brain states (Llinas and Ribary, 1993; Steriade et al., 1993b). The thalamic reticular nucleus (TRN), and its interaction with the thalamic nuclei, has been proposed as a central modulator of corticothalamic feedback that could modulate sleep states through the modulation of brain rhythms (McCormick and Bal, 1997; Steriade, 2000; Fernandez et al., 2018). The TRN is a nucleus with the shape of a shell around the dorsal thalamus that is formed solely by GABAergic neurons. These neurons are bursty, oscillatory neurons (Kim et al., 1997; Fogerson and Huguenard, 2016) that have large axonal arborizations that profusely innervate thalamic nuclei (Jones, 2012). Furthermore, interlaminar neurons in the visual thalamus are formed by neurons that are equivalent anatomically, electrophysiologically, pharmacologically, anatomically, immunocytochemically, and functionally to the TRN (perigeniculate) neurons (Sanchez-Vives et al., 1996) (Figure 2C). The involvement of the TRN, and thus of inhibitory interneurons, in the control of brain states, is such that GABAergic synaptic transmission has been proposed to be critical for anesthetic drugs to induce unconscious states, by enhancing the effects of TRN activity (Brown et al., 2011). Furthermore, some seizure activity such as absence seizures and its associated spike and wave activity also originate in the thalamus (Hosford et al., 1992; Labate et al., 2005; see below). Alteration of TRN conductances decreases EEG sleep rhythms

(Cueni et al., 2008; Espinosa et al., 2008) and other studies demonstrate that the TRN, and its interaction with thalamocortical neurons, initiates sleep spindles (Figure 2A, B) (Steriade et al., 1987; Von Krosigk et al., 1993; Bazhenov et al., 2000; Halassa et al., 2011; Bartho et al., 2014). Optogenetic studies have shown that TRN may modulate arousal states through selective inhibition of thalamic activity, facilitating the onset of slow waves (Lewis et al., 2015). But, to what extent is the GABAergic action of TRN neurons mediated by GABA_B-Rs? We will refer here to studies in the TRN section around the visual thalamus, which is called the perigeniculate nucleus (PGN), and which maintains the characteristics of TRN. PGN neurons innervate thalamocortical neurons (excitatory) reciprocally (Figure 2C): on their way to the cortex, the axons of the thalamocortical neurons cross the PGN, leaving collaterals at this level. In return, the PGN cells densely innervate the relay cells. In this way, a network of reciprocal interactions between excitatory and inhibitory neurons is established (Kim et al., 1997). When these neurons fire in the bursty mode, this reciprocal innervation results in the generation of spindle waves (Figure 2B) (Von Krosigk et al., 1993). The inhibition deactivates low threshold calcium current (Jahnsen and Llinás, 1984), inducing the rebound burst in thalamocortical cells that results in a cycle of inhibition and bursting that then excites PGN cells (Figure 2A,B), which also burst and further inhibit thalamocortical neurons, which then results in the 7–14 Hz spindle wave (Von Krosigk et al., 1993; McCormick and Bal, 1997). The inhibitory contribution to this cycle relies only on GABA_A-Rs, and the blockade of GABA_B-Rs does not affect spindle waves (Bal et al., 1995a, 1995b).

Thus, when are GABA_B-Rs activated? PGN neurons inhibit not only thalamocortical neurons, but also inhibit each other laterally, both through GABA_A and GABA_B receptors (Figure 3) (Sanchez-Vives et al., 1997). This is equivalent to what happens between TRN neurons (Shu and McCormick, 2002). When PGN neurons are disinhibited by blocking GABA_A receptors, they fire longer bursts of high-frequency action potentials. Under these conditions, it was observed that the inhibition exerted over thalamocortical neurons (Sanchez-Vives and McCormick, 1997) and also over other PGN neurons (Sanchez-Vives et al., 1997) was mediated through GABA_B-Rs (Figure 2B). Given that GABA_B-Rs have a slower time course, the recurrent circuit involving thalamocortical and PGN cells, oscillated at a highly synchronized, slower pace of 3 Hz (Figure 2B). (Sanchez-Vives and McCormick, 1997), a pattern that is highly similar to that generated during absence seizures (Hosford et al., 1992). Therefore, the switch of network activity from relying on GABA_A towards relying on GABA_B-Rs has a large impact on the oscillatory rhythm emerging from the thalamocortical network and the activation of GABA_B-Rs may be particularly important to the generation of some forms of generalized spike-and-wave seizures.

As mentioned above, in order for PGN cells to activate GABA_B-Rs they should be disinhibited in order for them to have longer and more intense firing patterns. This was highly suggestive that the activation of GABA_B-Rs




Figure 2. Pre- and postsynaptic GABA_B-R modulation of thalamic activity patterns. (A) Activation of a burst of action potentials in the PGN cell before and after bath application of bicuculline. (B) Spontaneous generation of a spindle wave is associated with repetitive burst firing in the PGN neuron and the occurrence of repetitive IPSPs in the LGNd cell. The IPSPs are generated by the activity of this PGN cell as well as others. After bath application of bicuculline, the geniculate slice spontaneously generates abnormal oscillations during which the PGN neuron generates prolonged high-frequency burst discharges. Source: Kim, U. et al. (1997) Functional dynamics of GABAergic inhibition in the thalamus. *Science* 278(5335):130–4. Reused with permission. (C) The cell body and dendrites located in the A-A1 interlaminar zone (IZ). The axon densely innervated lamina A1 and also innervated lamina C. One branch of the axon passed through lamina A and into the PGN. Source: Sanchez-Vives et al. (1996). Are the interlaminar zones of the ferret dorsal lateral geniculate nucleus actually part of the perigeniculate nucleus? *Journal of Neuroscience* 16(19): 5923–5941. Reused with permission. (D) (Top) IPSCs recorded in a thalamocortical cell with a microelectrode filled with 2M CsAc and held at 259 mV. Each barrage of IPSCs represents the generation of a spindle wave and was evoked with the local application of glutamate in the PGN. Application of baclofen (100 mM in the micropipette) results in a large reduction in the amplitude of the initial evoked IPSC as well as the generation of spindle wave-associated IPSCs. Local application of CGP 35348 reversed these effects. (Bottom) Expansion of spindle wave-associated IPSCs before and after application of baclofen and after recovery with application of CGP 35348.

required a larger release of GABA, generally accepting that several GABAergic neurons should contribute in order for the extrasynaptic GABA_B-Rs to be reached. However, Kim et al. (1997) demonstrated that the same single PGN neuron could indeed activate both GABA_A and GABA_B, with the difference that the activation of GABA_A and GABA_B required prolonged burst firing. Further, GABA_B-R activation had a delay of 52 ms with respect to the activation of GABA_A-Rs (Kim et al., 1997), which could be due to the properties of G-protein-mediated events that are the intermediaries between receptor binding and K⁺ channel opening or the location of receptors as extrasynaptic receptors (Figure 1) (Mody et al., 1994). Similar features to those described in the thalamus for GABA_B-Rs are described in the hippocampus for GABA_A/GABA_B activation and their impact on rhythmicity (Scanziani, 2000). In more

recent years, it has been proposed that the activation of GABA_B-Rs even contributes to the generation of thalamic spindle oscillations (Ulrich et al., 2018), modulates thalamocortical excitation of cortical inhibitory and excitatory neurons, and participates in the processing of sensory information in the barrel cortex (Porter and Nieves, 2004). Moreover, stimulation of GABA_B-Rs has been found to be effective in cortical seizure-like activity suppression (Chang et al., 2017).

GABA_B-Rs are also present on the presynaptic terminals of both GABAergic neurons (Figure 3E) as well as thalamocortical connections (Figure 2D), and the activation of these receptors results in the reduction of neurotransmitter release (Ulrich and Huguenard, 1996; Sanchez-Vives et al., 1997b). These results suggest that the activation of presynaptic GABA_B-Rs also plays an important role in the regulation of intrathalamic activity.




Figure 3. Lateral GABA_A and GABA_B (pre and postsynaptic) inhibition between inhibitory, perigeniculate neurons in the thalamus. (A) Higher power photomicrograph with Nomarski optics of boutons in the PGN cell. (B) Example of barrage of IPSPs that spontaneously occur in PGN neurons and presumably result from the burst discharge of single PGN neurons. (C) Evoked barrage of IPSPs in a neighboring PGN cell. Each compound IPSP is composed of several presumed unitary IPSPs. (D) Local application of bicuculline (400 mM in micropipette) reduces but does not completely block evoked IPSP. (E) Evoked EPSPs recorded in a PGN cell. The local application of baclofen results in a reduction in the amplitude of these evoked EPSPs. The local application of CGP 34348 reverses this action of baclofen. Source: [Sanchez-Vives et al. \(1997\)](#). Inhibitory interactions between perigeniculate GABAergic neurons. *Journal of Neuroscience* 17(22): 8894-8908. Reused with permission.

GABA_B RECEPTORS AND EMERGENT RHYTHMIC PATTERNS: CORTICAL ACTIVITY

The cortical network can synchronize its activity in different rhythmic patterns ranging from infraslow to ultrafast activity ([Buzsaki and Draguhn, 2004](#)). However, of all these possible frequencies, there is a prominent frequency that dominates the dynamics of the cortical emergent activity during sleep and pathological states: slow wave activity, driven by slow oscillations. Slow oscillations emerge from the recurrent interaction between cortical neurons, making them a network phenomenon ([Steriade et al., 1993c](#)). During the active periods, or Up states, neocortical neurons (both excitatory and inhibitory) are depolarized, receive barrages of synaptic inputs and fire action potentials. During the ensuing Down states, neurons remain hyperpolarized and the synaptic activity is almost silent. Cortical recordings from different species agree on the leading role of infragranular layers—in particular layer 5—in the initiation of Up states ([Sanchez-Vives and McCormick, 2000](#); [Sakata and Harris, 2009](#); [Chauvette et al., 2010](#)). During Up states, the activity expresses coherent oscillations at high frequencies in the beta (15–30 Hz) and gamma (30–90 Hz) range ([Cunningham et al., 2004](#); [Compte et al., 2008](#)). This phenomenon is multiscale such that slow oscillations display similar characteristics whether they are recorded from the intact cortex of a sleeping human or from a small piece of cortex in a plate. It has been proposed that fluctuations of

Up- and Down-states—the origin of the slow waves—is the “default” intrinsic pattern of cortical activity ([Sanchez-Vives and Mattia, 2014](#); [Sanchez-Vives et al., 2017](#)). The cortex defaults to slow-wave activity due to changes in neuromodulation, as in sleep ([Steriade et al., 2001](#); [Riedner et al., 2007](#)) or due to changes in the excitatory/inhibitory balance as in deep anesthesia ([Steriade, Conteras et al., 1993a](#); [Chauvette et al., 2011](#)), or as a consequence of structural lesions, as when a cortical gyrus is physically isolated by cutting the underlying white matter ([Timofeev and Steriade, 1996](#)) or in a cortical slab ([Timofeev et al., 2000](#)).

A key element in the balance and control of either spontaneous emergent or evoked cortical activity is the relationship between excitation and inhibition. Both excitatory and inhibitory neurons fire during Up states as originally described by Steriade et al (1993). Conductance measurements during Up states reveal that the weights of excitation and inhibition are well-balanced *in vivo* ([Haider et al., 2006](#)) and similarly *in vitro* ([Shu et al., 2003](#)) as has been proposed in a cortical computational model ([Compte et al., 2003](#)). Changes in excitatory and inhibitory conductance *in vitro* reveal that both increases and decreases at the beginning/end of Up states occur in close association with each other ([Shu et al., 2003](#); [Zucca et al., 2017](#)). Indeed, the timing of individual excitatory and inhibitory synaptic events also reveals a remarkable coincidence in the accumulation of both excitatory and inhibitory synaptic events during the

rise of an Up state, both *in vitro* and *in vivo*, although it is 1.4 times faster *in vivo* (Compte et al., 2009). A similar coincidence between the timing of excitatory and inhibitory events also occurs at the termination of the Up state. Such interlocking in time of excitation and inhibition is also found in simultaneous recordings of nearby pairs of cortical neurons (Okun and Lampl, 2008). When measuring excitatory versus inhibitory conductances, both have been found to be high at the beginning of the Up state and tend to progressively decrease, but their ratio remains constant and close to 1 in both anesthetized and *in vitro* preparations (Shu et al., 2003; Haider et al., 2006). However, other studies report that the inhibitory conductance is much larger than the excitatory conductance during Up states in natural sleep (Rudolph et al., 2007).

Mechanisms of termination of the Up states

Several mechanisms have been proposed that could account for the termination of Up states. They include: arrival of excitation (Shu et al., 2003; Haider et al., 2006), synaptic depression (Bazhenov et al., 2002, but see Benita et al. 2012), thalamic disfacilitation (Contreras et al., 1996), activation of K^+ currents (Sanchez-Vives and McCormick, 2000; Compte et al., 2003) or extracellular K^+ dynamics (Fröhlich et al., 2006). The time course of the slow afterhyperpolarization observed in intracellular recordings during Down states (e.g. fig. 7C in Contreras et al., 1996; Sanchez-Vives and McCormick, 2000; Sanchez-Vives et al., 2010), sug-

gests that slow K^+ currents can contribute to the termination of Up states and maintenance of Down states. Different mechanisms involving K^+ currents have been proposed, including ATP-dependent K^+ current (Cunningham et al., 2006) and Ca^{2+} and Na^+ -dependent K^+ currents (Sanchez-Vives and McCormick, 2000). Slow K^+ -mediated afterhyperpolarizations are blocked by neurotransmitters (acetylcholine, norepinephrine) that control the transition from sleep to awake states (Schwindt et al., 1988; Brumberg et al., 2000), providing a mechanism for stopping the bistability when entering the awake state (D'Andola et al., 2017).

GABA_B-Rs activation (Parga and Abbott, 2007; Mann et al., 2009; Wang et al., 2010; Craig et al., 2013; Perez-Zabalza et al., 2020) is a plausible mechanism involved in the termination of Up states. More than one of the mentioned mechanisms most probably interact and contribute to the termination of the Up-to-Down state transition. Mann and colleagues analyzed how electrical stimulation of layer 1 could trigger a transition from Up to Down states. This result suggested that persistent cortical activity can be switched off by GABA_B-R-mediated inhibition (Mann et al., 2009) (Figure 4A). Additional studies showed a specific role of the two types of GABA_B-Rs—GABA_{B1a} and GABA_{B1b}. Using mouse lines of knockout phenotypes for both GABA_B-Rs, it was found that the transition to the Down state was mediated by the activation of the GABA_{B1b} subunit, and not by the GABA_{B1a}, thus suggesting a postsynaptic mechanism (Craig et al., 2013). Modelling work has also suggested that GABA_B dynamics have the correct timescale to contribute to this




Figure 4. GABA_B-R modulation of cortical Up and Down states (A) Whole-cell recordings in control (top), 500 nM gabazine (middle), and 1 μM CGP55845 (bottom). Insets show equivalent Up states triggered by stimulation in the same cells. Source: Mann et al. (2009). Distinct roles of GABA_A and GABA_B receptors in balancing and terminating persistent cortical activity. *Journal of Neuroscience*, 29(23), 7513-7518. Reused with permission. **(B)** Raw signal (blue trace), relative firing rate (black trace) and detected Up and Down states (red trace) under control activity (top) and after application of 200 μM CGP 35348 (bottom). **(C)** Raster plots of the relative firing rate are represented for control activity and for 200 μM CGP 35348 corresponding to the ones in (B). **(D) (Top)** Average relative firing rate for Up states during the control and 200 μM CGP 35348 corresponding to the ones in (A). The shadow corresponds to the s.e.m. **(Bottom)** Down-state duration decreases whereas the oscillation frequency increases. Source: Perez-Zabalza et al. (2020) Modulation of cortical slow oscillatory rhythm by GABA_B receptors: an *in vitro* experimental and computational study. *J Physiol*. doi: 10.1111/JP279476. Reused with permission.

Down state termination process (Parga and Abbott, 2007) and that they could interact with firing rate adaptation to modulate the generation of slow oscillations. Indeed, Perez-Zabalza et al. (2020), not only found a significant elongation of Up states when GABA_B-Rs were blocked (Figure 4), but different cortical dynamics during the slow oscillatory regime: subsequent Down states were most commonly (75% cases) elongated (typical) -and thus the oscillatory frequency decreased (Figure 4B, C)-, while the remaining cases were shortened (atypical), therefore affecting the whole oscillatory cycle. Interestingly GABA_B-R blockade also strikingly increased the regularity of the slow oscillation dynamics, revealing that the activation of GABA_B-Rs introduces irregularity, variability and dynamical richness in the spontaneous slow oscillations (Perez-Zabalza et al., 2020). In this study it was also found that the Down-to-Up transition slope, which corresponds to the recruitment of the local network (Reig et al., 2010; Sanchez-Vives et al., 2010), was not affected by GABA_B-R blockade. Interestingly, the Up-to-Down (downward) transition slope was significantly decreased, further supporting the role of GABA_B-Rs in the termination of Up states.

GABA_B-Modulation Of Cortical Complexity

Throughout different brain states, cortical units interact in different ways resulting in a variety of cortical dynamics that are accompanied by switches of behavioral states (Gervasoni et al., 2004). Such cortical dynamics result from reciprocal interactions between excitatory and inhibitory cortical units, depolarizing simultaneously, but the most active inhibitory populations vary during different brain states (Gentet et al., 2010). During awake states, cortical activity is characterized by complex patterns of causal interactions, whereas this complexity collapses in deep sleep or anesthesia (Steriade, 2006). The study of the mechanisms that modulate the emergence of complex brain states is critical to the development of new methodologies to detect brain state and consciousness levels (i.e., during anesthesia or in brain-injured patients) and promote state transitions and recovery of function. The study of cortical complexity can be approached from the paradigm of the balance between segregation and integration of cortical activity, defined as brain complexity (Tononi, 2004). Different methods have been used to quantify brain complexity in humans. The Perturbational




Figure 5. GABA_B-R modulation of cortical complexity. Brain complexity measured in human using PCI, can be studied in isolated cortical slices *in vitro*. (A) Average of EEG responses following TMS (Red) and maximum current sources (color-coded according to their activation latency: light blue, 0 ms; red, 300 ms). Yellow cross: TMS target on the cortex. Source: Sanchez-Vives et al. (2017). Shaping the default activity pattern of the cortical network. *Neuron* 94:993–1001. Reused with permission. Originally modified from Pigorini et al. (2015) *Neuroimage* 112, 105–113. (B) Spontaneous emerging activity in isolated cortical slices, (top) Local Field Potential (LFP) and (bottom) Multiunit activity (MUA) under bath application of 50 μM NE and 0.5 μM CCh (red) and slow wave activity (SWA, blue). PCI was calculated using Lempel-Ziv compression of the significant MUA (SS(x,t)) contained in a binary matrix. Source: D’Andola et al. (2017). Bistability, causality, and complexity in cortical networks: an *in vitro* perturbational study. *Cereb Cortex* 91:1–10. Reused with permission. (C) Example of decay in sPCI in awake-like state of a cortical slice and during blockade of slow inhibition (CGP 1 μM). (D) Example of decay in sPCI reduction in slow oscillatory activity in a cortical slice and during blockade of slow inhibition (CGP 1 μM) Binary matrix SS(x,t) of MUA significantly higher than a threshold activity.

Complexity Index (PCI) is one such method, which determines brain complexity based on the ability of the network to integrate and differentiate the responses and cortical interaction following a particular perturbation (i.e., electrical stimulation) (Casali et al., 2013). Casali et al. (2013) measured brain complexity based on the spatiotemporal distribution of significant electroencephalography (EEG) responses following transcranial magnetic stimulation (TMS). They demonstrated that during NREM sleep, TMS induced bimodal stereotypical EEG responses (Figure 1A), while during wakefulness, TMS triggered complex heterogeneous EEG responses (Figure 1A) (Casali et al., 2013).

The measurement of complexity using PCI in humans can distinguish different levels of consciousness but does not provide information about the mechanisms involved. To provide a mechanistic approach, D'Andola et al., (2017) adapted the PCI measure to the cerebral cortex *in vitro* and found that the measure was able to distinguish variations in cortical complexity for different simulated “brain states” reached by varying neurotransmitter concentrations. The authors found that network complexity was low during slow wave activity, and increased at induced awake-like states by cholinergic and noradrenergic agonists (Figure 5B) (D'Andola et al., 2017). Indeed, it was also demonstrated that simply increasing neuronal excitability and firing rates do not imply higher complexity states measured by perturbational methods (D'Andola et al., 2017; Barbero-Castillo et al., 2020). Thus, while network excitability and neuronal firing patterns are key elements for the normal functioning of cortical networks and for the transition between different brain states, there is not a linear relationship between cortical excitability and complexity. For example, blocking of GABA_A-Rs can lead into rather high excitability and epileptiform discharges; however, cortical complexity decreases in these cases (Barbero-Castillo et al., 2020). Decreasing inhibition by blocking GABA_B-Rs was found to decrease cortical complexity independently from firing, either departing from an awake-like, asynchronous state (Figure 5C) or from a slow wave activity state (Figure 5D) (Barbero-Castillo et al., 2020), in agreement with the idea that the activation of GABA_B-Rs brings in dynamical richness and irregularity to the cortical network (Perez-Zabalza et al., 2020).

INDICATED TO MATCH THE OUTLINE

- 1 Synaptic transmission through metabotropic GABA_B receptors in thalamus and cortex
 - i Role of GABA_B receptors in synaptic plasticity in the thalamocortical loop
- 2 GABA_B receptors and emergent rhythmic patterns: thalamic activity
- 3 GABA_B receptors and emergent rhythmic patterns: cortical activity
 - i Mechanisms of termination of Up states
 - ii GABA_B modulation of cortical complexity

ACKNOWLEDGEMENTS

This work was supported by EU H2020 Research and Innovation Programme, Grant 785907 (HBP SGA2), BFU2017-85048-R (MINECO) and SloW-Dyn FLAGERA-PCIN-2015-162-C02-01. We would like to thank Tony Donegan for editing the paper.

REFERENCES

- Bal T, von Krosigk M, McCormick DA (1995a) Synaptic and membrane mechanisms underlying synchronized oscillations in the ferret lateral geniculate nucleus *in vitro*. *J Physiol* 483:641–663.
- Bal T, von Krosigk M, McCormick DA (1995b) Role of the ferret perigeniculate nucleus in the generation of synchronized oscillations *in vitro*. *J Physiol* 483:665–685.
- Barbero-Castillo A, Mateos-Aparicio P, Dalla Porta L, Camassa A, Perez Mendez L, Sanchez-Vives MV (2020). Impact of GABAA and GABAB inhibition on cortical dynamics and perturbational complexity during synchronous and asynchronous activity. <https://doi.org/10.5281/zenodo.3856665>.
- Bartho P, Slezia A, Matyas F, Faradz-Zade L, Ulbert I, Harris KD, Acsady L (2014) Ongoing network state controls the length of sleep spindles via inhibitory activity. *Neuron* 82:1367–1379.
- Bazhenov M, Timofeev I, Steriade M, Sejnowski T (2000) Spiking-bursting activity in the thalamic reticular nucleus initiates sequences of spindle oscillations in thalamic networks. *J Neurophysiol* 84:1076–1087.
- Bazhenov M, Timofeev I, Steriade M, Sejnowski TJ (2002) Model of thalamocortical slow-wave sleep oscillations and transitions to activated States. *J Neurosci* 22:8691–8704.
- Benita JM, Guillaumon A, Deco G, Sanchez-Vives MVM (2012) Synaptic depression and slow oscillatory activity in a biophysical network model of the cerebral cortex. *Frontiers in computational neuroscience* 6:64.
- Bettler B, Kaupmann K, Mosbacher J, Gassmann M (2004) Molecular structure and physiological functions of GABA(B) receptors. *Physiol Rev* 84:835–867.
- Bettler B, Tiao JY-H (2006) Molecular diversity, trafficking and subcellular localization of GABAB receptors. *Pharmacol Ther* 110:533–543.
- Bischoff S, Leonhard S, Reymann N, Schuler V, Shigemoto R, Kaupmann K, Bettler B (1999) Spatial distribution of GABA(B)R1 receptor mRNA and binding sites in the rat brain. *J Comp Neurol* 412:1–16.
- Blein S, Gingham R, Uhrin D, Smith BO, Soares DC, Vettel S, McIlhinney RAJ, White JH, Barlow PN (2004) Structural analysis of the complement control protein (CCP) modules of GABA(B) receptor 1a: only one of the two CCP modules is compactly folded. *J Biol Chem* 279:48292–48306.
- Bowery NG, Hill DR, Hudson AL, Doble A, Middlemiss DN, Shaw J, Turnbull M (1980) (-)Baclofen decreases neurotransmitter release in the mammalian CNS by an action at a novel GABA receptor. *Nature* 283:92–94.
- Boyes J, Bolam JP (2003) The subcellular localization of GABA(B) receptor subunits in the rat substantia nigra. *Eur J Neurosci* 18:3279–3293.
- Brown EN, Purdon PL, Van Dort CJ (2011) General Anesthesia and Altered States of Arousal: A Systems Neuroscience Analysis. *Annu Rev Neurosci* 34:601–628.
- Brumberg JC, Nowak LG, McCormick DA (2000) Ionic mechanisms underlying repetitive high-frequency burst firing in supragranular cortical neurons. *J Neurosci* 20:4829–4843.
- Buzsaki G, Draguhn A (2004) Neuronal oscillations in cortical networks. *Science* 304:1926–1929.
- Cai S, Fischer QS, He Y, Zhang L, Liu H, Daw NW, Yang Y (2017) GABAB receptor-dependent bidirectional regulation of critical

- period ocular dominance plasticity in cats. *PLoS One* 12:e0180162.
- Casali AG, Gosseries O, Rosanova M, Boly M, Sarasso S, Casali KR, Casarotto S, Bruno M-A, Laureys S, Tononi G, Massimini M (2013) A theoretically based index of consciousness independent of sensory processing and behavior. *Sci Transl Med* 5:198ra105.
- Chalifoux JR, Carter AG (2011) GABAB receptor modulation of voltage-sensitive calcium channels in spines and dendrites. *J Neurosci* 31:4221–4232.
- Chang WJ, Chang WP, Shyu BC (2017) Suppression of cortical seizures by optic stimulation of the reticular thalamus in PV-mChR2-YFP BAC transgenic mice. *Mol Brain* 10:42.
- Chauvette S, Crochet S, Volgushev M, Timofeev I (2011) Properties of Slow Oscillation during Slow-Wave Sleep and Anesthesia in Cats. *J Neurosci* 31:14998–15008.
- Chauvette S, Volgushev M, Timofeev I (2010) Origin of active states in local neocortical networks during slow sleep oscillation. *Cereb Cortex* 20:2660–2674.
- Compte A, Reig R, Descalzo VF, Harvey MA, Puccini GD, Sanchez-Vives MV (2008) Spontaneous high-frequency (10–80 Hz) oscillations during up states in the cerebral cortex in vitro. *J Neurosci* 28:13828–13844.
- Compte A, Reig R, Sanchez-Vives MV (2009) Timing excitation and inhibition in the cortical network. In: *Coherent Behavior in Neuronal Networks*. New York, NY: Springer. p. 17–46.
- Compte A, Sanchez-Vives MV, McCormick DA, Wang X-J (2003) Cellular and network mechanisms of slow oscillatory activity (. *J Neurophysiol* 89:2707–2725.
- Contreras D, Timofeev I, Steriade M (1996) Mechanisms of long-lasting hyperpolarizations underlying slow sleep oscillations in cat corticothalamic networks. *J Physiol* 494:251–264.
- Craig MT, Mayne EW, Bettler B, Paulsen O, McBain CJ (2013) Distinct roles of GABAB_{1a}- and {GABAB_{1b}-containing {GABAB} receptors in spontaneous and evoked termination of persistent cortical activity. *J Physiol* 591:835–843.
- Cueni L, Canepari M, Luján R, Emmenegger Y, Watanabe M, Bond CT, Franken P, Adelman JP, Lüthi A (2008) T-type Ca²⁺ channels, SK2 channels and SERCAs gate sleep-related oscillations in thalamic dendrites. *Nat Neurosci* 11:683–692.
- Cunningham MO, Pervouchine DD, Racca C, Kopell NJ, Davies CH, Jones RSG, Traub RD, Whittington MA (2006) Neuronal metabolism governs cortical network response state. *Proc Natl Acad Sci* 103:5597–5601.
- Cunningham MO, Whittington MA, Bibbig A, Roopun A, LeBeau FEN, Vogt A, Monyer H, Buhl EH, Traub RD (2004) A role for fast rhythmic bursting neurons in cortical gamma oscillations in vitro. *Proc Natl Acad Sci* 101:7152–7157.
- D'Andola M, Rebollo B, Casali AG, Weinert JF, Pigorini A, Villa R, Massimini M, Sanchez-Vives MV (2017) Bistability, Causality, and Complexity in Cortical Networks: An In Vitro Perturbational Study. *Cereb Cortex* 91:1–10.
- Davies CH, Starkey SJ, Pozza MF, Collingridge GL (1991) GABAB autoreceptors regulate the induction of LTP. *Nature* 349:609–611.
- Deisz RA (1999) The GABA(B) receptor antagonist CGP 55845A reduces presynaptic GABA(B) actions in neocortical neurons of the rat in vitro. *Neuroscience* 93:1241–1249.
- Deisz RA, Prince DA (1989) Frequency-dependent depression of inhibition in guinea-pig neocortex in vitro by GABAB receptor feed-back on GABA release. *J Physiol* 412:513–541.
- Dutar P, Nicoll RA (1988) A physiological role for GABAB receptors in the central nervous system. *Nature* 332:156–158.
- Emri Z, Turner JP, Crunelli V (1996) Tonic activation of presynaptic GABA(B) receptors on thalamic sensory afferents. *Neuroscience* 72:689–698.
- Espinosa F, Torres-Vega MA, Marks GA, Joho RH (2008) Ablation of Kv3.1 and Kv3.3 potassium channels disrupts thalamocortical oscillations in vitro and in vivo. *J Neurosci* 28:5570–5581.
- Fernandez LM, Vantomme G, Osorio-Forero A, Cardis R, Béard E, Lüthi A (2018) Thalamic reticular control of local sleep in mouse sensory cortex. *eLife* 2018;7.
- Fogerson PM, Huguenard JR (2016) Tapping the Brakes: Cellular and Synaptic Mechanisms that Regulate Thalamic Oscillations. *Neuron* 92:687–704.
- Fröhlich F, Bazhenov M, Timofeev I, Steriade M, Sejnowski TJ (2006) Slow state transitions of sustained neural oscillations by activity-dependent modulation of intrinsic excitability. *J Neurosci* 26:6153–6162.
- Gassmann M et al (2004) Comparative cellular distribution of GABAA and GABAB receptors in the human basal ganglia: immunohistochemical colocalization of the alpha 1 subunit of the GABAA receptor, and the GABABR1 and GABABR2 receptor subunits. *Nature* 396:589–601.
- Genet LJ, Avermann M, Matyas F, Staiger JF, Petersen CCH (2010) Membrane Potential Dynamics of GABAergic Neurons in the Barrel Cortex of Behaving Mice. *Neuron* 65:422–435.
- Gervasoni D, Lin S-C, Ribeiro S, Soares ES, Pantoja J, Nicolesis MAL (2004) Global forebrain dynamics predict rat behavioral states and their transitions. *J Neurosci* 24:11137–11147.
- Gil Z, Connors BW, Amitai Y (1997) Differential regulation of neocortical synapses by neuromodulators and activity. *Neuron* 19:679–686.
- Grover LM, Teyler TJ (1990) Two components of long-term potentiation induced by different patterns of afferent activation. *Nature* 347:477–479.
- Haider B, Duque A, Hasenstaub AR, McCormick DA (2006) Neocortical network activity in vivo is generated through a dynamic balance of excitation and inhibition. *J Neurosci* 26:4535–4545.
- Halassa MM, Siegle JH, Ritt JT, Ting JT, Feng G, Moore CI (2011) Selective optical drive of thalamic reticular nucleus generates thalamic bursts and cortical spindles. *Nat Neurosci* 14:1118–1120.
- Hosford DA, Clark S, Cao Z, Wilson WA, Lin FH, Morrisett RA, Huin A (1992) The role of GABAB receptor activation in absence seizures of lethargic (lh/lh) mice. *Science* 257:398–401.
- Huang YY, Malenka RC (1993) Examination of TEA-induced synaptic enhancement in area CA1 of the hippocampus: the role of voltage-dependent Ca²⁺ channels in the induction of LTP. *J Neurosci* 13:568–576.
- Huston E, Cullen GP, Burley JR, Dolphin AC (1995) The involvement of multiple calcium channel sub-types in glutamate release from cerebellar granule cells and its modulation by GABAB receptor activation. *Neuroscience* 68:465–478.
- Jahnsen H, Llinás R (1984) Ionic basis for the electro-responsiveness and oscillatory properties of guinea-pig thalamic neurones in vitro. *J Physiol* 349:227–247.
- Jia F, Xie X, Zhou Y (2004) Short-term depression of synaptic transmission from rat lateral geniculate nucleus to primary visual cortex in vivo. *Brain Res* 1002:158–161.
- Jones E (2012) *The Thalamus*. US: Springer.
- Jones KA et al (1998) GABA(B) receptors function as a heteromeric assembly of the subunits GABA(B)R1 and GABA(B)R2. *Nature* 396:674–679.
- Kang Y (1995) Differential paired pulse depression of non-NMDA and NMDA currents in pyramidal cells of the rat frontal cortex. *J Neurosci* 15:8268–8280.
- Kaupmann K, Huggel K, Heid J, Flor PJ, Bischoff S, Mickel SJ, McMaster G, Angst C, Bittiger H, Froestl W, Bettler B (1997) Expression cloning of GABA(B) receptors uncovers similarity to metabotropic glutamate receptors. *Nature* 386:239–246.
- Kaupmann K, Malitschek B, Schuler V, Heid J, Froestl W, Beck P, Mosbacher J, Bischoff S, Kulik A, Shigemoto R, Karschin A, Bettler B (1998) GABA(B)-receptor subtypes assemble into functional heteromeric complexes. *Nature* 396:683–687.
- Kim U, Sanchez-Vives MV, McCormick DA (1997) Functional dynamics of GABAergic inhibition in the thalamus. *Science* (80-) 278:130–134.
- Komatsu Y (1996) GABAB receptors, monoamine receptors, and postsynaptic inositol trisphosphate-induced Ca²⁺ release are involved in the induction of long-term potentiation at visual cortical inhibitory synapses. *J Neurosci* 16:6342–6352.

- Labate A, Briellmann RS, Abbott DF, Waites AB, Jackson GD (2005) Typical childhood absence seizures are associated with thalamic activation. *Epileptic Disord* 7:373–377.
- Larkum ME, Waters J, Sakmann B, Helmchen F (2007) Dendritic spikes in apical dendrites of neocortical layer 2/3 pyramidal neurons. *J Neurosci* 27:8999–9008.
- Lewis LD, Voigts J, Flores FJ, Schmitt LI, Wilson MA, Halassa MM, Brown EN (2015) Thalamic reticular nucleus induces fast and local modulation of arousal state. *Elife* 4 e08760.
- Lin Q, Peng YB, Willis WD (1996) Role of GABA receptor subtypes in inhibition of primate spinothalamic tract neurons: difference between spinal and periaqueductal gray inhibition. *J Neurophysiol* 75:109–123.
- Llinas R, Ribary U (1993) Coherent 40-Hz oscillation characterizes dream state in humans. *Proc Natl Acad Sci U S A* 90:2078–2081.
- López-Bendito G, Shigemoto R, Kulik A, Vida I, Fairén A, Luján R (2004) Distribution of metabotropic GABA receptor subunits GABA_{B1a/b} and GABA_{B2} in the rat hippocampus during prenatal and postnatal development. *Hippocampus* 14:836–848.
- Luscher C, Jan LY, Stoffel M, Malenka RC, Nicoll RA (1997) G protein-coupled inwardly rectifying K⁺ channels (GIRKs) mediate postsynaptic but not presynaptic transmitter actions in hippocampal neurons. *Neuron* 19:687–695.
- Maguire G, Maple B, Lukasiewicz P, Werblin F (1989) Gamma-aminobutyrate type B receptor modulation of L-type calcium channel current at bipolar cell terminals in the retina of the tiger salamander. *Proc Natl Acad Sci U S A* 86:10144–10147.
- Mann EO, Kohl MM, Paulsen O (2009) Distinct roles of GABA(A) and GABA(B) receptors in balancing and terminating persistent cortical activity. *J Neurosci* 29:7513–7518.
- McCormick DA, Bal T (1997) Sleep and arousal: thalamocortical mechanisms. *Annu Rev Neurosci* 20:185–215.
- Mintz IM, Bean BP (1993) GABAB receptor inhibition of P-type Ca²⁺ channels in central neurons. *Neuron* 10:889–898.
- Mody I, De Koninck Y, Otis TS, Soltesz I (1994) Bridging the cleft at GABA synapses in the brain. *Trends Neurosci* 17:517–525.
- Morrisett RA, Mott DD, Lewis DV, Swartzwelder HS, Wilson WA (1991) GABA(B)-receptor-mediated inhibition of the N-methyl-D-aspartate component of synaptic transmission in the rat hippocampus. *J Neurosci* 11:203–209.
- Mott DD, Lewis DV (1991) Facilitation of the induction of long-term potentiation by GABAB receptors. *Science* (80-) 252:1718–1720.
- Okun M, Lampl I (2008) Instantaneous correlation of excitation and inhibition during ongoing and sensory-evoked activities. *Nat Neurosci* 11:535–537.
- Oláh S, Komlósi G, Szabadics J, Varga C, Tóth E, Barzó P, Tamás G (2007) Output of neurogliaform cells to various neuron types in the human and rat cerebral cortex. *Front Neural Circuits* 1:4.
- Olpe HR, Karlsson G (1990) The effects of baclofen and two GABAB-receptor antagonists on long-term potentiation. *Naunyn Schmiedebergs Arch Pharmacol* 342:194–197.
- Olpe HR, Wörner W, Ferrat T (1993) Stimulation parameters determine role of GABAB receptors in long-term potentiation. *Experientia* 49:542–546.
- Palmer LM, Schulz JM, Murphy SC, Ledergerber D, Murayama M, Larkum ME (2012) The cellular basis of GABA(B)-mediated interhemispheric inhibition. *Science* 335:989–993.
- Parga N, Abbott LF (2007) Network model of spontaneous activity exhibiting synchronous transitions between up and down States. *Front Neurosci* 1:57–66.
- Pérez-García E, Gassmann M, Bettler B, Larkum ME (2006) The GABAB1b isoform mediates long-lasting inhibition of dendritic Ca²⁺ spikes in layer 5 somatosensory pyramidal neurons. *Neuron* 50:603–doi: 10.5281/zenodo.3856665.
- Perez-Zabalza M, Reig R, Manrique J, Jercog D, Winograd M, Parga N, Sanchez-Vives MV (2020) Modulation of cortical slow oscillatory rhythm by GABAB receptors: an experimental and computational study. *J Physiol*. <https://doi.org/10.1113/JP279476>.
- Porter JT, Nieves D (2004) Presynaptic GABAB receptors modulate thalamic excitation of inhibitory and excitatory neurons in the mouse barrel cortex. *J Neurophysiol* 92:2762–2770.
- Sarasso A, Pigorini S, Proserpio P, Szymanski C, Arnulfo G, Casarotto S, Fecchio M, Rosanova M, Mariotti M, Lo Russo G, et al. (2015) Bistability breaks-off deterministic responses to intracortical stimulation during non- REM sleep. *Neuroimage* 112:105–113.
- Ramoá AS, Sur M (1996) Short-term synaptic plasticity in the visual cortex during development. *Cereb Cortex* 6:640–646.
- Ramon y Cajal S, (1922). Estudios sobre la fina estructura de la corteza regional de los roedores I. Corteza suboccipital (retroesplenial de Brodmann). *Trab del Lab Inv bio XX:fasc. 1*.
- Reig R, Mattia M, Compte A, Belmonte C, Sanchez-Vives MV (2010) Temperature Modulation of Slow and Fast Cortical Rhythms. *J Neurophysiol* 103:1253–1261.
- Riedner BA, Vyazovskiy VV, Huber R, Massimini M, Esser S, Murphy M, Tononi G (2007) Sleep homeostasis and cortical synchronization: III. A high-density EEG study of sleep slow waves in humans. *Sleep* 30:1643–1657.
- Rudolph M, Pospischil M, Timofeev I, Destexhe A (2007) Inhibition Determines Membrane Potential Dynamics and Controls Action Potential Generation in Awake and Sleeping Cat Cortex. *J Neurosci* 27:5280–5290.
- Sakaba T, Neher E (2003) Direct modulation of synaptic vesicle priming by GABA(B) receptor activation at a glutamatergic synapse. *Nature* 424:775–778.
- Sakata S, Harris KD (2009) Laminar Structure of Spontaneous and Sensory-Evoked Population Activity in Auditory Cortex. *Neuron* 64:404–418.
- Sanchez-Vives MV, Massimini M, Mattia M (2017) Shaping the Default Activity Pattern of the Cortical Network. *Neuron* 94:993–1001.
- Sanchez-Vives MV, Mattia M (2014) Slow wave activity as the default mode of the cerebral cortex. *Arch Ital Biol* 152:2–3.
- Sanchez-Vives MV, McCormick DA (1997) Functional properties of perigeniculate inhibition of dorsal lateral geniculate nucleus thalamocortical neurons in vitro. *J Neurosci* 17:8880–8893.
- Sanchez-Vives MV, McCormick DA (2000) Cellular and network mechanisms of rhythmic recurrent activity in neocortex. *Nat Neurosci* 3:1027–1034.
- Sanchez-Vives MV, Bal T, Kim U, von Krosigk M, McCormick DA (1996) Are the interlaminar zones of the ferret dorsal lateral geniculate nucleus actually part of the perigeniculate nucleus? *J Neurosci* 16:5923–5941.
- Sanchez-Vives MV, Bal T, McCormick DA (1997) Inhibitory interactions between perigeniculate GABAergic neurons. *J Neurosci* 17:8894–8908.
- Sanchez-Vives MV, Mattia M, Compte A, Perez-Zabalza M, Winograd M, Descalzo VF, Reig R (2010) Inhibitory modulation of cortical up states. *J Neurophysiol* 104:1314–1324.
- Scanziani M (2000) GABA spillover activates postsynaptic GABA(B) receptors to control rhythmic hippocampal activity. *Neuron* 25:673–681.
- Schwindt PC, Spain WJ, Foehring RC, Chubb MC, Crill WE (1988) Slow conductances in neurons from cat sensorimotor cortex in vitro and their role in slow excitability changes. *J Neurophysiol* 59:450–467.
- Shu Y, Hasenstaub A, McCormick DA (2003) Turning on and off recurrent balanced cortical activity. *Nature* 423:288–293.
- Shu Y, McCormick DA (2002) Inhibitory interactions between ferret thalamic reticular neurons. *J Neurophysiol* 87:2571–2576.
- Stäubli U, Scafidi J, Chun D (1999) GABA(B) receptor antagonism: Facilitatory effects on memory parallel those on LTP induced by TBS but not HFS. *J Neurosci* 19:4609–4615.
- Steiger JL, Bandyopadhyay S, Farb DH, Russek SJ (2004) cAMP response element-binding protein, activating transcription factor-4, and upstream stimulatory factor differentially control hippocampal GABABR1a and GABABR1b subunit gene expression through alternative promoters. *J Neurosci* 24:6115–6126.
- Steriade M (2000) Corticothalamic resonance, states of vigilance and mentation. *Neuroscience* 101:243–276.
- Steriade M (2006) Grouping of brain rhythms in corticothalamic systems. *Neuroscience* 137:1087–1106.

- Steriade M, Contreras D, Dossi RC, Nunez A (1993a) The slow (< 1 Hz) oscillation in reticular thalamic and thalamocortical neurons: scenario of sleep rhythm generation in interacting thalamic and neocortical networks. *Journal of Neuroscience* 13(8):3284–3299.
- Steriade M, Domich L, Oakson G, Deschenes M (1987) The deafferented reticular thalamic nucleus generates spindle rhythmicity. *J Neurophysiol* 57:260–273.
- Steriade M, McCormick DA, Sejnowski TJ (1993b) Thalamocortical oscillations in the sleeping and aroused brain. *Science* 262:679–685.
- Steriade M, Nuñez A, Amzica F (1993c) A novel slow (< 1 Hz) oscillation of neocortical neurons in vivo: depolarizing and hyperpolarizing components. *J Neurosci* 13:3252–3265.
- Steriade M, Timofeev I, Grenier F (2001) Natural waking and sleep states: a view from inside neocortical neurons. *J Neurophysiol* 85:1969–1985.
- Tamas G, Lorincz A, Simon A, Szabadics J (2003) Identified sources and targets of slow inhibition in the neocortex. *Science* 299:1902–1905.
- Tang AC, Hasselmo ME (1994) Selective suppression of intrinsic but not afferent fiber synaptic transmission by baclofen in the piriform (olfactory) cortex. *Brain Res* 659:75–81.
- Timofeev I, Grenier F, Bazhenov M, Sejnowski TJ, Steriade M (2000) Origin of slow cortical oscillations in deafferented cortical slabs. *Cereb Cortex* 10:1185–1199.
- Timofeev I, Steriade M (1996) Low-frequency rhythms in the thalamus of intact-cortex and decorticated cats. *J Neurophysiol* 76:4152–4168.
- Tononi G (2004) An information integration theory of consciousness. *BMC Neurosci* 5:42.
- Ulrich D, Bettler B (2007) GABA(B) receptors: synaptic functions and mechanisms of diversity. *Curr Opin Neurobiol* 17:298–303.
- Ulrich D, Huguenard JR (1996) GABAB receptor-mediated responses in GABAergic projection neurones of rat nucleus reticularis thalami in vitro. *J Physiol* 493(Pt 3):845–854.
- Ulrich D, Lalanne T, Gassmann M, Bettler B (2018) GABAB receptor subtypes differentially regulate thalamic spindle oscillations. *Neuropharma*. 136:106–116.
- Vigot R, Barbieri S, Brauner-Osborne H, Turecek R, Shigemoto R, Zhang Y-P, Lujan R, Jacobson LH, Biermann B, Fritschy J-M, Vacher C-M, Muller M, Sansig G, Guetg N, Cryan JF, Kaupmann K, Gassmann M, Oertner TG, Bettler B (2006) Differential compartmentalization and distinct functions of GABAB receptor variants. *Neuron* 50:589–601.
- Von Krosigk M, Bal T, McCormick DA (1993) Cellular mechanisms of a synchronized oscillation in the thalamus. *Science* (80-) 261:361–364.
- Waldvogel HJ, Billinton A, White JH, Emson PC, Faull RLM (2004) Comparative Cellular Distribution of GABA and GABAB Receptors in the Human Basal Ganglia: Immunohistochemical Colocalization of the $\alpha 1$ Subunit of the GABA A Receptor, and the GABABR1 and GABABR2 Receptor Subunits. *J Comp Neurol* 470:339–356.
- Wang L, Maffei A (2014) Inhibitory plasticity dictates the sign of plasticity at excitatory synapses. *J Neurosci* 34:1083–1093.
- Wang Y, Neubauer FB, Lüscher H-R, Thurley K (2010) GABAB receptor-dependent modulation of network activity in the rat prefrontal cortex in vitro. *Eur J Neurosci*:no–no..
- White JH, Wise A, Main MJ, Green A, Fraser NJ, Disney GH, Barnes AA, Emson P, Foord SM, Marshall FH (1998) Heterodimerization is required for the formation of a functional GABA(B) receptor. *Nature* 396:679–682.
- Willcockson WS, Chung JM, Hori Y, Lee KH, Willis WD (1984) Effects of iontophoretically released amino acids and amines on primate spinothalamic tract cells. *J Neurosci* 4:732–740.
- Yung KK, Ng TK, Wong CK (1999) Subpopulations of neurons in the rat neostriatum display GABABR1 receptor immunoreactivity. *Brain Res* 830:345–352.
- Ziakopoulos Z, Brown MW, Bashir ZI (2000) GABAB receptors mediate frequency-dependent depression of excitatory potentials in rat perirhinal cortex in vitro. *Eur J Neurosci* 12:803–809.
- Zucca S, D'Urso G, Pasquale V, Vecchia D, Pica G, Bovetti S, Moretti C, Varani S, Molano-Mazón M, Chiappalone M, Panzeri S, Panzeri S, Fellin T (2017) An inhibitory gate for state transition in cortex. *Elife* 6 e26177.

(Received 4 December 2019, Accepted 6 March 2020)
(Available online xxxx)